

DAFTAR PUSTAKA

- Coglianesi, E., Larson, M.G., Vasan, R., Ho, J., Ghorbani, A., McCabe, E., Cheng, S., Fradley, M., Kretschman, D., Gao, W., O'Connor, G., Wang, T., dan Januzzi, J. 2013. Distribution and Clinical Correlates of the Interleukin Receptor Family Member Soluble ST2 in the Framingham Heart Study. *Clin Chem*, vol. 58, no. 12, hh. 1673 – 1681.
- Conaway, D., O'Keefe, J., Reid, K. dan Spertus, J. 2005. Frequency of Undiagnosed Diabetes Mellitus in Patients With Acute Coronary Syndrome. *The American Journal of Cardiology*, vol. 96, no.3, hh.363-365.
- Davies, M. 2000. Coronary Disease: The pathophysiology of acute coronary syndromes. *Heart*, vol. 83, no.3, pp.361-366.
- Deedwania, P., Kosiborod, M., Barrett, E., Ceriello, A., Isley, W., Mazzone, T. dan Raskin, P. 2008. Hyperglycemia and Acute Coronary Syndrome. *Anesthesiology*, vol. 109, no.1, hh.14-24.
- Duka, A., Zaimi, E., Hasimi, E., Kristo, A., Seiti, J. dan Petrela, E. 2015. Hyperglycemia, Diabetes Mellitus and the Outcomes of STEMI Patients. *International Journal of Science and Research*, vol.5, no.12, hh.1344-1348.
- Farcas, A.D., Anton, F.P., Goidescu, C.M., Gavrila, I.L., Vida-Simiti, L.A., Stoia, M.A. 2017. Serum Soluble ST2 and Diastolic Dysfunction in Hypertensive Patients. *Disease Markers*, vol. 2017, hh. 1-8.
- Fousteris, E., Melidonis, A., Panoutsopoulos, G., Tzirogiannis, K., Foussas, S., Theodosis-Georgilas, A., Tzerefos, S., Matsagos, S., Boutati, E., Economopoulos, T., Dimitriadis, G. dan Raptis, S. 2011. Toll/Interleukin-1 receptor member ST2 exhibits higher soluble levels in type 2 diabetes, especially when accompanied with left ventricular diastolic dysfunction. *Cardiovascular Diabetology*, vol. 10, no.1, h.101.
- Garcia-Compean, D., Jaquez-Quintana, J., Gonzalez-Gonzalez, J. dan Maldonado-Garza, H. 2009. Liver cirrhosis and diabetes: Risk factors, pathophysiology, clinical implications and management. *World Journal of Gastroenterology*, vol. 15, no.3, h.280.
- Grundy, S., Stone, N., Bailey, A., Beam, C., Birtcher, K., Blumenthal, R., Braun, L., de Ferranti, S., Faiella-Tommasino, J., Forman, D., Goldberg, R., Heidenreich, P., Hlatky, M., Jones, D., Lloyd-Jones, D., Lopez-Pajares, N., Ndumele, C., Orringer, C., Peralta, C., Saseen, J., Smith, S., Sperling, L., Virani, S. dan Yeboah, J. 2018. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: Executive Summary. *Journal of the American College of Cardiology*.

- Hartopo, A. B., Setianto, B. Y., Taufiq, N., 2016. Pengaruh kadar soluble ST2 serum terhadap risiko kejadian kardiovaskular mayor selama perawatan intensif pada infark miokard akut dengan elevasi segmen-ST. *Tesis*. Fakultas Kedokteran, Kesehatan Masyarakat dan Keperawatan Universitas Gadjah Mada, Program Pendidikan Dokter Spesialis Ilmu Penyakit Jantung dan Pembuluh Darah.
- Hartopo, A., Sukmasari, I. dan Puspitawati, I. 2018. The Utility of Point of Care Test for Soluble ST2 in Predicting Adverse Cardiac Events during Acute Care of ST-Segment Elevation Myocardial Infarction. *Cardiology Research and Practice*, hh.1-9.
- Ho, J., Sritara, P., deFilippi, C. dan Wang, T. 2015. Soluble ST2 Testing in the General Population. *The American Journal of Cardiology*, vol. 115, no.7, hh.22B-25B.
- Inggriani, M.P. 2019. Hubungan Kadar *Soluble* ST2 Serum dengan Disfungsi Ventrikel Kiri Akut pada Pasien Infark Miokard Akut dengan Elevasi Segmen-ST Anterior. Skripsi. Fakultas Kedokteran, Kesehatan Masyarakat dan Keperawatan Universitas Gadjah Mada.
- Ishihara, M., Kojima, S., Sakamoto, T., Kimura, K., Kosuge, M., Asada, Y., Tei, C., Miyazaki, S., Sonoda, M., Tsuchihashi, K., Yamagishi, M., Shirai, M., Hiraoka, H., Honda, T., Ogata, Y. dan Ogawa, H. 2009. Comparison of Blood Glucose Values on Admission for Acute Myocardial Infarction in Patients With Versus Without Diabetes Mellitus. *The American Journal of Cardiology*, vol. 104, no.6, hh.769-774.
- Iwakura, K. 2015. Stress hyperglycemia and microvascular obstruction after acute myocardial infarction. *Journal of Cardiology*, vol. 65, no.4, hh.270-271.
- Kakkar, R. dan Lee, R. 2008. The IL-33/ST2 pathway: therapeutic target and novel biomarker. *Nature Reviews Drug Discovery*, vol. 7, no.10, hh.827-840.
- Kementrian Kesehatan RI. 2013. Riset Kesehatan Dasar 2013. Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kementrian Kesehatan RI.
- Lazzeri, C., Valente, S., Chiostrri, M., D'Alfonso, M.G., dan Gensini, G.F. 2014. Clinical Significance of glycated hemoglobin in the acute phase of ST elevation myocardial infarction. *World J. Cardiol*, vol. 6, no.4, hh. 140 – 147.
- Libby, P. 2002. Inflammation and Atherosclerosis. *Circulation*, vol. 105, no.9, hh.1135-1143.
- Lilly, L. 2016. *Pathophysiology of heart disease*. 6th ed. Philadelphia: Wolters Kluwer.

- Martin, M. 2013. Special aspects of interleukin-33 and the IL-33 receptor complex. *Seminars in Immunology*, vol. 25, no.6, hh.449-457.
- Miller, A. dan Liew, F. 2011. The IL-33/ST2 pathway — A new therapeutic target in cardiovascular disease. *Pharmacology & Therapeutics*, vol. 131, no.2, hh.179-186.
- Miller, A., Purves, D., McConnachie, A., Asquith, D., Batty, G., Burns, H., Cavanagh, J., Ford, I., McLean, J., Packard, C., Shiels, P., Turner, H., Velupillai, Y., Deans, K., Welsh, P., McInnes, I. dan Sattar, N. 2012. Soluble ST2 Associates with Diabetes but Not Established Cardiovascular Risk Factors: A New Inflammatory Pathway of Relevance to Diabetes?. *PLoS ONE*, vol. 7, no. 10, h.e47830.
- Mueller, T. dan Jaffe, A. 2015. Soluble ST2—Analytical Considerations. *The American Journal of Cardiology*, vol. 115, no. 7, hh.8B-21B.
- Ota, S., Tanimoto, T., Orii, M., Hirata, K., Shiono, Y., Shimamura, K., Matsuo, Y., Yamano, T., Ino, Y., Kitabata, H., Yamaguchi, T., Kubo, T., Tanaka, A., Imanishi, T. dan Akasaka, T. 2015. Association between hyperglycemia at admission and microvascular obstruction in patients with ST-segment elevation myocardial infarction. *Journal of Cardiology*, vol. 65, no. 4, hh.272-277.
- Pascual-Figal, D. dan Januzzi, J. 2015. The Biology of ST2: The International ST2 Consensus Panel. *The American Journal of Cardiology*, vol. 115, no. 7, hh.3B-7B.
- Richards, A., Di Somma, S. dan Mueller, T. 2015. ST2 in Stable and Unstable Ischemic Heart Diseases. *The American Journal of Cardiology*, vol. 115, no. 7, hh.48B-58B.
- Rudijanto, A., Yuwono, A., Shahab, A., Manaf, A., Pramono, B., Lindarto, D., Purnamasari, D., Sanusi, H., Zufry, H., Novida, H., Suastika, K., Sucipto, K., Sasiarini, L., Dwipayana, M., Saraswati, M., Soetedjo, N., Soewondo, P., Soelistijo, S. dan Langi, Y. 2015. *Konsensus Pengelolaan dan Pencegahan Diabetes Mellitus Tipe 2 di Indonesia*. Jakarta: Pengurus Besar Perkumpulan Endokrinologi Indonesia.
- Sederholm Lawesson, S., Alfredsson, J., Szummer, K., Fredrikson, M. dan Swahn, E. 2015. Prevalence and prognostic impact of chronic kidney disease in STEMI from a gender perspective: data from the SWEDEHEART register, a large Swedish prospective cohort. *BMJ Open*, vol. 5, no. 6, h.e008188.
- Shah, A.M., Pfeffer, M.A., Hartley L.H., Moye, L.A., Gersh, B.J., Rutherford, J.D., Lamas, G.A., Rouleau, J.L., Braunwald, E., dan Solomon, S.D. 2010. Risk of All Cause Mortality, Recurrent Myocardial Infarction, and Heart Failure Hospitalization Associated With Smoking Status Following

- Myocardial Infarction With Left Ventricular Dysfunction. *The American Journal of Cardiology*, vol. 106, no. 7, hh. 911 – 916.
- Shen, J., Liu, J., dan Zhang, G. 2018. Interleukin-33 in Malignancies: Friends or Foes?. *Frontiers in Immunology*, vol. 9, no. 3051, hh. 1-7.
- Shimpo, M., Morrow, D., Weinberg, E., Sabatine, M., Murphy, S., Antman, E. dan Lee, R. 2004. Serum Levels of the Interleukin-1 Receptor Family Member ST2 Predict Mortality and Clinical Outcome in Acute Myocardial Infarction. *Circulation*, vol. 109, no. 18, hh.2186-2190.
- Singer, M., Deutschman, C., Seymour, C., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard, G., Chiche, J., Coopersmith, C., Hotchkiss, R., Levy, M., Marshall, J., Martin, G., Opal, S., Rubenfeld, G., van der Poll, T., Vincent, J. dan Angus, D. 2016. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*, vol. 315, no. 8, h.801.
- Stranders, I., Diamant, M., van Gelder, R., Spruijt, H., Twisk, J., Heine, R. dan Visser, F. 2004. Admission Blood Glucose Level as Risk Indicator of Death After Myocardial Infarction in Patients With and Without Diabetes Mellitus. *Archives of Internal Medicine*, vol. 164, no. 9, h.982.
- Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report. 2002. *Circulation*, vol. 106, no. 25, hh.3143-3143.
- Weinberg, E. 2002. Expression and Regulation of ST2, an Interleukin-1 Receptor Family Member, in Cardiomyocytes and Myocardial Infarction. *Circulation*, vol. 106, no. 23, hh.2961-2966.
- Weinberg, E., Shimpo, M., Hurwitz, S., Tominaga, S., Rouleau, J. dan Lee, R. 2003. Identification of Serum Soluble ST2 Receptor as a Novel Heart Failure Biomarker. *Circulation*, vol. 107, no. 5, hh.721-726.
- Willerson, J. 2007. *Cardiovascular medicine*. 3rd ed. London: Springer.
- Yuliani, D.D. 2019. Hubungan Antara Kadar ST2 Serum dengan Keberhasilan Fibrinolisis pada Pasien Infark Miokard Akut dengan Elevasi Segmen ST. Skripsi. Fakultas Kedokteran, Kesehatan Masyarakat dan Keperawatan Universitas Gadjah Mada.
- Zhao, S., Murugiah, K., Li, N., Li, X., Xu, Z., Li, J., Cheng, C., Mao, H., Downing, N., Krumholz, H. dan Jiang, L. 2017. Admission Glucose and In-hospital Mortality after Acute Myocardial Infarction in Patients with or without Diabetes. *Chinese Medical Journal*, vol. 130, no. 7, hh.767-775.