

DAFTAR PUSTAKA

- Abbas, AK. Andrew, HL., Shiv P. Cellular Molecular Immunology, *Elsevier Saunders*, 2007
- Abd-Elazeem MI, Mohamaed RA. Neutrophil Lymphocyte and Platelet Lymphocyte Ratio in Rheumatoid Arthritis Patients: Relation to Disease Activity. *The Egyptian Rheumatologist*. 2018; 40(4):227-231
- Abdulrahman MA, Afifi N. Ashry M. Neutrophil/ Lymphocyte and platelet/ lymphocyte ratio are useful predictor comparable to serum IL 6 for disease activity and damage in naïve and relapsing patients with lupus nephritis, *The Egyptian Rheumatologist*, 2019
- Al Arfaj, A. Khalil, N. Clinical and immunological manifestations in 624 SLE patients in Saudi Arabia, *Lupus*. 2009, 18(5), 465–473.
- Attilio LD, Santucci N, Bongiovanni B, Bay ML. Tuberculosis , the disrupted immune-endocrine response and the potential thymic repercussion as a contributing factor to disease physiopathology. *Front Endocrinol*. 2018;9:214.
- Ayhan MG, Cicek IE, Inanli I, Caliskan AM, Ercan SK, Eren I. Neutrophil/lymphocyte and platelet/lymphocyte ratios in all mood states of bipolar disorder. *Psychiatry Clin Psychopharmacol*. 2017;27(3):284–8. 13
- Ayna, AB., Ermurat, S., Coskun, BN., Harman, H., Pehlivan, Y. Neutrophil to Lymphocyte Ratio and Mean Platelet Volume as Inflammatory Indicators in Systemic Lupus Erythematosus Nephritis. *Arch Rheumatol* 2017;32(1):21-25

- Azab, B., Shah N. Akerman M, McGinn JT . Value of platelet/lymphocyte ratio as a predictor of all-cause mortality after non-ST-elevation myocardial infarction, *J. Thromb. Thrombolysis*, 2012. 34 (3) 326–334
- Barilla-Labarca, ML., Toder, K., Furie, R.. Targeting the complement system in systemic lupus erythematosus and other diseases, *Clin. Immunol.* 2013. 148 (3); 313–321.
- Barnes, EV., Narain S, Naranjo A. Shuster J, Segal MS, Sobel ES . High sensitivity C-reactive protein in systemic lupus erythematosus: relation to disease activity, clinical presentation and implications for cardiovascular risk, *Lupus* 2005. 14 (8) 576–582.
- Bashal, F. Hematological disorders in patients with systemic lupus erythematosus. *Open Rheumatol J* 2013;7:87-95.
- Bertsias, GK., Ioannidis, JPA., Boletis, J., Bombardieri, S., Cervera, R., Dostal, C. EULAR recommendations for the management of systemic lupus erythematosus (SLE). Report of a Task Force of the European Standing Committee for International Clinical Studies Including Therapeutics (ESCISIT). *Ann Rheum Dis.* 2008;67:195–205.
- Blank, M., Shoenfeld, Y., Perl, A. Crosstalk of the environment with the host genome and the immune system through endogenous retroviruses in systemic lupus erythematosus. *Lupus.* 2009; 18(13):1136-43.
- Boilard, E., Blanco, P., Nigrovic, PA. Platelets: active players in the pathogenesis of arthritis and SLE. *Nat Rev Rheumatol* 2012;8:534-42.

- Bosch, X. Systemic lupus erythematosus and the neutrophil. *N Engl J Med.* 2011; 365(8):758-60.
- Braunwald, E., Fauci, AS., Kasper, DL., Hauser, SL., Longo, DL., Jameson, JL. *Harrison's Principle of Internal Medicine.* Ed-18. USA: McGrawHill Professional Publishing; 2012.
- Bromberg L, Roufosse F, Pradier O, Delporte C, Van Antwerpen P, De Maertelaer V, Cogan E, Methylprednisolone-induced lymphocytosis in patients with immune mediated inflammatory disorders, *The American Journal of Medicine.* 2016
- Buyon, JP. Systemic lupus erythematosus a clinical and laboratory features : Klippel JH, editor. Primer Primer on the rheumatic diseases. Ed-13. Atlanta: *Arthritis Foundation*; 2008. 303-18.
- Buyukkaya, E., Karakaş, M. F., Karakaş, E., Akçay, A. B., Tanboga, I. H., Kurt, M. Sen N. 'Correlation of Neutrophil to Lymphocyte Ratio With the Presence and Severity of Metabolic Syndrome', *Clinical and Applied Thrombosis/Hemostasis*, 2014.20(2),pp.159–163.
- Byong Sun Oh, Jeong Won Jang, Jung Hyun Kwon, 2013. Prognostic value of C-reactive protein and neutrophil-to-lymphocyte ratio in patients with hepatocellular carcinoma, *BMC Cancer*, 13:78.
- Chen, M., Daha, MR., Kallenberg, CG. The complement system in systemic autoimmune disease, *J. Autoimmun.* 2010. 34 (3) J276–J286.
- De Jager CP, Van Wijk PT, Wever PC. Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit. *Critical Care* 2010, 14:R192. .

- Demir S, Atli A, Bulut M, Okan İbiloğlu A, Güneş M, Kaya MC. Neutrophil-lymphocyte ratio in patients with major depressive disorder undergoing no pharmacological therapy. *Neuropsychiatr Dis Treat*. 2015;11:2253-8.
- Durmus E, Kivrak T, Gerin F, Sunbul M, Sari I, Erdogan O. Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio are Predictors of Heart Failure. *Arq Bras Cardiol*. 2015; 105(6):606-613
- Elhafeez HA, El-Meghawry ES, Omran TM, Study of Neutrophil Lymphocyte Ratio and Platelet Lymphocyte Ratio as Inflammatory Markers in Systemic Lupus Erythematosus Egyptian Patients. *Electronic Journal of Biology*. 2017 Jul 1;13(3)
- Fauci AS, Dale DC. Alternate-day prednisone therapy and human lymphocyte subpopulations. *J Clin Invest*. 1975;55(1):22-32.
- Feng, JF., Huang, Y., Chen, QX. Preoperative platelet lymphocyte ratio (PLR) is superior to neutrophil lymphocyte ratio (NLR) as a predictive factor in patients with esophageal squamous cell carcinoma, *World J. Surg. Oncol*. 2014. 12, 58.
- Flower, C., Hennis, A. J.M., Hambleton, I.R., Nicholson, G.D. Liang, M.H. Systemic lupus erythematosus in an African Caribbean population: Incidence, clinical manifestations, and survival in the Barbados National Lupus Registry, *Arthritis Care and Research*, 2012. 64(8), 1151–1158. doi: 10.1002/acr.21656.

- Folman CC, Linthorst GE, van Mourik J, van Willigen G, de Jonge E, Levi M, de Haas M, Platelet release trombopoetin(TPO) upon activation, another regulatory loop in thrombocytopoesis/ *Thromb Hemost*, 2000, 89; 923-930
- Gibson, PH., Croal BL, Cuthbertson BH, Small GR, Preoperative neutrophil-lymphocyte ratio and outcome from coronary artery bypass grafting, *Am. Heart J.* 154 (5) 2007. 995–1002
- Giron Gonzales JA, Martinez Sierra C., Rodriguez-Ramos C, Marcioas MA, Rendon P, Diaz F, 2004, Implication of Inflammation related cytokines in The Natural History of Liver Cirrhosis, *Liver Int:* 24:473- 445.
- Goldblatt, F., Chambers S. Rahman A, Isenberg DA. Serious infections in British .patients with systemic lupus erythematosus: hospitalisations and mortality, *Lupus* 2009. 18 (8) 682–689.
- Gorial Faiq I. A Clinical Utility of Neutrophil Lymphocyte Ratio as A Prognostic Indicator of Systemic Lupus Erythematosus Disease Activity *J. Pharm. Sci. dan Res.* Vol. 10(3), 2018, 637-639
- Hanahan, D., Weinberg, R.A. The hallmarks of cancer. *Cell.*2000,100(1):57–70
- Hariadi, H. *Lupus dan Penatalaksanaannya* Edisi ke-3. Surabaya: Fakultas Kedokteran UNAIR; 2007.
- Harris, ED., Hoes, JN., Jacobs, JWG., Boers, M., Boumpas, D., Buttgerit, F., Caeyers, N., et al. EULAR evidence based recommendations on the management of systemic glucocorticoid therapy in rheumatic diseases. *Ann Rheum Dis.* 2007; 66: 1560-7.

- Hepburn AL, Narat S, Mason JC. The management of peripheral blood cytopenias in systemic lupus erythematosus. *Rheumatol.* 2010;49:2243-54)
- Hung HY, Chen JS, Yeh CY, Changchien CR, Tang R, Hsieh PS, et al. Effect of preoperative neutrophillymphocyte ratio on the surgical outcomes of stage II colon cancer patients who do not receive adjuvant chemotherapy. *Int J Colorectal Dis* 2011;26:1059-65
- Illei, GG. Tackey E.Lapteva L, Lipsky PE, Biomarkers in systemic lupus erythematosus. I. General overview of biomarkers and their applicability, *Arthritis Rheum.* 2004. 50 (6) ; 1709–1720.
- Jacobs, JWG., Bijlsma, JWJ. Glucocorticoid therapy. Dalam: Firestein GS, Budd RC, Harris ED, Hoes JN, Jacobs JWG, Boers M, Boumpas D, Buttgerit F, Caeyers N, et al. EULAR evidence based recommendations on the management of systemic glucocorticoid therapy in rheumatic diseases. *Ann Rheum Dis.* 2007; 66: 1560-7.
- Jaillon, S. Galdiero MR.Del Prete D, Cassatela MA, Garlanda C, Mantovani A, Neutrophils in innate and adaptive immunity, *Semin. Immunopathol.* 2013. 35 (4) 377–394.
- Janoudi N, Haematological Manifestations in Systemic Lupus Erythematosus, *KFSH RC, Jeddah Saudi Arabia*, 2012 , hal. 363–382. doi: 10.1016/0268-960X(93)90006-P.
- Kaplan J Mariana. Neutrophils in the pathogenesis and manifestations of SLE, *Nature Review Rheumatology*, Macmilan Publisher, 2011; 691-669

- Kasjmir, YI. Handono, K., Wijaya, LK., Hamijoyo, L., Albar, Z., Kalim, H., Hermansyah., Kertia, N., Achadiono, DNW., Manuaba, IARW., Dewi, S., Ongkowijaya, JA., Isbagio, H., Setyohadi, B., Suarjana, N. 2014. *Diagnosis dan Pengelolaan Lupus Eritematosus Sistemik dalam Buku Ajar Ilmu Penyakit Dalam Jilid III edisi VI*. Jakarta. Interna Publishing.2260-3375
- Kayhan F, Gündüz S, Ersoy SA, Kandeğer A, Annagür BB. Relationships of neutrophil–lymphocyte and platelet–lymphocyte ratios with the severity of major depression. *Psychiatry Res*. 2017;247:332-5
- Kedziora-Kornatowska, KZ. Production of superoxide and nitric oxide by granulocytes in non-insulin-dependent diabetic patients with and without diabetic nephropathy, *IUBMB Life* 1999. 48 (3), 359–362.
- Kemenkes RI. . *Situasi Lupus di Indonesia*. Jakarta.Pusat 2017. Data dan Informasi Kementerian Kesehatan RI.
- Keenan, RT., Swearingen, CJ., Yazici, Y. 2008. Erythrocyte sedimentation rate and Creactive protein levels are poorly correlated with clinical measures of disease activity in rheumatoid arthritis, systemic lupus erythematosus and osteoarthritis patients, *Clin. Exp. Rheumatol*. 26 (5) (2008) 814–819.
- Kocyigit I, Eroglu E, Unal A, Sipahioglu MH, Tokgoz B, Oymak O. Role of neutrophil/lymphocyte ratio in prediction of disease progression in patients with stage–4 chronic kidney disease. *Journal Nephrology*. 2013; 26(02): 358-65.
- Koops HS. Lymphopenia and autoimmune diseases, *Arthritis Res Ther* 2004; 6: 178-

Krishnan, S., Chowdhury, B., Juang, YT., Tsokos, GC. *Overview of the Pathogenesis of Systemic Lupus Erythematosus*. Philadelphia: Mosby, Inc.; 2007.

Kyrtaris, VC., Tsokos, GC. *New Treatments in Systemic Lupus Erythematosus*. Philadelphia: Mosby, Inc.; 2007.

Lee, YH., Song, GG. .Association of Neutrophil to Lymphocyte Ratio, Platelet to Lymphocyte Ratio, and Mean Platelet Volume with Systemic Lupus Erythematosus Disease Activity: A Meta-analysis. *Journal of Rheumatic Diseases* Vol. 24, No. 5, October, 2017

Lee JS, Kim NY, Na SH, Youn HY, Shin CS. Reference values of neutrophil-lymphocyte ratio, lymphocyte-monocyte ratio, platelet-lymphocyte ratio, and mean platelet volume in healthy adults in South Korea, *Medicine*, 2018, 97 : 26

Li J, Pan Z, Liu H, Ding F, Shu Q, Li X. Retrospective analysis of the risk of hemorrhage associated with moderate and severe thrombocytopenia of 173 patients with systemic lupus erythematosus. *Medicine* (Baltimore) 2018;97.

Li L, Xia Y, Chen C, Cheng P, Peng C. Neutrophil-lymphocyte ratio in systemic lupus erythematosus disease: a retrospective study. *Int J Clin Exp Med* 2015;8:11026-31.

Lipsky, PE. Systemic lupus erythematosus: an autoimmune disease of B cell hyperactivity. *Nat Immunol* 2001;2:764-6.

Lou, M., Luo, P., Tang, R., Peng, Y., Yu, S., Huang, W., He, L.. Relationship between neutrophil-lymphocyte ratio and insulin resistance in newly diagnosed type 2 diabetes mellitus patients. *BMC Endocrine Disorders* 2015. 15:9

McInnes, IB., Ruddy, S., Sergent, JS., editor. *Kelley's Textbook of rheumatology*.

Edisi ke-8. Philadelphia: WB Saunders Elsevier; 2009. 863-81.

Mok, C. C. dan Lau, C. S. Lupus in Hong Kong Chinese, *Lupus*, 2003, 12(9), 717–722. doi: 10.1191/0961203303lu451xx.

Newman K ,Owlia M , EL-Hemaidi I, Akhtari M. : Management of cytopenias in patients with systemic lupus erythematosus Old and new. *Autoimmunity Reviews*,2013, 12(7):784-91.

Nikolsky E, Grines CL, Cox DA, Garcia A. Platelet-lympocyte ratio : A simple, inexpensive and rapid prognostic marker for cardiovascular event : *Pletelets* : Vol 26, No7. 2014

Ntali, S., Tzabakakis, M., Bertias, G, Boumpas DT. What's new in clinical trials in lupus. *Int J Clin Rheum*. 2009;4(4):473-85.

Nugroho Adi, Suwarman, Nawawi AM. Hubungan Antara Rasio Netrofil Limfosit dan skor sequential organ failure Assessment Pada Pasien yang Dirawat di Ruang Intensive Care Unit. *Jurnal Anastesi Perioperatif*. 2013; 1 (3):189-196

Nunez, J., Bodi V.Nunez E,Llacer A. Usefulness of the neutrophil to lymphocyte ratio in predicting longterm mortality in ST segment elevation myocardial infarction, *Am. J. Cardiol*. 2008. 101 (6) (2008) 747–752.

Ohadian A ,Suryadinata H , Dewi S. : The role of neutrophyl lymphocyte count ratio as an inflammatory marker in systemic lupus erythematosus. *Acta Med. Indonesia*, 2013. 45(3):170-174.

Ohl, K, Tenbrock, K. Inflammatory cytokines in systemic lupus erythematosus. *J Biomed Biotechnol* ; 2011: 432595

- Oylumlu M, Yıldız A, Oylumlu M, Yüksel M, Polat N, Bilik MZ, et al. Platelet to lymphocyte ratio is a predictor of in-hospital mortality patients with acute coronary syndrome. *Anatol J Cardiol*. 2015;15(4):277-83
- Park BH, Kang YA, Kim YS. Delta neutrophil index as an early marker of disease severity in critically ill patients with sepsis *BMC Infectious Diseases*, 11:299. 2011
- Peredo R, Vilá S, Goñi M, Reactive thrombocytosis: an early manifestation of Takayasu arteritis. *J Clin Rheumatol* 2005;11:270–3.
- Petri, M., Orbai, AM., Alarcón, GS., Gordon, C., Merrill, JT., Fortin, PR., et al. Derivation and validation of the Systemic Lupus International Collaborating Clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum*. 2012; 64(8):2677-86. .
- Qin, B., Ma, N., Tang, Q., Wei, T., Yang, M., Fu, H., Hu, Z., Liang, Y., Yang, Z. 2015. Neutrophil to lymphocyte ratio (NLR) and platelet to lymphocyte ratio (PLR) were useful markers in assessment of inflammatory response and disease activity in SLE patients. *Mod Rheumatol*, 2016
- Rahman, A., Isenberg, DA. Systemic lupus erythematosus. *N Engl J Med*. 2008; 358(9):929-39.
- Rezaieyazdi, Z., Farid R., Mirfeizi Z, Hatef MR, Mirheidari M, Mansouri H, Is there any correlation between high sensitive CRP and disease activity in systemic lupus erythematosus? *Lupus* 2011. 20 (14); 1494–1500.
- Rosales C. Neutrophil : A Cell with many roles in inflammation or several Cell Types? *Frontier Physiol* , 2016, 9 : 113

Salido, O.Reyes, M. “Epidemiology of systemic lupus erythematosus in Asia,”

Lupus, 2010. 13651372.

Sanchez, E., Nadig, A., Richardson, BC., Freedman, BI., Kaufman, KM., Kelly, JA.,

et al. Phenotypic associations of genetic susceptibility loci in systemic lupus erythematosus. *Ann Rheum Dis*. 2011; 70(10):1752-7.

Sefil, F., Ulutas, K. T., Dokuyuku, R. Investigation of neutrophil lymphocyte ratio and blood glucose regulation in patients with type 2 diabetes mellitus.

Journal of International Medical Research. 2014 Vol. 42(2) 581–588

Sherer Y, Gorstein A, Fritzler MJ, Shoenfield Y . Autoantibody explosion in systemic

lupus erythematosus: more than 100 different antibodies found in SLE patients, *Semin. Arthritis Rheum*. 2004. 34 (2) 501–537.

Soliman WA, Nahed MS, Iman M, Ghanima. Mohja A, El-badawy. Neutrophil to

lymphocyte ratio (NLR) and platelet to lymphocyte ratio (PLR) in Systemic Lupus Erythematosus : relation with disease Activity and Lupus Nephritis, *Reumatol Clin*, 2018.

Suarjana, IN. *Imunopatogenesis Lupus Eritematosus Sistemik dalam Buku Ajar Ilmu*

Penyakit Dalam Jilid III edisi VI. Jakarta. Interna Publishing. 2014. hal 3331-3343.

Takele Y, Adem E, Getahun M, Tajebe F. Kiflie, Hailu A. Malnutrition in healthy

Individuals Result in increased Mixex Cytokine Profiles. Altered Neutrophyl Subsets and Fuction. *Plos ONE*, 2016. 11(8)

- Tan TP, Arekapudi A, Metha J. Prasad A, Venkatraghavan. Neutrophil-Lymphocyte ratio as predictor of mortality and morbidity in cardiovascular surgery : a systemic review, *ANZ J Surg*, 2015, 85 (6)
- Tucker PS, Scanlan AT, Dalbo VJ. Review Article Chronic Kidney Disease Influences Multiple Systems: Describing the Relationship between Oxidative Stress, Inflammation, Kidney Damage, and Concomitant Disease. *Oxidative Medicine and Cellular Longevity Hindawi Publishing Corporation*. 2015: 1-8.
- Üsküdar Cansu D, Üsküdar Teke H, Musmul A, Korkmaz C. Is thrombocytosis always an indicator of autosplenectomy in patients with systemic lupus erythematosus? *Rheumatol Int* 2018;38:239-47.
- Uslu, AA. Adem K, Sbahin A, Ugan Y, Yilmaz R. . Two new inflammatory markers associated with Disease Activity Score-28 in patients with rheumatoid arthritis: neutrophil-lymphocyte ratio and platelet-lymphocyte ratio, *Int. J. Rheum. Dis.* 2015
- Varaprasad, IR., Agrawal, S., Prabu, VN., Rajasekhar, L., Kanikannan, MA., Narsimulu, G. Posterior reversible encephalopathy syndrome in systemic lupus erythematosus. *J Rheumatol*. 2011; 38(8):1607-11.
- Vigano A, Bruera E, Jhangri GS, Newman SC, Fiel AL, Suarez ME. Clinical survival predictors in patient with advanced cancer. *Arch Intern Med*. 2000; 160: 861-868
- Williams Jr, RC.,Harmon ME, Burlingame R, Du Clos TW, Studies of serum C-reactive protein in systemic lupus erythematosus, *J. Rheumatol*. 2005. 32 (3) 454-461.

- Wong, CK., Ho, CY., Li, EK., Lam, CW. Elevation of proinflammatory cytokine (IL-18, IL-17, IL-12) and Th2 cytokine (IL-4) concentrations in patients with systemic lupus erythematosus. *Lupus* 2000;9:589-93.
- Xia Y, Guo XG, Xing JT, Chen Q. Neutrophil Count to Lymphocyte Count Ratio is a Potential Diagnostic Index for Bacteremia in Adult. *Life Science Journal*. 2014;11(1). 172-7.
- Xiang R. Depression' s influence on the immune system. *J Undergrad Life Sci*. 2007;1(1):24–6.
- Yildiz, A., Yuksel M, Oylumlu M, Polat N, Akyuz A. The utility of the platelet-lymphocyte ratio for predicting no reflow in patients with ST-segment elevation myocardial infarction, *Clin. Appl. Thromb. Hemost*. 2015. 21 (3) 223–228.
- Yolbas, S., Yildirim, A., Gozel, N., Uz, B., Koca, SS. 2016. Hematological Indices May Be Useful in the Diagnosis of Systemic Lupus Erythematosus and in Determining Disease Activity in Behçet's Disease. *Med Princ Pract* 2016;25:510–516
- Yu Haitao, Jiang Lili, Yao Liqiong, Gan chao, Han Xinwen, Liu Ruiqi, Su Na : Predictive value of the neutrophil to lymphocyte ratio and hemoglobin in systemic lupus erythematosus and hemoglobin in systemic lupus erythematosus , Experimental and Therapeutic, *Medicine* 16: 1547-1553, 2018
- Zahorec R,. Ratio of neutrophil to lymphocyte counts rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl. LekListy* , 2001; 102: 5-14.



Hubungan antara MEX-SLEDAI dengan Neutrophyl Lymphocyte Ratio dan Platelet Lymphocyte Ratio pada

penderita Lupus Eritematosus Sistemik

Andris Senggetan, Dr. dr. Deddy Nur Wachid A, M.Kes, SpPD, KR

Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Zhang, H., Sun, Q., Mao, W., Fan, J., Ye, B.,. Neutrophil-to-Lymphocyte Ratio Predicts Early Mortality in Patients with HBV-Related Decompensated Cirrhosis. *Gastroenterol Res Pract* , 2016. 16: 2200-2205