

REFERENCES

- Butterworth, J. F., Mackey, D. C. and Wasnick, J. D. (2013) *Clinical Anesthesiology, Anesthesia & Analgesia*. doi: 10.1213/00000539-199210000-00057.
- Beverly, A., Kaye, A., Ljungqvist, O. and Urman, R. (2017). Essential Elements of Multimodal Analgesia in Enhanced Recovery After Surgery (ERAS) Guidelines. *Anesthesiology Clinics*, 35(2), pp.e115-e143.
- Bithal *et al.*, (2002). Haemodynamic Changes In Response To Skull-Pins Application - Comparison Between Normotensive And Hypertensive Patients. *Indian Journal of Anaesthesia*, 46(5): 381-383.
- Brown, E., Lydic, R. and Schiff, N. (2010). General Anesthesia, Sleep, and Coma. *New England Journal of Medicine*, 363(27), pp.2638-2650.
- Can, B. and Bilgin, H. (2017). Effects of scalp block with bupivacaine versus levobupivacaine on haemodynamic response to head pinning and comparative efficacies in postoperative analgesia: A randomized controlled trial. *Journal of International Medical Research*, 45(2), pp.439-450.
- Canakci, E. (2017). Our Scalp Block Results in Craniotomy Cases. *Journal of Anesthesia & Intensive Care Medicine*, 2(4).
- Chowdhury, T., Baron, K. and Cappellani, R. (2013). Severe bradycardia during scalp nerve block in patient undergoing awake craniotomy. *Saudi Journal of Anaesthesia*, 7(3), p.356.
- Duke, J. C. and Keech, B. M. (2016) *Duke's Anesthesia Secrets, Saunders, an Imprint of Elsevier Inc.*
- Feldman, L., Lee, L. and Fiore, J. (2014). What outcomes are important in the assessment of Enhanced Recovery After Surgery (ERAS) pathways?. *Canadian*

Journal of Anesthesia/Journal canadien d'anesthésie, 62(2), pp.120-130.

Flexman, A. M., Ng, J. L. and Gelb, A. W. (2010) 'Acute and chronic pain following craniotomy', *Current Opinion in Anaesthesiology*, 23(5), pp. 551–557. doi: 10.1097/ACO.0b013e32833e15b9.

Gazoni, F., Pouratian, N. and Nemergut, E. (2008). Effect of ropivacaine skull block on perioperative outcomes in patients with supratentorial brain tumors and comparison with remifentanyl: a pilot study. *Journal of Neurosurgery*, 109(1), pp.44-49.

Geze, S., Yilmaz, A. and Tuzuner, F. (2009). The effect of scalp block and local infiltration on the haemodynamic and stress response to skull-pin placement for craniotomy. *European Journal of Anaesthesiology*, 26(4), pp.298-303.

de Gray, L. C. and Matta, B. F. (2005) 'Acute and chronic pain following craniotomy : a review', *Blackwell Publishing*, 23(5), pp. 693–704. doi: 10.1097/ACO.0b013e32833e15b9.

Guilfoyle, M., Helmy, A., Duane, D. and Hutchinson, P. (2013). Regional Scalp Block for Postcraniotomy Analgesia. *Anesthesia & Analgesia*, 116(5), pp.1093-1102.

Halldin, C. B. *et al.* (2009) 'Nerve blocks enable adequate pain relief during topical photodynamic therapy of field cancerization on the forehead and scalp', *British Journal of Dermatology*, 160(4), pp. 795–800. doi: 10.1111/j.1365-2133.2008.09024.x.

Hassani, V., Movassaghi, G., Safaian, R., Safari, S., Zamani, M.M., Hajiashrafi, M., Sedaghat, M. (2014). Bupivacaine-Sufentanyl Versus Bupivacaine-Fentanyl in Spinal Anesthesia of Patients Undergoing Lower Extremity Surgery. *Anesthesiology and Pain Medicine* 4.

Kaye, A. (2005). *Essential Neurosurgery*. 3rd ed. Wiley-Blackwell.

Kaushic A. Theerth, Kamath Sriganesh, K. Madhusudan Reddy, Dhritiman Chakrabarti, Ganne S. Umamaheswara Rao (2018). Analgesia Nociception Index-guided intraoperative fentanyl consumption and postoperative analgesia in patients receiving scalp block versus incision-site infiltration for craniotomy. *Minerva Anesthesiol* 84.

Leone, S. *et al.* (2008) ‘Pharmacology, toxicology, and clinical use of new long acting local anesthetics, ropivacaine and levobupivacaine’, *Acta Biomedica de l’Ateneo Parmense*, 79(2), pp. 92–105.

Osborn, I. and Sebeo, J. (2010) ‘“Scalp block” during craniotomy: A classic technique revisited’, *Journal of Neurosurgical Anesthesiology*, 22(3), pp. 187–194. doi: 10.1097/ANA.0b013e3181d48846.

Papangelou, A., Radzik, B., Smith, T. and Gottschalk, A. (2013). A review of scalp blockade for cranial surgery. *Journal of Clinical Anesthesia*, 25(2), pp.150-159.

Sihombing, R., Yulianti Bisri, D. and Herman Sitanggang, R. (2018). Perbandingan Antara Fentanil 2 µg/kgBB/jam dan Scalp Block Terhadap Peningkatan Hemodinamik dan Kadar Glukosa Darah Sewaktu Saat Pemasangan Pin Kepala Pada Kraniotomi. *Jurnal Neuroanestesi Indonesia*, 8(2), pp.80-87.

Suwarman and Bisri, T. (2016) ‘Pengelolaan Nyeri Pascakraniotomi Postcraniotomy Pain Management’, 5(1), pp. 68–76.

Talke, P. and Gelb, A. (2005). Postcraniotomy pain remains a real headache!. *European Journal of Anaesthesiology*, 22(5), pp.325-327.

Tuchinda, L., Somboonviboon, W., Supbornsug, K., Worathongchai, S. and Limutaitip, S. (2010). Bupivacaine scalp nerve block: hemodynamic response during craniotomy, intraoperative and post-operative analgesia. *Asian Biomedicine*, 4(2), pp.243-251.

Vahabi, S. and Beiranvand, S. (2017) 'Effect of Local Ropivacaine on Hemodynamic Responses in Craniotomy Patients, Journal of Investigative Surgery'. Journal of Investigative Surgery, p. 5.

Young, W. B. (2010) 'Blocking the greater occipital nerve: Utility in headache management', *Current Pain and Headache Reports*, 14(5), pp. 404–408. doi: 10.1007/s11916-010-0130-x.