

REFERENCES

- Abecassis, M., Bartlett, S.T., Collins, A.J., Davis, C.L., Delmonico, F.L., Friedewald, J.J., Hays, R., Howard, A., Jones, E., Leichtman, A.B. and Merion, R.M., 2008. Kidney transplantation as primary therapy for end-stage renal disease: a National Kidney Foundation/Kidney Disease Outcomes Quality Initiative (NKF/KDOQI™) conference. *Clinical Journal of the American Society of Nephrology*, 3(2), pp.471-480.
- Al Wakeel, J., Al Harbi, A., Bayoumi, M., Al-Suwaida, K., Al Ghonaim, M. and Mishkiry, A. (2012). Quality of life in hemodialysis and peritoneal dialysis patients in Saudi Arabia.
- Bakewell, A.B., Higgins, R.M. and Edmunds, M.E., 2002. Quality of life in peritoneal dialysis patients: decline over time and association with clinical outcomes. *Kidney international*, 61(1), pp.239-248.
- Bauer, C., Melamed, M.L. and Hostetter, T.H., 2008. Staging of chronic kidney disease: time for a course correction. *Journal of the American Society of Nephrology*, 19(5), pp.844-846.
- Blake, P.G., Bargman, J.M., Brimble, K.S., Davison, S.N., Hirsch, D., McCormick, B.B., Suri, R.S., Taylor, P. and Zalunardo, N., 2011. CANADIAN SOCIETY OF NEPHROLOGY GUIDELINES/RECOMMENDATIONS. *Peritoneal Dialysis International*, 31, pp.218-239.
- Broers, N.J., Martens, R.J., Canaud, B., Cornelis, T., Dejagere, T., Diederens, N.M., Hermans, M.M., Konings, C.J., Stifft, F., Wirtz, J.J. and Leunissen, K.M., 2018. Health-related quality of life in end-stage renal disease patients: the effects of starting dialysis in the first year after the transition period. *International urology and nephrology*, 50(6), pp.1131-1142.
- Burbank, J.H., Brugger, J.M., Treu, D.M. and Wyeth, M.T., NxStage Medical Inc, 2019. *Peritoneal Dialysis Systems, Devices, and Methods*. U.S. Patent Application 16/360,396.
- Burkart, J.M., Patient education: Peritoneal dialysis (Beyond the Basics).
- CHAN, J., CHE, K., LAM, K., CHOW, K., CHUNG, K., LI, P. and SZETO, C. (2007). Comprehensive malnutrition inflammation score as a marker of nutritional status in Chinese peritoneal dialysis patients.
- ElHafeez, S.A., Sallam, S.A., Gad, Z.M., Zoccali, C., Torino, C., Tripepi, G., ElWakil, H.S. and Awad, N.M., 2012. Cultural adaptation and validation of the “Kidney Disease and Quality of Life-Short Form (KDQOL-SF™) version 1.3” questionnaire in Egypt. *BMC nephrology*, 13(1), p.170.

González-Ortiz, A.J., Arce-Santander, C.V., Vega-Vega, O., Correa-Rotter, R. and Espinosa-Cuevas, M.A., 2015. Assessment of the reliability and consistency of the “malnutrition inflammation score” (MIS) in Mexican adults with chronic kidney disease for diagnosis of protein-energy wasting syndrome (PEW). *Nutricion hospitalaria*, 31(3), pp.1352-1358.

Grant, K.L., Wilt, M.J., Tracey, B.D., Rudolf, B.A., Collins, D.E., Gustin, L.A., Dale, J.D. and Bodwell, J.T., Deka Products LP, 2017. *Hemodialysis system*. U.S. Patent 9,724,458.

Ibrahim, S. and El Salamony, O. (2008). Depression, Quality of Life and Malnutrition-Inflammation Scores in Hemodialysis Patients. *American Journal of Nephrology*, 28(5), pp.784-791.

Iorember, F.M., 2018. Malnutrition in chronic kidney disease. *Frontiers in pediatrics*, 6.

Jankowska-Polańska, B., Uchmanowicz, I., Wysocka, A., Uchmanowicz, B., Lomper, K. and Fal, A.M., 2017. Factors affecting the quality of life of chronic dialysis patients. *European Journal of Public Health*, 27(2), pp.262-267.

Kdigo.org. (2020). [online] Available at: https://kdigo.org/wp-content/uploads/2017/04/KDIGO-CKD-Guideline-Manila_Kasiske.pdf [Accessed 22 Jan. 2020].

Kistler, B.M., Benner, D., Burrowes, J.D., Campbell, K.L., Fouque, D., Garibotto, G., Kopple, J.D., Kovesdy, C.P., Rhee, C.M., Steiber, A. and Stenvinkel, P., 2018. Eating during hemodialysis treatment: a consensus statement from the International Society of Renal Nutrition and Metabolism. *Journal of renal nutrition*, 28(1), pp.4-12.

Kovesdy, C.P., 2016, January. Malnutrition in dialysis patients—the need for intervention despite uncertain benefits. In *Seminars in dialysis* (Vol. 29, No. 1, pp. 28-34).

Neri L, e. (2005). Work ability and health status in dialysis patients. - PubMed - NCBI. [online] Ncbi.nlm.nih.gov. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16015564> [Accessed 7 Jan. 2020].

Phillips, S. and Knuchel, N., 2011. Chronic kidney disease: nutrition basics. *Journal of Renal Nutrition*, 21(4), pp. e15-e17.

Rambod, M., Bross, R., Zitterkoph, J., Benner, D., Pithia, J., Colman, S., Kovesdy, C., Kopple, J. and Kalantar-Zadeh, K. (2008). Association of Malnutrition-Inflammation Score with Quality of Life and Mortality in Hemodialysis Patients: A 5-Year Prospective Cohort Study.

Rees, L., 2019. Assessment of dialysis adequacy: beyond urea kinetic measurements. *Pediatric Nephrology*, 34(1), pp.61-69.

Slee, A. (2012). Exploring metabolic dysfunction in chronic kidney disease. *Nutrition & Metabolism*, 9(1), p.36.

Sohrabi, Z., Eftekhari, M.H., Eskandari, M.H., Rezaeianzadeh, A. and Sagheb, M.M., 2015. Malnutrition-inflammation score and quality of life in hemodialysis patients: is there any correlation? *Nephro-urology monthly*, 7(3).

Sthepens, C. (2018). Healthline Media. *Kidney Overview*

White, C.A., Akbari, A., Talreja, H., Lalani, N. and Knoll, G.A., 2016. Classification of kidney transplant recipients using a combination of estimated GFR and albuminuria reflects risk. *Transplantation direct*, 2(8).

Webster, A.C., Nagler, E.V., Morton, R.L. and Masson, P., 2017. Chronic kidney disease. *The lancet*, 389(10075), pp.1238-1252.

Yang, F., Griva, K., Lau, T., Vathsala, A., Lee, E., Ng, H.J., Mooppil, N., Foo, M., Newman, S.P., Chia, K.S. and Luo, N., 2015. Health-related quality of life of Asian patients with end-stage renal disease (ESRD) in Singapore. *Quality of Life Research*, 24(9), pp.2163-2171.

Yang, F., Lau, T. and Luo, N., 2016. Cost-effectiveness of hemodialysis and peritoneal dialysis for patients with end-stage renal disease in Singapore. *Nephrology*, 21(8), pp.669-677.