

DAFTAR PUSTAKA

- Ali, M., Bukhari, S., Ali, M. and Lee, H. (2017). Upstream signalling of mTORC1 and its hyperactivation in type 2 diabetes (T2D). *BMB Reports*, 50(12), pp.601-609.
- Batuman, V. (2019). *Diabetic Nephropathy: Practice Essentials, Pathophysiology, Etiology*. [online] Emedicine.medscape.com. Available at: <https://emedicine.medscape.com/article/238946-overview#a3> [Accessed 18 Sep. 2019].
- Blandino-Rosano, M., Chen, A., Scheys, J., Alejandro, E., Gould, A., Taranukha, T., Elghazi, L., Cras-Méneur, C. and Bernal-Mizrachi, E., (2012) mTORC1 signaling and regulation of pancreatic β -cell mass. *Cell cycle* 11, 1892-1902.
- Castedo, M., Ferri, K. and Kroemer, G., (2002). Mammalian Target of Rapamycin (mTOR): Pro- and Anti-Apoptotic. *Cell Death & Differentiation*, 9(2), pp.99-100.
- Ding, Y. and Choi, M., (2014). Autophagy in diabetic nephropathy. *Journal of Endocrinology*, 224(1), pp.R15-R30.
- Erekat, N., Rababa'h, R. and Al-Jarrah, M., (2019). Overexpression of renal proapoptotic factors is attenuated subsequent to endurance exercise in Type I diabetes: An immunohistochemistry study. *Journal of Natural Science, Biology and Medicine*, 10(1), p.24.
- Eslami, H. and Mohammad Sharifi, A., (2013). Role of Bax Protein and Caspase-3 at High Glucose-Induced Apoptosis in Human Embryonic Kidney (HEK) 293 Cells. *Zahedan Journal of Research in Medical Sciences*, 15(5), pp.25-29.
- Feng, M., Li, J., Wang, J., Ma, C., Jiao, Y., Wang, Y., Zhang, J., Sun, Q., Ju, Y., Gao, L. and Zhao, Y., (2014). High glucose increases LPS-induced DC apoptosis through modulation of ERK1/2, AKT and Bax/Bcl-2. *BMC Gastroenterology*, 14(1).
- Gao, X., Kuang, H., Zou, W., Liu, X., Lin, H. and Yang, Y., (2008). The timing of re-institution of good blood glucose control affects apoptosis and expression of Bax and Bcl-2 in the retina of diabetic rats. *Molecular Biology Reports*, 36(7), pp.1977-1982.
- Gödel, M., Hartleben, B., Herbach, N., Liu, S., Zschiedrich, S., Lu, S., Debreczeni-Mór, A., Lindenmeyer, M., Rastaldi, M., Hartleben, G., Wiech, T., Forni, A., Nelson, R., Kretzler, M., Wanke, R., Pavenstädt, H., Kerjaschki, D., Cohen, C., Hall, M., Rüegg, M., Inoki, K., Walz, G. and Huber, T. (2011). Role of mTOR in podocyte function and diabetic nephropathy in humans and mice. *Journal of Clinical Investigation*, 121(6), pp.2197-2209.

- Gokalp-Ozkorkmaz, E., Kirman, G., Pekkolay, Z., Asir, F. and Deveci, E., (2018). Expression of Apoptotic Proteins Bax and Bcl-2 in Blood Cells of Type 2 Diabetic Patients. *Proceedings*, 2(25), p.1563.
- Gong, L., Liu, F., Wang, J., Wang, X., Hou, X., Sun, Y., Qin, W., Wei, S., Zhang, Y., Chen, L. and Zhang, M. (2011). Hyperglycemia induces apoptosis of pancreatic islet endothelial cells via reactive nitrogen species-mediated Jun N-terminal kinase activation. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1813(6), pp.1211-1219.
- Inoki, K., Mori, H., Wang, J., Suzuki, T., Hong, S., Yoshida, S., Blattner, S., Ikenoue, T., Rüegg, M., Hall, M., Kwiatkowski, D., Rastaldi, M., Huber, T., Kretzler, M., Holzman, L., Wiggins, R. and Guan, K. (2011). mTORC1 activation in podocytes is a critical step in the development of diabetic nephropathy in mice. *Journal of Clinical Investigation*, 121(6), pp.2181-2196.
- Jung, D., Lee, S., Kwak, S., Li, J., Kim, D., Nam, B., Kang, H., Chang, T., Park, J., Han, S., Yoo, T. and Kang, S. (2011). Apoptosis occurs differentially according to glomerular size in diabetic kidney disease. *Nephrology Dialysis Transplantation*, 27(1), pp.259-266.
- Khalfaoui, T., Basora, N. & Ouertani-Meddeb, A., (2010). Apoptotic factors (Bcl-2 and Bax) and diabetic retinopathy in type 2 diabetes. *J Mol Hist* 41, 143–152. <https://doi.org/10.1007/s10735-010-9271-9>
- Khadori, R., (2019). *Type 1 Diabetes Mellitus: Practice Essentials, Background, Pathophysiology*. [online] Emedicine.medscape.com. Available at: <https://emedicine.medscape.com/article/117739-overview#a3> [Accessed 18 Sep. 2019].
- Kubisz, P., (2015). Endothelial and platelet markers in diabetes mellitus type 2. *World Journal of Diabetes*, 6(3), p.423.
- Kulkarni, R., Mizrachi, E., Ocana, A. and Stewart, A., (2012). Human β -cell proliferation and intracellular signaling driving in the dark without a road map. *Diabetes* 61, pp.2205-2213.
- Laplante, M., and Sabatini, D.M. (2012). mTOR signaling in growth control and disease cell 149, pp.274-293.
- Mao, Z. and Zhang, W. (2018). Role of mTOR in Glucose and Lipid Metabolism. *International Journal of Molecular Sciences*, 19(7), p.2043.
- Mundel, P., (2002). Podocyte Biology and Response to Injury. *Journal of the American Society of Nephrology*, 13(12), pp.3005-3015.
- Nagata, S., (2018). *Apoptosis and Clearance of Apoptotic Cells*. pp.489-517.
- Parchwani, D. and Upadhyah, A. (2012). Diabetic nephropathy: Progression and pathophysiology. *International Journal of Medical Science and Public Health*, 1(2), p.59.

- Paul, W., Littman, D. and Yokoyama, W. (2012). *Annual review of immunology*. Palo Alto, Calif.: Annual Reviews.
- Peng, J., Li, X., Zhang, D., Chen, J., Su, Y., Smith, S. and Dong, Z. (2015). Hyperglycemia, p53, and mitochondrial pathway of apoptosis are involved in the susceptibility of diabetic models to ischemic acute kidney injury. *Kidney International*, 87(1), pp.137-150.
- Qian, Y., Feldman, E., Pennathur, S., Kretzler, M. and Brosius, F. (2008). From Fibrosis to Sclerosis: Mechanisms of Glomerulosclerosis in Diabetic Nephropathy. *Diabetes*, 57(6), pp.1439-1445.
- Sanchez-Niño, M., Fernandez-Fernandez, B., Perez-Gomez, M., Poveda, J., Sanz, A., Cannata-Ortiz, P., Ruiz-Ortega, M., Egido, J., Selgas, R. and Ortiz, A., (2015). Albumin-induced apoptosis of tubular cells is modulated by BASP1. *Cell Death & Disease*, 6(2), pp.e1644-e1644.
- Sanchez-Niño, M., Sanz, A., Lorz, C., Gnrke, A., Rastaldi, M., Nair, V., Egido, J., Ruiz-Ortega, M., Kretzler, M. and Ortiz, A., (2010). BASP1 Promotes Apoptosis in Diabetic Nephropathy. *Journal of the American Society of Nephrology*, 21(4), pp.610-621
- Tunçdemir, M. and Öztürk, M. (2016). Regulation of the Ku70 and apoptosis-related proteins in experimental diabetic nephropathy. *Metabolism*, 65(10), pp.1466-1477.
- Vergès, B. and Cariou, B., 2015. mTOR inhibitors and diabetes. *Diabetes Research and Clinical Practice*, 110(2), pp.101-108.
- Wei, P. and Szeto, C. (2019). Mitochondrial dysfunction in diabetic kidney disease. *Clinica Chimica Acta*, 496, pp.108-116.
- Walters, H. and Cox, L. (2018). mTOR Inhibitors as Broad-Spectrum Therapeutics for Age-Related Diseases. *International Journal of Molecular Sciences*, 19(8), p.2325.
- Westphal, D., Dewson, G., Czabotar, P. and Kluck, R. (2011). Molecular biology of Bax and Bak activation and action. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1813(4), pp.521-531.
- Xie, J., and Herbert, T.P. (2012). The role of mammalian target of rapamycin (mTOR) in the regulation of pancreatic beta-cell death by activation of autophagy via 5'-AMP-activated protein kinase modulation. *Endocrine* 44, pp. 87-98.
- Yamahara, K., Yasuda, M., Kume, S., Koya, D., Maegawa, H. and Uzu, T., (2013). The Role of Autophagy in the Pathogenesis of Diabetic Nephropathy. *Journal of Diabetes Research*, 2013, pp.1-9.