

## DAFTAR PUSTAKA

- Abramson, N. dan Melton, B., 2000. Leukocytosis: Basics of Clinical Assessment. *American Family Physician*, 62(9), pp. 2053–2060. PMID: 11087187
- Amatullah, H., Shan, Y., Beauchamp, B. L. *et al.*, 2017. DJ-1/PARK7 impairs bacterial clearance in sepsis. *American Journal of Respiratory and Critical Care Medicine*, 195(1), pp. 889–905. doi: 10.1164/rccm.201604-0730OC
- Angus, D. C., Linde-Zwirble, W. T., Lidicker, J. *et al.*, 2001. Epidemiology of severe sepsis in the United States: Analysis of incidence, outcome dan associated costs of care. *Critical Care Med.*, 29, pp. 1303–1310.
- Babior, B. M., 1984. Oxidants From Phagocytes: Agents of Defense and Destruction. *Blood*, 64, pp. 959–966.
- Bae, M. H., Park, S. H., Park, C. J. *et al.*, 2016. Flow cytometric measurement of respiratory burst activity and surface expression of neutrophils for septic patient prognosis. *Cytometry Part B - Clinical Cytometry*, 90(4), pp. 368–375. doi: 10.1002/cyto.b.21274.
- Baehner, R. L. dan Nathan, D. G., 1968. Quantitative Nitroblue Tetrazolium Test in Chronic Granulomatous Disease. *The New England Journal of Medicine*, 278(18), pp. 971–976. doi: 10.1056/NEJM196805022781801.
- Ballmer, P. E., 2001. Causes and Mechanism of Hypoalbuminemia. *Clinical Nutrition*, 20(3), pp. 271–273. doi: 10.1054/clnu.2001.0439
- Boomer, J. S., Green, J. M. dan Hotchkiss, R. S., 2014. The changing immune system in sepsis. *Virulence*, 5(1), pp. 45–56. doi: 10.4161/viru.26516.
- Chaplin, D. D., 2010. Overview of the Immune Response. *Journal of Allergy and Clinical Immunology*, 125(2), pp. S3-23. doi: 10.1016/j.jaci.2009.12.980.
- Choi, H. S., Kim, J. W., Cha, Y. N. *et al.*, 2006. A quantitative nitroblue tetrazolium assay for determining intracellular superoxide anion production in phagocytic cells. *Journal of Immunoassay and Immunochemistry*, 27(1), pp. 31–44. doi: 10.1080/15321810500403722.
- Danikas, D. D., Karakantza, M., Theodorou, G. L. *et al.*, 2008. Prognostic value of phagocytic activity of neutrophils and monocytes in sepsis. Correlation to CD64 and CD14 antigen expression. *Clinical and Experimental Immunology*, 154(1), pp. 87–97. doi: 10.1111/j.1365-2249.2008.03737.x.
- Ding, R., Meng, Y. dan Ma, X., 2018. The Central Role of the Inflammatory Response in Understanding the Heterogeneity of Sepsis-3. *BioMed Research International*, 2018. doi: 10.1155/2018/5086516.

- Djordjevic, D., Karakantza, M., Theodorou, G. L. *et al.*, 2018. Neutrophil-to-Lymphocyte Ratio, Monocyte-to-Lymphocyte Ratio, Platelet-to-Lymphocyte Ratio, and Mean Platelet Volume-to-Platelet Count Ratio as Biomarkers in Critically Ill and Injured Patients: Which Ratio to Choose to Predict Outcome and Nature of Bacteremia?. *Mediators of Inflammation*. Hindawi, 2018. doi: 10.1155/2018/3758068.
- Drewry, A. M., Samra, N., Skrupky, L. P. *et al.*, 2014. Persistent lymphopenia after diagnosis of sepsis predicts mortality. *Shock*, 42(5), pp. 383–391. doi: 10.1097/SHK.0000000000000234.
- Fang, F. C., 2004. Antimicrobial reactive oxygen and nitrogen species: Concepts and controversies. *Nature Reviews Microbiology*, 2(10), pp. 820–832. doi: 10.1038/nrmicro1004.
- Fang, F. C., 2011. Antimicrobial Actions of Reactive Oxygen Species. *mBio*, II(5).
- Fialkow, L., Wang, Y. dan Downey, G. P., 2007. Reactive oxygen and nitrogen species as signaling molecules regulating neutrophil function. *Free Radical Biology and Medicine*, 42(2), pp. 153–164. doi: 10.1016/j.freeradbiomed.2006.09.030.
- Fleischmann, C., Scherag, A., Adhikari, N. K. *et al.*, 2016. Assessment of global incidence and mortality of hospital-treated sepsis current estimates and limitations. *American Journal of Respiratory and Critical Care Medicine*, 193(3), pp. 259–272. doi: 10.1164/rccm.201504-0781OC.
- Forget, P., Khalifa, C., Defour, J. *et al.*, 2017. What is the normal value of the neutrophil-to-lymphocyte ratio?. *BMC Research Notes*. BioMed Central, 10(1), pp. 1–4. doi: 10.1186/s13104-016-2335-5.
- Furukawa, M., Kinoshita, K., Yamaguchi, J. *et al.*, 2019. Sepsis patients with complication of hypoglycemia and hypoalbuminemia are an early and easy identification of high mortality risk. *Internal and Emergency Medicine*. Springer International Publishing, 14(4), pp. 539–548. doi: 10.1007/s11739-019-02034-2.
- Gutierrez, M. G., 2013. Functional role(s) of phagosomal Rab GTPases. *Small GTPases*, 4(3), pp. 148–158. doi: 10.4161/sgtp.25604.
- Hampson, P., Dinsdale, R. J., Wearn, C. M. *et al.*, 2017. Neutrophil dysfunction, immature granulocytes, and cell-free DNA are early biomarkers of sepsis in burn-injured patients: A prospective observational cohort study. *Annals of Surgery*, 265(6), pp. 1241–1249. doi:10.1097/SLA.0000000000001807

- Heffernan, D. S., Monaghan, S. F., Thakkar, R. K. *et al.*, 2012. Failure to normalize lymphopenia following trauma is associated with increased mortality, independent of the leukocytosis pattern. *Critical Care*. BioMed Central Ltd, 16(1), p. R12. doi: 10.1186/cc11157.
- Hotchkiss, R. S., Monneret, G. dan Payen, D., 2013. Immunosuppression in sepsis: a novel understanding of the disorder and a new therapeutic approach. *NIH Public Access Author*, 13(3), pp. 260–268. doi: 10.1016/S1473-3099(13)70001-X.Immunosuppression.
- Hotchkiss, R. S., Monneret, G. dan Payen, D., 2013. Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy Richard. *Nature Reviews Immunology*, 13(12), pp. 862–874. doi: 10.1037/a0032811.Child.
- LaRosa, D. F. dan Orange, J. S., 2008. 1. Lymphocytes. *Journal of Allergy and Clinical Immunology*, 121(2 SUPPL. 2), pp. 364–369. doi: 10.1016/j.jaci.2007.06.016.
- Liu R, Fu S, Zhan H, dan Lucia LA., 2009. Spectrophotometric determination of superoxide anion radical with nitro blue tetrazolium General Spectroscopic Protocol to Obtain the Concentration of the Superoxide Anion Radical. *Industrial and Engineering Chemistry Research*, 48:9331-9334. doi:10.1021/ie9007826
- Liu, T., Zhang, L., Joo, D. *et al.*, 2017. NF-κB signaling in inflammation. *Signal Transduction and Targeted Therapy*, 2(April). doi: 10.1038/sigtrans.2017.23.
- Liu, X., Shen, Y., Wang, H. *et al.*, 2016. Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis: A Prospective Observational Study. *Mediators of Inflammation*. Hindawi Publishing Corporation, 2016. doi: 10.1155/2016/8191254.
- Ljungström, L. L., Karlsson, D., Pernestig, A. *et al.*, 2015. Neutrophil to lymphocyte count ratio performs better than procalcitonin as a biomarker for bacteremia and severe sepsis in the emergency department. *Critical Care*, 19(1), p. 66. doi: 10.1186/cc14146.
- Luan, Y. Y., Yao, Y. M., Xiao, X. Z. *et al.*, 2015. Insights into the apoptotic death of immune cells in sepsis. *Journal of Interferon and Cytokine Research*, 35(1), pp. 17–22. doi: 10.1089/jir.2014.0069.
- Marshall, J. C., 2005. Neutrophils in the pathogenesis of sepsis. *Crit Care Med*, 33(12 SUPPL.), pp. 12–15. doi:10.1097/01.CCM.0000186266.34541.5F

- Marshall, J. S., Warrington, R., Watson, W. *et al.*, 2018. An introduction to immunology and immunopathology. *Allergy, Asthma and Clinical Immunology*. BioMed Central, 14(s2), pp. 1–10. doi: 10.1186/s13223-018-0278-1.
- Martin, G. S., Mannino, D. M. dan Moss M., 2005. The effect of age on the development and outcome of adult sepsis. *Critical Care Medicine*, 34(1), pp. 15–21. doi:10.1097/01.CCM.0000194535.82812.BA
- Martins, P. S., Kallas, E. G., Neto, M. C. *et al.*, 2003. Upregulation of reactive oxygen species generation and phagocytosis, and increased apoptosis in human neutrophils during severe sepsis and septic shock. *Shock (Augusta, Ga.)*, 20(3), pp. 208–212. doi: 10.1097/01.shk.0000079425.52617.db.
- Mayadas, T. N., Cullere, X. dan Lowell, C. A., 2014. The Multifaceted Functions of Neutrophils. *Annual Review Pathology*, 9, pp. 181–218. doi: 10.1146/annurev-pathol-020712-164023.
- Monneret, G., Venet, F., Pachot, A. *et al.*, 2008. Monitoring immune dysfunctions in the septic patient: A new skin for the old ceremony. *Molecular Medicine*, 14(1–2), pp. 64–78. doi:10.2119/2007-00102.Monneret
- Muller Kobold, A. C., Tulleken, J. E., Zijlstra, J. G. *et al.*, 2000. Leukocyte activation in sepsis: Correlations with disease state and mortality. *Intensive Care Medicine*, 26(7), pp. 883–892. doi: 10.1007/s001340051277.
- Murphy, K. dan Weaver, C., 2016. *Janeway's Immunobiology*. 9th ed. New York: Garland Science. doi: 10.1201/b13424-12.
- Park, B. H., Fikrig, S. M. dan Smithwick, E. M., 1968. Infection and nitroblue-tetrazolium reduction by neutrophils. A diagnostic acid. *Lancet*, 2(7567), pp. 532–534. doi: 10.1016/s0140-6736(68)92406-9.
- Parkin, J. dan Cohen, B., 2001. An Overview of the Immune System. *Immunology*. *Lancet*, 347(9270), pp. 1777–1789. doi: 10.1016/S0140-6736(00)04904-7.
- Pawelec, G., Solana, R., Remarque, E. *et al.*, 1998. Impact of aging on innate immunity. *Journal of Leukocyte Biology*, 64(6), pp. 703–712. doi:10.1002/jlb.64.6.703
- Polat, G., Ugan, R. A., Cadirci, E. *et al.*, 2017. Sepsis and Septic Shock: Current Treatment Strategies and New Approaches. *Eurasian Journal of Medicine*, 49(1), pp. 53–58. doi: 10.5152/eurasianjmed.2017.17062.

- van der Poll, T., van de Veerdonk, F. L., Scicluna, B. P. *et al.*, 2017. The immunopathology of sepsis and potential therapeutic targets. *Nature Reviews Immunology*. Nature Publishing Group, 17(7), pp. 407–420. doi: 10.1038/nri.2017.36.
- Rajab, R., 2012. Hubungan kadar laktat dengan disfungsi organ multipel pada pasien sepsis di Makassar. Ph.D. Thesis, Universitas Hasanuddin, Makassar.
- Rosales, C. dan Uribe-Querol, E., 2017. Phagocytosis: A Fundamental Process in Immunity. *BioMed Research International*. Hindawi, 2017. doi: 10.1155/2017/9042851.
- Shen, X. F., Cao, K., Jiang, J. P. *et al.*, 2017. Neutrophil dysregulation during sepsis: an overview and update. *Journal of Cellular and Molecular Medicine*, 21(9), pp. 1687–1697. doi: 10.1111/jcmm.13112.
- Singer, M., Deutschman, C. S., Seymour, C. W. *et al.*, 2016. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *Jama*, 315(8), pp. 801–810. doi: 10.1001/jama.2016.0287.The.
- Sônego, F., Castanheira, F. V., Ferreira, R. G. *et al.*, 2016. Paradoxical roles of the neutrophil in sepsis: Protective and deleterious. *Frontiers in Immunology*, 7(APR), pp. 1–7. doi: 10.3389/fimmu.2016.00155.
- Thomas, E. L., Lehrer, R. I. dan Rest, R. F., 1988. Human neutrophil antimicrobial activity. *Rev Infect Dis*, 10 Suppl 2(August), pp. 450–456. doi:10.1093/cid/10.supplement\_2.s450
- Tosi, M. F., 2005. Innate immune responses to infection. *Journal of Allergy and Clinical Immunology*, 116(2), pp. 241–249. doi: 10.1016/j.jaci.2005.05.036.
- Wang, X., Zhang, G., Jiang, X. *et al.*, 2014. Neutrophil to lymphocyte ratio in relation to risk of all-cause mortality and cardiovascular events among patients undergoing angiography or cardiac revascularization: A meta-analysis of observational studies. *Atherosclerosis*. Elsevier Ltd, 234(1), pp. 206–213. doi: 10.1016/j.atherosclerosis.2014.03.003.
- Witter, A. R., Okunnu, B. M. dan Berg, R. E., 2016. The Essential Role of Neutrophils During Infection with the Intracellular Bacterial Pathogen *Listeria monocytogenes*. *Journal of Immunology*, 197(5), pp. 1557–1565. doi: 10.4049/jimmunol.1600599.

- Zador, Z., Landry, A., Cusimano, M. D. *et al.*, 2019. Multimorbidity states associated with higher mortality rates in organ dysfunction and sepsis: A data-driven analysis in critical care. *Critical Care*. Critical Care, 23(1), pp. 1–11. doi: 10.1186/s13054-019-2486-6.
- Zahorec, R., 2001. Ratio of neutrophil to lymphocyte counts — rapid and simple parameter of systemic inflammation and stress in critically ill Pomer neutrofilov a lymfocytov — rýchly a jednoduchý ukazovate systémového zápalu a stresu v kritických stavoch. *Bratislava Medical Journal*, 102(1), pp. 5–14.