

REFERENSI

American Society of Anaesthesiologist. (2017). [online] Available at: <https://www.asahq.org/standards-and-guidelines/continuum-of-depth-of-sedation-definition-of-general-anesthesia-and-levels-of-sedationanalgesia> [Accessed 8 Oct. 2019].

American Society of Anaesthesiologist. (2018). [online] Available at: <https://www.asahq.org/resources/clinical-information/asa-physical-status-classification-system> [Accessed 18 Oct. 2018].

Bajwa, S., Costi, D. and Cyna, A. (2010). A comparison of emergence delirium scales following general anesthesia in children. *Pediatric Anesthesia*, 20(8), pp.704-711.

Carolina, T., Carolina R., Barbosa, C., Sergio, P., Gonelli, R., Augusta, F. (2018). Risk factors associated with anesthesia emergence delirium in children undergoing outpatient surgery. *Brazilian Journal of Anesthesiology*, 68(2), pp. 162-167.

Cohen, I., Hannallah, R., Hummer, K. (2001). The incidence of emergence agitation associated with desflurane anesthesia in children is reduced by Fentanyl. *Anesthesia & Analgesia*, 93(1), pp. 88-91.

Cravero, J., Surgenor, S., Whalen, K. (2000). Emergence agitation in Paediatric patients after sevoflurane anesthesia and no surgery: a comparison with halothane. *Paediatric Anaesthesia*, 10(4):419-24.

Cravero, J., Rosen, H., Mervitz, D. (2015). Pediatric emergence delirium: Canadian Pediatric Anesthesiologists' experience. *Pediatric Anesthesia*, 26(2), pp.207-212.

Dahmani, S., Delivet, H. and Hilly, J. (2014). Emergence delirium in children. *Current Opinion in Anaesthesiology*, 27(3), pp.309-315.

Gooden, R., Tennant, I., James, B., Augier, R., Crawford, A., Ehikhametalor, K., Gordon, G., Harding, H. (2014). The incidence of emergence delirium and risk factors following sevoflurane use in pediatric patients for day case surgery, Kingston, Jamaica. *Revista Brasileira de Anestesiologia*, 64(6), pp.413-418

Garra, G., Adam, J., Singer, M., Breena, R., Chohan, J., Cardoz, H., Chisena, E., Henry, C. (2010). Validation of the Wong Baker FACES Pain Rating Scale in Pediatric Emergency Departement Patients. *Academic Emergency Medicine*, 17(1), pp.50

International Association for the Study of Pain (IASP). (2001). *FACES Pain Scale Revised*. [online] Available at: <https://www.iasp-pain.org/Education/Content.aspx?ItemNumber=1519>

Jin, H., Suk, K., Hee, Y., Tae, J. (2012). The influence of DNA polymorphism of multidrug resistant 1 (MDR1) on the effect of midazolam pretreatment in children. *Korean Journal of Anesthesiology*, 62(4), pp. 332.

Kanaya, A. (2015). Emergence agitation in children: risk factors, prevention, and treatment. *Journal of Anesthesia*, 30(2), pp.261-267.

Logan, K., Rich, C., Cristofaro, C., and Collins, S. (2010). Use of Propofol and Emergence Agitation in Children: A Literature Review. *AANA Journal*, 78(6), pp. 468-473.

Lee, C. (2018). Paediatric Emergence Delirium: An Approach to Diagnosis and Management in the Postanaesthesia Care Unit. *Journal of Perioperative & Critical Intensive Care Nursing*, 04(01).

Locatelli, B., Ingelmo, P., Emre, S., Meroni, V., Minardi, C., Frawley, G., Benigni, A., Di Marco, S., Spotti, A., Busi, I. and Sonzogni, V. (2012). Emergence delirium in children: a comparison of sevoflurane and desflurane anesthesia using the Paediatric Anesthesia Emergence Delirium scale. *Pediatric Anesthesia*, 23(4), pp.301-308.

Malarbi, S., Stargatt, R., Howard, K. and Davidson, A. (2011). Characterizing the behavior of children emerging with delirium from general anesthesia. *Pediatric Anesthesia*, 21(9), pp.942-950.

Mason, K. (2017). Paediatric emergence delirium: a comprehensive review and interpretation of the literature. *British Journal of Anaesthesia*, 118(3), pp.335-343.

McHugh, M. (2012). Interrater reliability: the Kappa statistic. *Biochemia Medica*, 22(3), pp. 276-282.

Nair, S., Wolf, A. (2017). Emergence Delirium after Pediatric anaesthesia: New strategies in avoidance and treatment. *BJA Education*. 33(4), pp. 107-112.

O'Regan, N., Fitzgerald, J., Timmons, S., O'Connell, H. and Meagher, D. (2013). Delirium: A key challenge for perioperative care. *International Journal of Surgery*, 11(2), pp.136-144.

Paul, A., Mizell, M., Thomas, S., Goldfinger, M., Florentino, I. (2004). Assessment of risk factors for emergence distress and postoperative behavioural changes in children following general anaesthesia. *Pediatric Anesthesia*, 14(1), pp. 235-240.

Pickard, A., Davies, P., Birnie, K., and Beringer, R. (2014). Systematic review and meta-analysis of the effect of intraoperative α -adrenergic agonists on postoperative behaviour in children. *British Journal of Anaesthesia*, 112(6), pp.982-990.

Reduque, L., Verghese, S. (2013). Pediatric emergence delirium. *Continuing Education in Anaesthesia Critical Care and Pain: British Journal of Anaesthesia*, 13(2), pp.39-41.

Reynolds, T., Sankaran, S., Chimbira, W., Phan, T., Nafiu, O. (2017). Severe Obesity and Sleep Disordered Breathing as risk factors for Emergence Agitation in pediatric ambulatory surgery. *Journal of PeriAnesthesia Nursing*, 33(3), pp.304-311.

Rim, C., Kim, J., Hong, J., Park, S., Lee, J., Chung, C. (2016). Risk factors of emergence agitation after general anesthesia in adult patients. *Anesthesia Pain Medications*, 11(1), pp. 410-416.

Sinha, A., Sood, J. (2012). Caudal block and emergence delirium in pediatric patients. *Saudi Journal of Anaesthesia*, 6(4), p. 403.

Slade, T., Andrews, G. (2001). DSM-IV and ICD-10 generalized anxiety disorder: discrepant diagnoses and associated disability. *Social Psychiatry and Psychiatric Epidemiology*, 36(1), pp. 45-51.

Somaini, M., Engelhardt, T., Fumagalli, R., Ingelmo, P. (2016). Emergence delirium or pain after anaesthesia-how to distinguish between the two in young children: a retrospective analysis of observational studies. *British Journal of Anaesthesia*, 116(3), pp. 377-383.

Vlajkovic, G., Sindjelic, R. (2007). Emergence Delirium in Children: Many Questions, Few Answers. *Anesthesia & Analgesia*, 104(1), pp.84-91.

Yu, D., Chai, W., Sun, X., Yao, L. (2010). Emergence agitation: risk factors in 2000 patients. *Canadian Journal of Anaesthesia*, 57(9), pp. 843-848.