

Daftar Pustaka

- Abdelrahman TM. Prognostic value of T peak-to-end interval for risk stratification after acute myocardial infarction. *The Egyptian Journal of Critical Care Medicine*. 2014 Apr 1;2(1):19-27.
- Achmad C, Iqbal M, Karwiky G, Prameswari HS, Febrianora M. 2020. T-Peak to T-End Improvements After Beta-Blocker Administration in Peripartum Cardiomyopathy Patients. *Cardiol Res*
- Akboğa MK, Balcı KG, Yılmaz S, Aydın S, Yayla Ç, Ertem AG, Ünal S, Balcı MM, Balbay Y, Aras D, Topaloğlu S. Tp-e interval and Tp-e/QTc ratio as novel surrogate markers for prediction of ventricular arrhythmic events in hypertrophic cardiomyopathy. *Anatolian journal of cardiology*. 2017 Jul;18(1):48.
- Alonso JL, Martínez P, Vallverdú M, Cygankiewicz I, Pitzalis MV, Genís AB, Cinca J, Rizzon P, Caminal P, Zareba W, De Luna AB. Dynamics of ventricular repolarization in patients with dilated cardiomyopathy versus healthy subjects. *Annals of noninvasive electrocardiology*. 2005 Apr;10(2):121-8.
- Antzelevitch, C., Viskin, S., Shimizu, W., Yan, Gan-Xin., Kowey, P., Zhang, Li., Sicouri, S., Di Diego, J.M., Burashnikov, A. 2007. Does Tpeak-Tend provide an index of transmural dispersion of repolarization? *Heart Rhythm*;4(8): 1114-1119.
- Bacon BR. Sirosis dan penyulitnya. In: Longo DL, Fauci AS, editors. *Harrison gastroenterologi dan hepatologi*. Jakarta: EGC; 2010 .p. 374-86.
- Baldassarre M, Giannone FA, Napoli L, Tovoli A, Ricci CS, Tufoni M, et.al. The endocannabinoid system in advanced liver cirrhosis: Pathophysiological implication and future perspectives. *Liver Int*. 2013; 33: 1298–308. doi: 10.1111/liv.12263.
- Barr, C. S., Naas, A. A., Fenwick, M., & Struthers, A. D. (1997). Enalapril reduces QTc dispersion in mild congestive heart failure secondary to coronary artery disease. *The American journal of cardiology*, 79(3), 328-333.
- Bicca J, Jarske L, Silva T, Gismondi R, Mocarzel L, Lanzieri P. Cirrhotic cardiomyopathy. *International Journal of Cardiovascular Sciences*. 2016 Feb 21;29(2):139-48.
- Bonilla-Palomas JL, Gámez-López AL, Moreno-Conde M, López-Ibáñez MC, Anguita-Sánchez M, de la Sacristana ÁG, García-Catalán F, Villar-Ráez A. Hypoalbuminemia in acute heart failure patients: causes and its impact on hospital and long-term mortality. *Journal of cardiac failure*. 2014 May 1;20(5):350-8.
- Bousier, J., Casbron, E., Tropet, A.L., & Piette, C. 2009. *Comparison and improvement of MELD and Child Pugh score accuracies for the prediction for 6 month mortality in cirrhotic patients*. *Journal of Clinical Gastroenterology*; 43(6); 500-5.

- Burak, A., Yüksel, M., Yayla, Ç., Kırbaş, Ö., Ünal, S., Ertem, A. G., ... & Kayaçetin, E. (2017). Parameters of ventricular repolarization in patients with autoimmune hepatitis. *Turk Kardiyol Dern Ars*, 45(4), 333-338.
- Chou R, Wasson N. Blood tests to diagnose fibrosis or cirrhosis in patients with chronic hepatitis C virus infection: a systematic review. *Ann Intern Med*. 2013;158:807-20.
- Ciobanu, A., Tse, G., Liu, T., Deaconu, M. V., Gheorghe, G. S., Ilieşiu, A. M., & Nanea, I. T. (2017). Electrocardiographic measures of repolarization dispersion and their relationships with echocardiographic indices of ventricular remodeling and premature ventricular beats in hypertension. *Journal of geriatric cardiology: JGC*, 14(12), 717.
- Demir, C., & Demir, M. (2013). Evaluation of Tp-e interval and Tp-e/QT ratio in patients with chronic hepatitis B. *Prague Med Rep*, 114(4), 239-45.
- Demirtas, D., Sumbul, H. E., Bulut, A., Demirtas, A. O., Gulumsek, E., Koca, H. & Koc, M. (2020). Tp-e interval, Tp-e/QT and Tp-e/QTc ratio in hypertensive patients with primary aldosteronism. *Clinical and Experimental Hypertension*, 42(1), 93-98.
- Dimmitt, S. B., Stampfer, H. G., & Warren, J. B. (2014). β -adrenoceptor blockers valuable but higher doses not necessary. *British journal of clinical pharmacology*, 78(5), 1076-1079.
- Dinshaw, L., Münch, J., Dickow, J., Lezius, S., Willems, S., Hoffmann, B. A., & Patten, M. (2018). The T-peak-to-T-end interval: a novel ECG marker for ventricular arrhythmia and appropriate ICD therapy in patients with hypertrophic cardiomyopathy. *Clinical Research in Cardiology*, 107(2), 130-137.
- Dumont CA, Monserrat L, Soler R, Rodríguez E, Fernandez X, Peteiro J, Bouzas A, Bouzas B, Castro-Beiras A. Interpretation of electrocardiographic abnormalities in hypertrophic cardiomyopathy with cardiac magnetic resonance. *European heart journal*. 2006 Jun 14;27(14):1725-31.
- Fede G, Privitera G, Tomaselli T, Spadaro L, Purrello F. Cardiovascular dysfunction in patients with liver cirrhosis. *Ann Gastroenterol*. 2014; 27(5): 1-10.
- Fede, G., Privitera, G., Tomaselli, T., Spadaro, L., & Purrello, F. (2015). Cardiovascular dysfunction in patients with liver cirrhosis. *Annals of Gastroenterology: Quarterly Publication of the Hellenic Society of Gastroenterology*, 28(1), 31.
- Ferrucci, A., Canichella, F., Battistoni, A., Palano, F., Francia, P., Ciavarella, G.M., Volpe, M., Tocci, G. 2015. A novel electrocardiographic T-wave measurement (Tp-Te interval) as a predictor of heart abnormalities in hypertension: a new opportunity for first-line electrocardiographic evaluation. *The Journal of Clinical Hypertension*, 17, 441-449.

- Fouad YM, Yehia R. Hepato-cardiac disorders. *World journal of hepatology*. 2014 Jan 27;6(1):41
- Funck-Brentano C, Jaillon P. Rate-corrected QT interval: techniques and limitations. *American Journal of Cardiology*. 1993 Aug 26;72(6):B17-22.
- Gupta, P., Patel, C., Patel, H., Narayanaswamy, S., Malhotra, B., Green, J.T., Yan, G.X. 2008. Tp-e/QT ratio as an index of arrhythmogenesis. *Journal of Electrocardiology*, 41(6), 567–574.
- Han, J. dan Moe, G. 1964. Nonuniform recovery of excitability in ventricular muscle. *Circulation research* 14:44-60.
- Icli, A., Kayrak, M., Akilli, H., Aribas, A., Coskun, M., Ozer, S. F., Ozdemir, K. 2015. Prognostic value of Tpeak-Tend interval in patients with acute pulmonary embolism. *BMC cardiovascular disorders*, 15, 99.
- Karaagac K, Emul A, Tenekecioglu E, Agca FV, Ozluk OA, Tutuncu A, Yontar OC, Yilmaz M. The Effects of Metabolic Syndrome on TpTe Interval and TpTe/QT Ratio in Patients with Normal Coronary Arteries. *The Eurasian journal of medicine*. 2014 Oct;46(3):182.
- Lanzieri, P. G., Gismondi, R. A., Chimelli, M. D. C. A. R., Cysne, R. P., Guaraná, T., Mesquita, C. T., & Mocarzel, L. O. (2017). Cirrhotic patients with Child-Pugh C have longer QT intervals. *International Journal of Cardiovascular Sciences*, 30(6), 496-503.
- Liu H, Ma Z, Lee SS. Contribution of nitric oxide to the pathogenesis of cirrhotic cardiomyopathy in bile duct-ligated rats. *Gastroenterology* 2000; 118(5): 937-44.
- LŐORINCZ, I., Matyus, J., Zilahi, Z., Kun, C., KARÁNYI, Z., & KAKUK, G. (1999). QT dispersion in patients with end-stage renal failure and during hemodialysis. *Journal of the American Society of Nephrology*, 10(6), 1297-1302.
- Ma Z, Lee SS, Meddings JB. Effects of altered cardiac membrane fluidity on beta-adrenergic receptor signalling in rats with cirrhotic cardiomyopathy. *J Hepatol*. 1997; 26(4): 904-12.
- Mallat A, Teixeira-Clerc F, Deveaux V, Manin S, Lotersztajn S.. The endocannabinoid system as a key mediator during liver diseases: New insights and therapeutic openings. *Br J Pharmacol*. 2011; 163(7): 1432–40. doi: 10.1111/j.1476-5381.2011.01397.x.
- Mesquita, E. T., Deus, F. C. C. D., Guedes, C. R., Maia, E. R., Subieta, C. G. P., Villacorta, H., ... & Romeo, L. J. M. (1999). Effects of propranolol on the QT dispersion in congestive heart failure. *Arquivos brasileiros de cardiologia*, 73(3), 295-298.
- Møller S, Henriksen JH. Cardiovascular complications of cirrhosis. *Gut* 2008; 57: 268-78.
- Møller S, Henriksen JH. Cirrhotic cardiomyopathy: a pathophysiological review of circulatory dysfunction in liver disease. *Heart*. 2002 Jan 1;87(1):9-15.

- Mozos, Ioana. "Arrhythmia risk in liver cirrhosis." *World journal of hepatology* 7.4 (2015): 662.
- Mozos, I.,Serban, C. 2011. The relation between QT interval and T-wave variables in hypertensive patients. *Journal of Pharmacy and Bioallied Sciences*, 3, 339.
- Nurdjanah S. Sirosis hati. In: Sudoyo AW, Setiyohadi B, Alwi I, editors. Buku ajar ilmu penyakit dalam. 4th ed. Jakarta: Pusat Penerbitan Ilmu Penyakit Dalam FKUI; 2006. p. 443-6.
- Ophthof, T., Coronel, R., Wilms-Schopman, F.J., Plotnikov, A.N., Shlapakova, I.N., Danilo, P.Jr,m Rosen, M.R., Jansen, M.J. 2007. Dispersion of repolarization in canine ventricle and the electrographic T wave: Tp-e interval does not reflect transmural dispersion. *Heart rhythm*;4: 341-348.
- Ozcan, E.E., Szilagyi, S., Sallo, Z., Molnar, L., Zima, E., Szeplaki, G., Osztheimer, I., Ozturk, A., Merkely, B., Geller, L. 2015. Comparison of the effect of epicardial and endocardial cardiac resynchronization therapy on transmural dispersion of repolarization. *Pace*;38(9):1099-1105.
- Padmanabhan, S., Silvet, H., Amin, J.,Pai, R. G. 2003. Prognostic value of QT interval and QT dispersion in patients with left ventricular systolic dysfunction: results from a cohort of 2265 patients with an ejection fraction of $\leq 40\%$. *American heart journal*, 145, 132-138.
- Panikkath R, Reinier K, Uy-Evanado A, Teodorescu C, Hattenhauer J, Mariani R, Gunson K, Jui J, Chugh SS. Prolonged Tpeak-to-tend interval on the resting ECG is associated with increased risk of sudden cardiac death. *Circulation: Arrhythmia and Electrophysiology*. 2011 Aug;4(4):441-7.
- Peters,M.G., & Carbone, J.V. 2013. *Evaluation and prognosis of patients with cirrhosis*. Hepatology Research, San Fransisco.
- Poernomo, Tiara. 2015. Peran Skor Child Pugh dan Sor Model of End Stage Liver Disease. Kedokteran Klinis Universitas Gadjah Mada. Yogyakarta
- Ruiz-del-Árbol L, Serradilla R. Cirrhotic cardiomyopathy. *World J Gastroenterol*. 2015; 21(41): 11502-21.
- Sakata K, Shimizu M, Ino H, Yamaguchi M, Terai H, Hayashi K, Kiyama M, Hayashi T, Inoue M, Mabuchi H. QT dispersion and left ventricular morphology in patients with hypertrophic cardiomyopathy. *Heart*. 2003 Aug 1;89(8):882-6.
- Saksana, R. A., Bayupurnama, P., Indrarti, F., Ratnasari, N., Maduseno, S., Triwikatmani, C., & Nurdjanah, S. (2012). Correlation Between the Severity of Liver Cirrhosis (Chil-Pugh Score) and QTc Interval Prolongation. *Indonesian Journal of Gastroenterology, Hepatology, and Digestive Endoscopy*, 13(3), 157-160.
- Salgado AA, Barbosa PR, Ferreira AG, Reis CA, Terra C. Prognostic Value of a New Marker of Ventricular Repolarization in Cirrhotic Patients. *Arquivos brasileiros de cardiologia*. 2016 Dec;107(6):523-31.

- Savelieva, I., Yap, Y. G., YI, G., Guo, X., Camm, A. J., & Malik, M. (1998). Comparative reproducibility of QT, QT peak, and T peak-T end intervals and dispersion in normal subjects, patients with myocardial infarction, and patients with hypertrophic cardiomyopathy. *Pacing and clinical electrophysiology*, 21(11), 2376-2381.
- Scaglione, Steven, *et al.* "The epidemiology of cirrhosis in the United States." *Journal of clinical gastroenterology* 49.8 (2015): 690-696.
- Sherlock, S., & Dooley, J. 2007. *Diseases of liver and billiary system*, United States.
- Sicouri, S. dan Antzelevitch, C. 1991. A subpopulation of cells with unique electrophysiological properties in the deep subepicardium of the canine ventricle. The M cell. *Circulation Research*, 68(6), 1729–1741
- Smetana, P., Schmidt, A., Zabel, M., Hnatkova, K., Franz, M., Huber, K., Malik, M. 2011. Assessment of repolarization heterogeneity for prediction of mortality in cardiovascular disease: peak to the end of the T wave interval and nondipolar repolarization components. *Journal of Electrocardiology*, 44(3), 301–8.
- Soga, K., Tomikashi, K., Miyawaki, K., Fukumoto, K., Wakabayashi, N., Konishi, H., Mitsufuji, S., Yoshida, N., Kokura, S., Naito, Y., Kataoka, K., & Yoshikawa, T. 2009. MELD score, Child Pugh score, and decreased albumin as risk factors for gastric variceal bleeding. *Hepatogastroenterology*; 56(94-95):1552-6.
- Sulaiman, Akbar, Lesmana dan Noer. 2007. *Buku Ajar Ilmu Penyakit Hati*. Jakarta: Jayabadi
- Tokatli, A., Kilicaslan, F., Alis, M., Yiginer, O., Uzun, M. 2016. Prolonged Tp-e interval, Tp-e/QT ratio and Tp-e/QTc ratio in patients with type 2 diabetes mellitus. *Endocrine Metab*;31(1):105-112.
- Tsochatzis EA, Bosch J, Burroughs AK. Liver cirrhosis. *The Lancet*. 2014 May 17;383(9930):1749-61.
- Tuttolomondo, A., Buttà, C., Casuccio, A., Di Raimondo, D., Serio, A., D'Aguanno, G., ... & Cirrincione, A. (2015). QT Indexes in cirrhotic patients: relationship with clinical variables and potential diagnostic predictive value. *Archives of Medical Research*, 46(3), 207-213.
- Viitasalo, M., Oikarinen, L., Swan, H., Väänänen, H., Järvenpää, J., Hietanen, H., ... & Toivonen, L. (2006). Effects of beta-blocker therapy on ventricular repolarization documented by 24-h electrocardiography in patients with type 1 long-QT syndrome. *Journal of the American College of Cardiology*, 48(4), 747-753.
- Walley KR, McDonald TE, Wang Y, Dai S, Russell JA. Albumin resuscitation increases cardiomyocyte contractility and decreases nitric oxide synthase II expression in rat endotoxemia. *Critical care medicine*. 2003 Jan 1;31(1):187-94.

- Wiese S, Hove JD, Bendtsen F, Møller S. Cirrhotic cardiomyopathy: Pathogenesis and clinical relevance. *Nat Rev Gastroenterol Hepatol*. 2014; 11: 177-86. doi: 10.1038/nrgastro.2013.210.
- Wong F. Cirrhotic cardiomyopathy. *Hepatol Int*. 2009; 3: 294-304.
- Xia, Y., Liang, Y., Kongstad O., Liao, Q., Holm, M., Olsson, B., Yuan, S. 2005. In vivo validation of the coincidence of the peak and end of T wave with full repolarization of the epicardium and endocardium in swine. *Heart Rhythm*;2:162-169.
- Xue, J., Gao, W., Chen, Y., Han, X. 2008. Study of repolarization heterogeneity and electrocardiographic morphology with a modeling approach. *Journal of Electrocardiology*, 41(6), 581–587.
- Yan GX, Antzelevitch C. Cellular basis for the normal T wave and the electrocardiographic manifestations of the long-QT syndrome. *Circulation*. 1998;98:1928–1936.
- Yan, G.X. dan Martin, J. 2003. Electrocardiographic T wave: a symbol of transmural dispersion of repolarization in the ventricles. *Journal of Cardiovascular Electrophysiology*, 14(6), 639–40.
- Yontar, O. C., Karaagac, K., Tenekecioglu, E., Tutuncu, A., Karakus, A., Melek, M. 2014. T wave peak to T wave end interval is prolonged in patients with atrioventricular nodal reentry. *Biomedical Research*, 25.
- Zaidi M, Robert A, Fesler R, Derwael C, Brohet C. Dispersion of ventricular repolarization in hypertrophic cardiomyopathy. *Journal of electrocardiology*. 1996 Jan 1;29:89-94.
- Zaky A, Lang JD. Cirrhosis-associated cardiomyopathy. *J Anesth Clin Res*. 2012; 3(12): 1-7.
- Zambruni ,A., Trevisani ,F., Di Micoli ,A., *et al*. 2008.Effect of chronic betablockade on QT interval in patients with liver cirrhosis. *Journal of Hepatology* ;48:415-21.
- Zardi EM, Zardi DM, Chin D, Sonnino C, Dobrina A, Abbate A. Cirrhotic cardiomyopathy in the pre-and post-liver transplantation phase. *Journal of cardiology*. 2016 Feb 1;67(2):125-30.
- Zardi EM, Abbate A, Zardi DM, Dobrina A, Margiotta D, Van Tassel BW, Afeltra A, Sanyal AJ. Cirrhotic cardiomyopathy. *Journal of the American College of Cardiology*. 2010 Aug 10;56(7):539-49.
- Zhao, Z., Yuan, Z., Ji, Y., Wu, Y., Qi, Y. 2010. Left ventricular hypertrophy amplifies the QT, and Tp-e intervals and the Tp-e/QT ratio of left chest ECG. *Journal of biomedical research*, 24, 69-72.