

DAFTAR PUSTAKA

- Admani, S., Feldstein, S., M Gonzalez, E. and Friedlander, S. (2014) 'Beta blockers: An innovation in the treatment of infantile hemangiomas', *The Journal of clinical and aesthetic dermatology*, 7, pp. 37–45.
- Akaishi, S., Akimoto, M., Ogawa, R. and Hyakusoku, H. (2008) 'The relationship between keloid growth pattern and stretching tension: Visual analysis using the finite element method', *Annals of Plastic Surgery*, 60(4), pp. 445–451.
- Akaishi, S., Koike, S., Dohi, T., Kobe, K., Hyakusoku, H. and Ogawa, R. (2012) 'Nd:YAG Laser Treatment of Keloids and Hypertrophic Scars.', *Eplasty*, 12, p. e1.
- Alster, T. S. and Handrick, C. (2000) 'Laser treatment of hypertrophic scars, keloids, and striae.', *Seminars in cutaneous medicine and surgery*, 19(4), pp. 287–292.
- Andrea, F. D. and Ferraro, G. (2002) 'Prevention and Treatment of Keloids with Intralesional Verapamil', *dermatology*, pp. 60–62.
- Andrews, J. P., Marttala, J., Macarak, E., Rosenbloom, J. and Uitto, J. (2016) 'Keloids: The paradigm of skin fibrosis - Pathomechanisms and treatment', *Matrix Biology*. Elsevier B.V., 51, pp. 37–46.
- Ardekani, G. S. (2009) *Treatment of a Postburn Keloid Scar with Topical Captopril: Report of the First Case*.
- Arima, J., Huang, C., Rosner, B. and Akaishi, S. (2015) 'Hypertension : a systemic key to understanding local keloid severity', pp. 213–221.
- Ariwibowo, L., Budiyanto, A. and Danarti, R. (2016) 'Efikasi Terapi Topikal Kortikosteroid Ultrapoten, Solusio, dan Gel Timolol Maleat 0,5% Terhadap Hemangioma Infantil Superfisial', 17(6).
- Arno, A. I., Gauglitz, G. G., Barret, J. P. and Jeschke, M. G. (2014) 'Up-to-date approach to manage keloids and hypertrophic scars: A useful guide', *Burns*, 40(7), pp. 1255–1266.
- Atiyeh, B. S. (2007) 'Nonsurgical management of hypertrophic scars: Evidence-based therapies, standard practices, and emerging methods', *Aesthetic Plastic Surgery*, 31(5), pp. 468–492.
- Bainbridge, P. (2014) 'Wound healing and the role of fibroblasts', *Journal of Wound Care*, 22(8), pp. 407–412.

- Bayat, A., Arscott, G., Ollier, W. E. R., Grouther, D. A. M. and Ferguson, M. W. J. (2005) 'Keloid disease : clinical relevance of single versus multiple site scars', *The British Association of Plastic Surgeons*, pp. 28–37.
- Berman, B. and Duncan, M. R. (1988) 'Short-term keloid treatment in vivo with human interferon alfa-2b results in a selective and persistent normalization of keloidal fibroblast collagen , glycosaminoglycan , and collagenase production in vitro Patient treatment', *Journal of the American Academy of Dermatology*. American Academy of Dermatology, Inc, 21(4), pp. 694–702.
- Biernacka, A., Dobaczewski, M. and Frangogiannis, N. G. (2015) 'TGF- β signaling in fibrosis', *Growth Factors*, 29(5), pp. 196–202.
- Bock, O., Yu, H., Zitron, S., Bayat, A., Ferguson, M. W. J. and Mrowietz, U. (2005) 'Studies of transforming growth factors beta 1-3 and their receptors I and II in fibroblast of keloids and hypertrophic scars', *Acta Dermato-Venereologica*, 85(3), pp. 216–220.
- Brien, O. L., Dj, J., Brien, L. O. and Jones, D. J. (2013) 'Silicone gel sheeting for preventing and treating hypertrophic and keloid scars (Review) Silicone gel sheeting for preventing and treating hypertrophic and keloid scars', (9), pp. 3–5.
- Brown, J. J. (2008) 'Positive association of HLA-DRB1 * 15 with keloid disease in Caucasians', *International Journal of Immunogenetics*, pp. 303–307.
- Chantasart, D., Hao, J. and Li, S. K. (2013) 'Evaluation of skin permeation of β -blockers for topical drug delivery', *Pharmaceutical Research*, 30(3), pp. 866–877.
- Cheng, W., Yan-hua, R., Fang-gang, N. and Guo-an, Z. (2011) 'The content and ratio of type I and III collagen in skin differ with age and injury', *African Journal of Biotechnology*, 10(13), pp. 2524–2529.
- Cox, T. R. *et al.* (2013) 'LOX-mediated collagen crosslinking is responsible for fibrosis-enhanced metastasis', *Cancer Research*, 73(6), pp. 1721–1732.
- Craig, R. D. P., Schofield, J. D. and Jackson, S. S. (1975) 'Collagen Biosynthesis in Normal Human Skin, Normal and Hypertrophic Scar and Keloid', *European Journal of Clinical Investigation*, 5(1), pp. 69–74.
- Cruz-Korchin, N. (1996) 'Effectiveness of silicone sheets in the prevention of hypertrophic breast scars.', *Ann Plast Surg*, 37, pp. 345–348.

- Darougheh, A., Asilian, A. and Shariati, F. (2014) 'Intralesional triamcinolone alone and in combination with 5-fluorouracil for the treatment of keloid and hypertrophic scars', *Journal of the Pakistan Medical Association*, 64(9), pp. 1003–1007.
- El-aleem, S. A., Abdelwahab, S., Mohamed, N. and Osman, S. (2017) 'Journal of Cytology & Histology Co-upregulation of Transforming Growth Factor Beta-1 and Nitric Oxide Synthase in Keloid by Comparison to Normal Human Skin - A Possible Role for TGF β 1 and NOS in Pathogenesis of Keloid', 8(4), pp. 1–8.
- Erick A, Mafong, M. and Robin Ashinoff, M. (2000) 'Treatment of Hypertrophic Scars and Keloids', *Aesthetic surgery journal*, (April), p. 116.
- España, A., Solano, T. and Quintanilla, E. (2001) 'Bleomycin in the Treatment of Keloids and Hypertrophic Scars by Multiple Needle Punctures', *Dermatologic Surgery*, 27(1).
- Fiala, O. *et al.* (2019) 'Incidental use of beta-blockers is associated with outcome of metastatic colorectal cancer patients treated with bevacizumab-based therapy: A single-institution retrospective analysis of 514 patients', *Cancers*, 11(12), pp. 1–13.
- Finsson, K. W., McLean, S., Di Guglielmo, G. M. and Philip, A. (2013) 'Dynamics of Transforming Growth Factor Beta Signaling in Wound Healing and Scarring', *Advances in Wound Care*, 2(5), pp. 195–214.
- Fitzpatrick, R. E. (1999) 'Treatment of inflamed hypertrophic scars using intralesional 5-FU', *Dermatologic Surgery*, 25(3), pp. 224–232.
- Freshney, R. I. (2000) 'Quantitation', in *culture of animal cells : a manual basic of technique*. fourth, pp. 309–327.
- Gauglitz, G. G. (2013a) 'Management of keloids and hypertrophic scars : current and emerging options', *Clinical, Cosmetic and Investigational Dermatology*, (6), pp. 103–114.
- Gauglitz, G. G. (2013b) 'Management of keloids and hypertrophic scars: Current and emerging options', *Clinical, Cosmetic and Investigational Dermatology*, 6, pp. 103–114.
- Gauglitz, G. and Korting, H. (2011a) 'Hypertrophic Scarring and Keloids: Pathomechanisms and Current and Emerging Treatment Strategies', *Molecular Medicine*, 17(1–2), p. 1.
- Gauglitz, G. and Korting, H. (2011b) 'Hypertrophic Scarring and Keloids:

Pathomechanisms and Current and Emerging Treatment Strategies’,
Molecular Medicine, 17(1–2), p. 1.

Gelse, K., Pöschl, E. and Aigner, T. (2003) ‘Collagens - Structure, function, and biosynthesis’, *Advanced Drug Delivery Reviews*, 55(12), pp. 1531–1546.

Gira, A. K., Brown, L. F., Washington, C. V., Cohen, C. and Arbiser, J. L. (2004) ‘Keloids demonstrate high-level epidermal expression of vascular endothelial growth factor’, *Journal of the American Academy of Dermatology*, 50(6), pp. 850–853.

Gomolak, J. R. and Didion, S. P. (2014) ‘Angiotensin II-induced endothelial dysfunction is temporally linked with increases in interleukin-6 and vascular macrophage accumulation’, *Frontiers in Physiology*, 5(October), pp. 1–10.

Har-Shai, Y., Amar, M. and Sabo, E. (2003) ‘Intralesional cryotherapy for enhancing the involution of hypertrophic scars and keloids’, *Plastic and Reconstructive Surgery*, 111(6), pp. 1841–1852.

Hu, J. Z. (2010) *Assessment of ascorbic acid effects on the properties of cell-derived tissue rings*. woechester polytechnic institute.

Huang, C., Murphy, G. F., Akaishi, S. and Ogawa, R. (2013a) ‘Keloids and hypertrophic scars: Update and future directions’, *Plastic and Reconstructive Surgery*, 1(4), pp. 1–7.

Huang, C., Murphy, G. F., Akaishi, S. and Ogawa, R. (2013b) ‘Keloids and hypertrophic scars: Update and future directions’, *Plastic and Reconstructive Surgery*, 1(4), pp. 1–7.

Huang, L., Cai, Y. J., Lung, I., Leung, B. C. S. and Burd, A. (2013) ‘A study of the combination of triamcinolone and 5-fluorouracil in modulating keloid fibroblasts in vitro *’, *British Journal of Plastic Surgery*. Elsevier Ltd, 66(9), pp. e251–e259.

Huang, L., Wong, Y. P., Cai, Y. J., Lung, I., Leung, C. S. and Burd, A. (2010) ‘Low-dose 5-fluorouracil induces cell cycle G2 arrest and apoptosis in keloid fibroblasts’, *British Journal of Dermatology*, 163(6), pp. 1181–1185.

Ibrahim, D. G., Ko, J. A., Iwata, W., Okumichi, H. and Kiuchi, Y. (2019) ‘An in vitro study of scarring formation mediated by human Tenon fibroblasts: Effect of Y-27632, a Rho kinase inhibitor’, *Cell Biochemistry and Function*, 37(2), pp. 113–124.

- Jagadeesan, J. and Bayat, A. (2007) 'Transforming growth factor beta (TGF β) and keloid disease', *International Journal of Surgery*, 5(4), pp. 278–285.
- Johnston, E. F. and Gillis, T. E. (2017) 'Transforming growth factor beta-1 (TGF- β 1) stimulates collagen synthesis in cultured rainbow trout cardiac fibroblasts', *The Journal of Experimental Biology*, 220(14), pp. 2645–2653.
- Jones, C. D., Guiot, L., Samy, M., Gorman, M. and Tehrani, H. (2015) 'The use of chemotherapeutics for the treatment of keloid scars', *Dermatology Reports*, 7(2), pp. 15–19.
- Ko, C. J. (2012) 'Dermal Hypertrophies and Benign Fibroblastic/Myofibroblastic Tumors', in LOWELL A. GOLDSMITH, MD, M., STEPHEN I. KATZ, MD, P., BARBARA A. GILCHREST, M., AMY S. PALLER, M., DAVID J. LEFFELL, M., and KLAUS WOLFF, MD, F. (eds) *Fitzpatrick's Dermatology in General Medicine*. Eighth Edi, pp. 707–710.
- Krishnaswamy, V. R., Lakra, R. and Korrapati, P. S. (2014) 'Keloid collagen–cell interactions: structural and functional perspective', *RSC Advances*, 4(45), p. 23642.
- 'Recent understandings of biology, prophylaxis and treatment strategies for hypertrophic scars and keloids', *International Journal of Molecular Sciences*, 19(3).
- Lee, S. Y. and Park, J. (2015) 'Postoperative electron beam radiotherapy for keloids: Treatment outcome and factors associated with occurrence and recurrence', *Annals of Dermatology*, 27(1), pp. 53–58.
- Van Leeuwen, M. C. E., Bulstra, A. E. J., Ket, J. C. F., Ritt, M. J. P. F., Van Leeuwen, P. A. M. and Niessen, F. B. (2015) 'Intralesional cryotherapy for the treatment of keloid scars: Evaluating effectiveness', *Plastic and Reconstructive Surgery - Global Open*, 3(6), pp. 1–9.
- Li, Z., Liu, Q., Wang, X. and Luan, J. (2017) 'The Characteristics of Blood Supply and Tissue Hypoxia in Pathological Scars', *Chinese Medical Sciences Journal*. Chinese Academy Medical Sciences, 32(2), pp. 113–118.
- Liu, C. *et al.* (2014) 'Activation of the AT1R/HIF-1 α /ACE Axis Mediates Angiotensin II-Induced VEGF Synthesis in Mesenchymal Stem Cells', *BioMed Research International*, 2014, pp. 1–10.
- Liu, Y. *et al.* (2016) 'TGF- β 1 promotes scar fibroblasts proliferation and

- transdifferentiation via up-regulating MicroRNA-21', *Scientific Reports*. Nature Publishing Group, 6(August), pp. 1–9.
- Longley, D. B., Harkin, D. P. and Johnston, P. G. (2003) '5-Fluorouracil: Mechanisms of action and clinical strategies', *Nature Reviews Cancer*, 3(5), pp. 330–338.
- Lu, Y. *et al.* (2010) 'Phosphatidylinositol-3-kinase/Akt regulates bleomycin-induced fibroblast proliferation and collagen production', *American Journal of Respiratory Cell and Molecular Biology*, 42(4), pp. 432–441.
- Marnaros, A. G., Norris, A. J. E. C., Watanabe, S., Reichenberger, E. and Å, B. R. O. (2004) 'Genome Scans Provide Evidence for Keloid Susceptibility Loci on Chromosomes 2q23 and 7p11', *The Society for Investigative Dermatology*, pp. 1126–1132.
- Mayasari, I. (2018) *pengaruh timolol pada proliferasi biakan fibroblas keloid*. Gadjah Mada.
- Meaume, S., Le Pillouer-Prost, A., Richert, B., Roseeuw, D. and Vadoud, J. (2014) 'Management of scars: Updated practical guidelines and use of silicones', *European Journal of Dermatology*, 24(4), pp. 435–443.
- de Mesquita, C. J. G. (2010) 'About strawberry, crab claws, and the Sir James Black's invention. Hypothesis: Can we battle keloids with propranolol?', *Medical Hypotheses*. Elsevier Ltd, 74(2), pp. 353–359.
- Mlacker, S., Shah, V. V., Aldahan, A. S., Samarkandy, S., Nouri, K. and Alsaidan, M. (2016) '5-Fluorouracil in the Treatment of Keloids and Hypertrophic Scars: A Comprehensive Review of the Literature', *Dermatology and Therapy*. Springer Healthcare, 6(2), pp. 169–183.
- Modamio, P., Lastra, C. . and Mariño, E. . (2000) 'A comparative in vitro study of percutaneous penetration of β -blockers in human skin', *International Journal of Pharmaceutics*, 194(2), pp. 249–259.
- Monstrey, S. *et al.* (2014) 'Updated Scar Management Practical Guidelines: Non-invasive and invasive measures', *Journal of Plastic, Reconstructive and Aesthetic Surgery*, 67(8), pp. 1017–1025.
- Murphy, A. M., Wong, A. L. and Bezuhly, M. (2015) 'Modulation of angiotensin II signaling in the prevention of fibrosis', *Fibrogenesis and Tissue Repair*. ???, 8(1), pp. 1–7.
- Nanda, S. and Reddy, B. (2004) 'Intralesional 5-fluorouracil as a treatment modality of keloids.', *Dermatol Surg*, 30, pp. 54–7.

- Nast, A. *et al.* (2012) ‘German S2k guidelines for the therapy of pathological scars (hypertrophic scars and keloids)’, 2012(Band 10), pp. 1–14.
- Niazi, F., Hooshyar, sayed hasan, Hedayatyanfard, K. and Doroodgar, F. (2018) ‘Detection of Angiotensin’, *J Clin Aesthet Dermatol*, 11(10), pp. 36–39.
- Noishiki, C., Takagi, G., Kubota, Y. and Ogawa, R. (2017) ‘Endothelial Dysfunction May Promote Keloid Growth’, *accepted article*.
- Ogawa, R. (2010) ‘The most current algorithms for the treatment and prevention of hypertrophic scars and keloids’, *Plastic and Reconstructive Surgery*, 125(2), pp. 557–568.
- Ogawa, R. (2017) ‘Keloid and hypertrophic scars are the result of chronic inflammation in the reticular dermis’, *International Journal of Molecular Sciences*, 18(3).
- Ogawa, R. and Akaishi, S. (2016) ‘Endothelial dysfunction may play a key role in keloid and hypertrophic scar pathogenesis – Keloids and hypertrophic scars may be vascular disorders’, *Medical Hypotheses*. The Authors, 96, pp. 51–60.
- Painter, S. L. and Hildebrand, G. D. (2016) ‘Review of topical beta blockers as treatment for infantile hemangiomas’, *Survey of Ophthalmology*. Elsevier Ltd, 61(1), pp. 51–58.
- Park, H. Y. L., Kim, J. H. and Park, C. K. (2013) ‘VEGF induces TGF- β 1 expression and myofibroblast transformation after glaucoma surgery’, *American Journal of Pathology*, 182(6), pp. 2147–2154.
- Pasquier, E. *et al.* (2011) ‘Propranolol potentiates the anti-angiogenic effects and antitumor efficacy of chemotherapy agents: Implication in breast cancer treatment’, *Oncotarget*, 2(10), pp. 797–809.
- Pasquier, E. *et al.* (2013) ‘B-blockers increase response to chemotherapy via direct antitumour and anti-angiogenic mechanisms in neuroblastoma’, *British Journal of Cancer*. Nature Publishing Group, 108(12), pp. 2485–2494.
- Pozos, E. S. (2014) ‘The effect of skin tension on the formation of keloid scars’.
- Prince, G. T., Cameron, M. C., Fathi, R. and Alkousakis, T. (2018) ‘Pharmacology and therapeutics Topical 5-fluorouracil in dermatologic disease’, pp. 1–6.
- Rei Ogawa (2013) *High Blood Pressure (Hypertension) May Influence the Results*

of Clinical Trials for Scar and Keloid Treatments, Plastic and Reconstructive Surgery.

- Sansan, M. V., Radiono, S., Irawanto, M. E. and Wirohadidjojo, Y. W. (2018) 'Bovine vitreous gel can reavcate replicave senescence of human dermal fibroblast', *Indonesian Journal of Biotechnology*, 23(1), pp. 48–53.
- Sayah, D. N. *et al.* (1999) 'Downregulation of apoptosis-related genes in keloid tissues', *Journal of Surgical Research*, 87(2), pp. 209–216.
- Seeland, U. *et al.* (2009) 'Effects of AT1- and β -adrenergic receptor antagonists on TGF- β 1-induced fibrosis in transgenic mice', *European Journal of Clinical Investigation*, 39(10), pp. 851–859.
- Shirakata, Y. *et al.* (2006) 'A Novel Function of Angiotensin II in Skin Wound Healing', *Journal of Biological Chemistry*, 281(19), pp. 13209–13216.
- Sipkova, Z., Xue, K., Mudhar, H. S., Wagner, B. and Hildebrand, G. D. (2017) 'Early and Late Histological and Ultrastructural Findings in Resected Infantile Capillary Hemangiomas Following Treatment with Topical Beta-Blocker Timolol Maleate 0.5%', *Ocular Oncology and Pathology*, 4(2), pp. 100–106.
- Son, D. and Harijan, A. (2014) 'Overview of surgical scar prevention and management', *Journal of Korean Medical Science*, 29(6), pp. 751–757.
- Stekelenburg, C. and Wal, M. Van Der (2012) 'On the surgical treatment of hypertrophic scars: a comprehensive guideline for the surgical treatment of hypertrophic scars', *European Surgery*, pp. 79–84.
- Stoddart, M. J. (2011) 'Mammalian Cell Viability: Methods and Protocols, Methods in Molecular Biology', *Springer Science+Business Media*, 740, pp. 1–6.
- Takeda, H., Katagata, Y., Hozumi, Y. and Kondo, S. (2004) 'Effects of angiotensin II receptor signaling during skin wound healing', *American Journal of Pathology*, 165(5), pp. 1653–1662.
- Tătaru, C. and Purcărea, V. (2012) 'Antiglaucoma pharmacotherapy', *Journal of medicine and life*, 5(3), pp. 247–251.
- Tullberg-Reinert, H. and Jundt, G. (1999) 'In situ measurement of collagen synthesis by human bone cells with a sirius red-based colorimetric microassay: effects of transforming growth factor beta 2 and ascorbic acid 2-phosphate', *Histochemistry and Cell Biology*, 112(4), pp. 271–276.

- Turner, N. A., Porter, K. E., Smith, W. H. T., White, H. L., Ball, S. G. and Balmforth, A. J. (2003) 'Chronic β 2-adrenergic receptor stimulation increases proliferation of human cardiac fibroblasts via an autocrine mechanism', *Cardiovascular Research*, 57(3), pp. 784–792.
- Um, S. and Henz, B. M. (2004) 'Human skin : source of and target organ for angiotensin II', *Experimental Dermatology*, pp. 148–154.
- Verhaegen, P. D. H. M. *et al.* (2009) 'Differences in collagen architecture between keloid, hypertrophic scar, normotrophic scar, and normal skin: An objective histopathological analysis', *Wound Repair and Regeneration*, 17(5), pp. 649–656.
- Vincent, A. S., Phan, T. T., Mukhopadhyay, A., Lim, H. Y., Halliwell, B. and Wong, K. P. (2008) 'Human skin keloid fibroblasts display bioenergetics of cancer cells', *Journal of Investigative Dermatology*. Elsevier Masson SAS, 128(3), pp. 702–709.
- Weibel, L. *et al.* (2016) 'Topical timolol for infantile hemangiomas: Evidence for efficacy and degree of systemic absorption', *Pediatric Dermatology*, 33(2), pp. 184–190.
- Wendling, J. (2003) '5-Fluorouracil Blocks Transforming Growth Factor- - Induced 2 Type I Collagen Gene (COL1A2) Expression in Human Fibroblasts via c-Jun NH2-Terminal Kinase/Activator Protein-1 Activation', *Molecular Pharmacology*, 64(3), pp. 707–713.
- Widgerow, A. D., Chait, L. A., Stals, R. and Stals, P. J. (2000) 'New innovations in scar management', *Aesthetic Plastic Surgery*, 24(3), pp. 227–234.
- Wirohadidjojo, Y. W., Radiono, S., Budiyo, A. and Soebono, H. (2011) 'Cellular viability, collagen deposition, and transforming growth factor α 1 production among ultraviolet b-irradiated keloid fibroblasts', *Aesthetic Plastic Surgery*, 35(6), pp. 1050–1055.
- Wolfram, D., Tzankov, A., Püzl, P. and Piza-Katzer, H. (2009) 'Hypertrophic scars and keloids - A review of their pathophysiology, risk factors, and therapeutic management', *Dermatologic Surgery*, 35(2), pp. 171–181.
- Wong, T.-S., Li, J. Z.-H., Chen, S., Chan, J. Y.-W. and Gao, W. (2016) 'The Efficacy of Triamcinolone Acetonide in Keloid Treatment: A Systematic Review and Meta-analysis', *Frontiers in Medicine*, 3(December), pp. 1–8.
- Wu, K. Y., Wu, P. C., Lai, Y. H. and Hong, S. J. (2013) 'Novel usage of intraocular pressure-lowering drugs as wound-healing inhibitors after

trabeculectomy with cell culture and animal models', *Kaohsiung Journal of Medical Sciences*. Elsevier Taiwan LLC, 29(7), pp. 353–361.

- Xie, G. *et al.* (2017) 'Hypoxia-induced angiotensin II by the lactate-chymase-dependent mechanism mediates radioresistance of hypoxic tumor cells', *Nature Publishing Group*. Nature Publishing Group, (July 2016), pp. 1–13.
- Xu, J., Yang, E., Yu, N.-Z., Wang, Y.-B. and Long, X. (2017) 'The radiation therapy in keloids treatment: a comprehensive review of pathomechanism, damage mechanisms and cellular response Topic: Keloid and its Treatment Open Access Plastic and Aesthetic Research Radiation therapy in keloids treatment', *Plast Aesthet Res*, 4(4), pp. 116–26.
- Younai, S. (1994) 'modulation of collagen synthesis by transforming growth factor beta in keloid and hypertrophic scar fibroblast', *Annals of Plastic Surgery*, 33(2), pp. 148–154.