

DAFTAR PUSTAKA

- Airheart, JW. (2016). Lived Experiences of Individuals with End-Stage Renal Disease on Peritoneal Dialysis. *Nursing Theses and Capstone Projects*, 253.
- Avrarnovic M, Radovanovic VR., Djordjevic, V., Mitic B, Kostic S. (1997). Fungal Peritonitis in Continuous Ambulatory Peritoneal Dialysis. *The Scientific J Facta Universitatis Medicine and Biology*, 4(1):62-5.
- Barreti P, Moraes TMC, Camargo CH, Caramori JCT, Mondelli AL, dkk. (2012). Peritoneal Dialysis- Related Peritonitis Due to Staphylococcus aureus: A single center Experience over 15 years. *Plos One* 7(5):10.
- Srinivasan B, Matthew HS, Mark SR, Gregory JS, Nirupama R, Lisa MP, Alfred KC. (2003). Impact of Timing of Initiation of Dialysis on Mortality. *J Am Soc Nephrol*: 14(9):2305-12.
- Beth P, George R, Bailie. (2000). Guidelines for Peritoneal Dialysis Adequacy. NKF K/DOQI Guidelines 2000. Available at: <http://www.kidney.org/professionals/kdoq>. Accessed on: 15 th Juli 2007.
- Chern YB, Ho PS, Kuo LC, Chen JB. (2013). Lower Education Level Is a Major Risk Factor for Peritonitis Incidence in Chronic Peritoneal Dialysis Patients: A Retrospective Cohort Study with 12-Year Follow-Up. *Perit Dial Int*, 33(5):552-558.
- Chow KM, Szeto CC., Leung CB., Kwan BCH., Law MC, and Li PKT. (2005). A Risk Analysis of Continuous Ambulatory Peritoneal Dialysis-Related Peritonitis. *PerDial Int*, 2005: 25, (4) 374–379.
- Despina K, Kevan RP, Tony MK, Robert CA, Fiona B. (2007). Risk Factors for Peritoneal Dialysis-Related Peritonitis: Can We Reduce the Incidence and Improve Patient Selection? *Nephrology*: 12(3):239-45.
- Devanarayana NM, Adhikari C, Pannala W, Rajindrajith S. (2011). Prevalence of Functional Gastrointestinal Diseases in A Cohort of Sri Lankan Adolescents: Comparison Between Rome II And Rome III Criteria. *J Trop Ped*;57:34–9.

- DeVore, VS. 2008. Continuouse Ambulatory Peritoneal Dialysis (CAPD) and Its Camplications. <http://www.renal.org/guedelines/module3b.html>. Diperoleh tanggal 6 Juni 2019.
- Duranay M, Kanbay M, Turgut F., Altay M, and Akcay A. (2007). Comparison of incidence of peritonitis between peritoneal dialysis solution types,” *Nephron—Clinical Practice*;106:157–160.
- Ellam T dan Wilkie M. (2015). Chronic Kidney Disease: Peritoneal Dialysis. *Medicine*: 43(8), 484-488.
- Fan X, Huang R, Wang J, Ye H, Guo Q, Yi C, Lin J, dkk. (2014). Risk Factors for the First Episode of Peritonitis in Southern Chinese Continuous Ambulatory Peritoneal Dialysis Patients. *Plos One*;9:9.
- Friedman AN dan Fadem SZ. (2010). Reassessment of Albumin as a Nutritional Marker in Kidney Disease. *JASN* : 21 (2) 223-230.
- Fasianos E, Angelini P, Artesano L, Payota G, Berloco P, dkk. (2008). A Case Report: Recurrent Peritonitis in Peritoneal Dialysis Patient During Antibiotic Therapy: Is Catheter Removal Always Reasonable? *Dialysis and Transplantation*;36:2515-20.
- Guest, Steven. (2013). Hypoalbuminemia in Peritoneal Dialysis Patients. *Adv Perit Dial*; 29:55-60.
- Han SH, Lee SC, Ahn SV, Lee JE, Kim DK, Lee TH, dkk. (2007). Reduced Residual Renal Function Is A Risk Of Peritonitis in Continuous Ambulatory Peritoneal Dialysis Patients. *Nephrol Dial Transplant*;22(9):2653-8.
- Haryanti E, Kandarini Y, Widiana IGR, Sudhana W, Loekman J, Suwitra K. (2010). Kejadian Peritonitis pada Pasien Continuous Ambulatory Peritoneal Dialysis: Identifikasi Mikroorganisme dan Sensitifitas Antibiotik. *J Peny Dal*: 11 (2)
- Huerta RS, Rubio GAF, Flores AG (2010). Severe Hypoalbuminemia: Risk Factor of Peritonitis in Patients In Peritoneal Dialysis. *Med Int Mex* 26 (2).

- Javelosa R, Pena A, Javelosa G, Villanueva N. (1988). A Clinical and Microbiologic Profile of Peritonitis in Chronic Peritoneal Dialysis at UERM Memorial Medical Center: a retrospective study. *Phil J Microbiol Infect Dis*;17(1):34-6.
- Jie D, Yuan C, Suping L, Rong X, Ying X. (2013). Peritoneal Protein Leakage, Systemic Inflammation, and Peritonitis Risk in Patients on Peritoneal Dialysis. *Per Dial Int*;33:273-279.
- Kan GW, Thomas Mark AB, Heath CH. (2005). A 12- Month Review of Peritoneal Dialysis-Related Peritonitis in Western Australia: Is Empiric Vancomycin Still Indicated for Some Patient? *Peritoneal Dialysis International*; 22:465-8.
- KatzIvor J, Sofianou L, Hopley M. (2001). An African Community-Based Chronic Ambulatory Peritoneal Dialysis Programme. *Nephrol Dial Transplant*;16:2395-400.
- Kaysen GA (1998). Biological Basis of Hypoalbuminemia in ESRD. *J Am Soc Nephro*;9(12):2368–2376.
- Keane W.F, Bailie G.R, Boeschoten E, Gokal R. (2000). Adult Peritoneal Dialysis-Related Peritonitis Treatment Recommendations: 2000 update. *Peritoneal Dialysis International*;20: 396-411.
- Keles M, Cetinkaya M, Uyanik A, Acemoglu H, Eroglu H, Uyanik MH. (2010). Peritoneal Dialysis-Related Peritonitis: An Analysis of Risk Factors in Northeast Anatolia. *Turk J Med Sci*; 40 (4): 643-650.
- Kerschbaum J, König P, and Rudnicki M. (2012). Risk Factors Associated with Peritoneal-Dialysis-Related Peritonitis. *Hindawi Publishing Corporation Int J of Nephrology*.
- Kim HJ, Lee J, Park M, Kim Y, Lee H, Kim DK, dkk. (2017). Lower Education Level Is a Risk Factor for Peritonitis and Technique Failure but Not a Risk for Overall Mortality in Peritoneal Dialysis under Comprehensive Training System. *Plos One*: 12(1): e0169063.

- Kofteridis DP, Valachis A, Perakis K, Maraki S, Daphnis E dan Samonis G. (2010). Peritoneal Dialysis-Associated Peritonitis: Clinical Features and Predictors Of Outcome. *Int J of Infectious Diseases* 14 e489–e493.
- Krediet, R.T. (2007). 30 Years of Peritoneal Dialysis Development: The Past and The Future. *Peritoneal Dialysis International*;27:2.
- Lee JS (2012). Albumin for End-Stage Liver Disease. *Journal of the Korean association of internal medicine*;27:1.p.14.
- Lee, Anna. (2011). Constipation in Patients on Peritoneal Dialysis: A literature review. *Renal Society of Australasia Journal*: 7(3)
- Li PKT, Szeto CC, Piraino B, Arteaga J, Fan S, Figueredo AE, dkk. (2016). ISPD Peritonitis Recommendations: 2016 Update on Prevention and Treatment. *Perit Dial Int Sep-Oct*; 36 (5) 481-508
- Lim WH, Boudville N, McDonald SP, Gorham G, Johnson DW, Jose M. (2011). Remote indigenous peritoneal dialysis patients have higher risk of peritonitis, technique failure, all-cause and peritonitis-related mortality. *Nephrol Dial Transplant* 26: 3366–3372.
- Lim WH, Boudville N, McDonald SP. (2005). Higher rate and earlier peritonitis in Aboriginal patients compared to non-Aboriginal patients with end-stage renal failure maintained on peritoneal dialysis in Australia: analysis of ANZDATA. *Nephrology (Carlton)*: 10 (2): 192-7 26: 3366–3372.
- Lobo JV, Villar KR, MP. de Andrade J' unior, Bastos KA. (2010). Predictor Factors of Peritoneal Dialysis-Related Peritonitis. *Jornal Brasileiro de Nefrologia*, vol. 32, no. 2, pp. 156–164.
- Mardiana, Nunuk. (2008). Nutrisi pada Penderita Dialisis, Meeting and Symposium, diunduh dari <http://www.b11nk.wordpress.com>, tanggal 12 Juni 2018
- Maruyama, Yukio dan Yokoyama, Keitaro. (2016). Clinical Efficacy of Combined Therapy with Peritoneal Dialysis and Hemodialysis. *Renal Replacement Therapy, Bio Med Cen.*: 2:11

- Martin LC, Caramori JCT, Fernandes N, Divino-Filho JC, Pecoits-Filho R, Barretti P. (2011). Geographic and Educational Factors and Risk of The First Peritonitis Episode in Brazilian Peritoneal Dialysis Study (BRAZPD) Patients. *Clin J Am Soc Nephrol* 6:1944–1951.
- Mehrotra R, Khawar O, Duong U, Fried L, Norris K, Nissenson A, dkk. (2009). Ownership Patterns Of Dialysis Units and Peritoneal Dialysis in The United States: Utilization and outcomes. *Am J Kidney Dis*;54: 289- 98.
- Mehrotra Rajnish, Duong Uyen, Jiwakanon Sirin, Kovesdy Csaba P, Moran John, Kopple Joel D., Kalantar-Zadeh Kamyar. (2011). Serum Albumin as a Predictor of Mortality in Peritoneal Dialysis: Comparisons With Hemodialysis. *Am J Kidney Dis*; 58: 418–428.
- Mothusi WM, Shepherd K, Jean JN, Ikechukwu OM, Udeme E, Andre PK, Aminu KB, Ikechi GO. (2018). Prevalence of Peritonitis and Mortality Patients Treated with Continuous Ambulatory Peritoneal Dialysis (CAPD) in Africa: a Protocol for a Systematic Review and Meta-Analysis. *BMJ*;8;5.
- Munib, Syeb. Continous Ambulatory Peritoneal Dialysis (CAPD). (2006). Review artikel. *Gomal Jof Medical Sciences*;4(2):82-5.
- Murad MH, Coburn JA, Coto–Yglesias F, Dzyubak S, Hazem A, Lane MA. (2012). Glycemic Control in Non-Critically Ill Hospitalized Patients: A Systematic Review and Meta-Analysis. *J Clin endocrinol metab*; 97:49–58.
- Murray RK, Granner DK, Rodwell VW (2009). Protein Plasma dan Immunoglobulin. Dalam: Buku ajar Biokimia harper. Edisi 27. Jakarta: EGC:608.
- Nagao Y, Sata M. (2010). Serum Albumin and Mortality Risk in a Hyperendemic Area of HCV Infection in Japan. *Virology J*: 7:375.
- National Kidney Foundation. (2002). K/DOQI Clinical Practice Guidelines for Chronic Kidney Disease: Evaluation, Clasification and Stratification. *Am J of Kid Dis*; 39: 1-266.

- Nessim SJ, Bargman JM, Austin PC, Nisenbaum R, Jassal SV. (2009). Predictors of Peritonitis in Patients on Peritoneal Dialysis: Results of a Large, Prospective Canadian Database. *Clin J Am Soc Nephrol*: 4: 1195–1200
- Nessim SJ, Komenda P, Rigatto C, Verrelli M, Sood MM. (2013). Frequency and microbiology of peritonitis and exit-site infection among obese peritoneal dialysis patients. *Perit Dial Int*; 33:167.
- Oygar DD, Yalin AS, Altiparmak MR., Ataman R, and Serdengeçti K. (2011). Obligatory Referral Among Other Factors Associated with Peritonitis in Peritoneal Dialysis Patients. *Nefrologia*: 31:4; 435–440.
- Ozturk S, Soyuk O, Karakaya D, Yazici H, Caliskan YK, Yildiz A, Bozfakioglu S. (2009). Is Decline in Serum Albumin an Ominous Sign for Subsequent Peritonitis in Peritoneal Dialysis Patients?. *Advances in Peritoneal Dialysis, Conference on Peritoneal Dialysis*: 25;172–177.
- Pollock CA, Ibels LS, Caterson RJ. (1989). Continuous Ambulatory Peritoneal Dialysis: Eight Years of Experience at a Single Center. *Medicine (Baltimore)*;68(5):293-308.
- Pessegueiro P, Amoedo M, Barros S, Aniceto J, Pires C. (2006). Aseptic Peritonitis in a Peritoneal Dialysis Patient. *Nefrologia*;26(4):493-6.
- Peter BS, Robert RW, Alan S. (2019) Hypoalbuminemia: Pathogenesis and Clinical Significance. *J of Par and Ent Nut*; 43: 181–193
- Piraino B, Bailie G.R, Bernardini J, Boeschoten E, Gupta A, Holmes C, dkk. (2005). Peritoneal Dialysis-Related Infections Recommendations: 2005 update. *Peritoneal Dialysis International*;25:107-31.
- Piraino B, Bernardini J, Centa PK, Johnston JR, Sorkin MI. (1991). The Effect of Body Weight on CAPD Related Infections and Catheter Loss. *Perit Dial Int* ;11(1):64-8.
- Prasad N, Sinha A, Gupta A, Sharma RK, Bhadauria D, Chandra A, Prasad KN, Kaul A. (2014). Effect of Body Mass Index on Outcomes of Peritoneal Dialysis Patients in India. *Perit Dial Int*. Jun;34(4):399-408.
- Rachnindar, D. 2013. Hypoalbuminemia. Available from: <http://id.scribd.com/doc/124643683/Hypoalbuminemia>

- Rodríguez-Carmona A, Pérez-Fontán M, López-Muñiz A, Ferreiro-Hermida T, García-Falcón T. (2014). Correlation Between Glycemic Control and the Incidence of Peritoneal and Catheter Tunnel and Exit-Site Infections in Diabetic Patients Undergoing Peritoneal Dialysis. *Perit Dial Int. Sep-Oct*; 34(6): 618–626.
- Sacher, Ronald A. (2004). *Tinjauan Klinik Hasil Pemeriksaan Laboratorium*. Jakarta:EGC.
- Saklayen MG. (1990). Incidence, Pathogens, Diagnosis, and Management. *Med Clin North Am*, Jul;74(4):997-1010.
- Salzer WL. (2018). Peritoneal Dialysis Related Peritonitis: Challenges and solutions. *Int J Nephrol Renovasc Dis*;11:173-186.
- Sanjeev G, Derek S, Judith A, Donna S, Elizabeth H, Williamson B. (2001). Children with Hypoalbuminemia on Continuous Peritoneal Dialysis are at Risk for Technique Failure. *Kidney International*; 59:2361–2367.
- Santoianni JE, Predari SC, Veron D, Zucchini A, De Paulis AN. (2008). A 15 Year-Review of Peritoneal Dialysis-Related Peritonitis: Microbiological Trends and Patterns of Infection in a Teaching Hospital in Argentine. *Revista Argentina de Microbiologia*;40:17-23.
- Sibuea, Herdin; Panggabean, Marulan; Gultom. (2005). *Ilmu Penyakit Dalam*, Jakarta:PT Rineka Cipta.
- Sirivongs D, Pongskul C, Keobounma T, Chunlertrith D, Sritaso K, and Johns J. (2006). Risk factors of first peritonitis episode in Thai CAPD patients. *J of the Med Assoc Thai*;89;2;138–145.
- Spencer RC. (1988). Infections in Continuous Ambulatory Peritoneal Dialysis. *J Med Microbiol*;27: 1-9.
- Suwitra K. (2014). *Penyakit Ginjal Kronik: Buku Ajar Ilmu Penyakit Dalam Edisi ke VI*. Jakarta: Interna Publishing;. p. 2159.
- Than NO, Tricia LR, Allan JC. (2005). A Comparison of Peritonitis Rates from The United States Renal Data System Database: CAPD Versus Continuous Cycling Peritoneal Dialysis Patients. *Am J Kidney Dis*: 45(2):372-80.

- Tian Y, Xie X, Xiang S, Yang X, Lin J, Zang X, Shou Z dan Chen J. (2017). Risk Factors and Outcomes of Early-Onset Peritonitis In Chinese Peritoneal Dialysis Patients. *Kidney Blood Press Res*;42:1266-1276.
- Tian Y, Xie X, Xiang S, Yang X, Zhang X, Shou Z dan Chen J. (2016). Risk Factors and Outcomes of High Peritonitis Rate in Continuous Ambulatory Peritoneal Dialysis Patients. *Medicine*: 95:49.
- Wang Q, Bernardini J, Piraino B, Fried L. (2003). Albumin at The Start of Peritoneal Dialysis Predicts The Development of Peritonitis. *Am J of Kidney Disease*. March 2003; 41 (3): 664-669
- Widiana, I Gede. *Distribusi Geografis Penyakit Ginjal Kronik Di Bali: Komparasi Formula Cockcroft-Gault Dan Formula*. Artik Asli Bag/SMF Ilmu Penyakit Dalam FK Unud / RSUP Sanglah, Denpasar. 2007;8(September 2007):2-8.
- Woo PCY, Wong SSY, Lau SKP, Yuen K. (2004). Continous Ambulatory Peritoneal Dialysis-Related Peritonitis Associated with Lancefield Group G Betahemolytic Streptococcus:Report of Two Cases Requiring Tenckhoff Catheter Removal. *J of Clinical Microbiology*;42(9):4399-402.
- Yuwono, Agus (2000). *Kualitas Hidup Menurut Spitzer pada Penderita Gagal Ginjal Terminal yang Menjalani Hemodialisis di RSUD. Kariadi Semarang*, diunduh dari <http://www.eprint.undip.ac.id>, tanggal 28 Juni 2018.