

DAFTAR PUSTAKA

- Albertsmeier, M. *et al.* (2012) 'Evaluation of the safety and efficacy of MonoMax® suture material for abdominal wall closure after primary midline laparotomy - A controlled prospective multicentre trial: ISSAAC [NCT005725079]', *Langenbeck's Archives of Surgery*, 397(3), pp. 363–371. doi: 10.1007/s00423-011-0884-6.
- Anderson, J.M. (2000) The cellular cascades of wound healing. In J. E. Davies (Ed.), *Bone Engineering*. Toronto: em squared inc., p:81–93.
- Beck, Rosenberg, *et al* (2003) Transforming Growth Factor- β Mediates Intestinal Healing and susceptibility to Injury in Vitro and in Vivo Through Epithelial Cells. *The American Journal of Pathology* ; 162: 597–608.
- Bisono, Puspongoro, A.P. (1997) Luka, syok, bencana. Dalam Buku ajar bedah. Jakarta: EGC, p:73-75.
- Bosanquet, D. C. *et al.* (2015) 'Systematic review and meta-regression of factors affecting midline Incisional hernia rates: Analysis of 14 618 Patients', *PLoS ONE*, 10(9), pp. 1–18. doi: 10.1371/journal.pone.0138745.
- Boyle, M. (2006) Wound healing in Midwifery. United Kingdom. Radcliffe. Publishing. Page :14.
- Ceydeli, A., Rucinski, J. and Wise, L. (2007) 'Finding the best abdominal closure - An evidence-based overview of the literature', *Recurrent Hernia: Prevention and Treatment*, pp. 117–122. doi: 10.1007/978-3-540-68988-1_14.
- Chalya, P. L. *et al.* (2015) 'Abdominal fascia closure following elective midline laparotomy: A surgical experience at a tertiary care hospital in Tanzania', *BMC Research Notes*. BioMed Central, 8(1), pp. 1–9. doi: 10.1186/s13104-015-1243-4.
- Dahlan, S. (2011) '*Statistik Untuk Kedokteran dan Kesehatan*', Edisi 5, Jakarta: Salemba Medika.
- Dart, A. J. and Dart, C. M. (2011) 'Suture Material: Conventional and Stimuli Responsive', *Comprehensive Biomaterials*, pp. 573–587. doi: 10.1016/B978-0-08-055294-1.00245-2.
- David, L. Dunn (2004) "*The Wound Closure Manual*". Ethicon, inc a Johnson & Johnson company.
- Deerenberg, E. B. *et al.* (2015) 'Small bites versus large bites for closure of abdominal midline incisions (STITCH): A double-blind, multicentre, randomised controlled trial', *The Lancet*. Elsevier Ltd, 386(10000), pp. 1254–1260. doi: 10.1016/S0140-6736(15)60459-7.
- Djaya, W., Rudiman, R., & Lukman, K. (2012) Efek Oksigen Konsentrat Tinggi Paskaoperasi Laparatomi terhadap Tingkat Infeksi Luka Operasi. *MKBB*,

- 40(3).Dubay, D. A. and Franz, M. G. (2003) 'Acute wound healing: The biology of acute wound failure', *Surgical Clinics of North America*, pp. 463–481. doi: 10.1016/S0039-6109(02)00196-2.
- Dubay, D. A. *et al.* (2004) 'Fascial fibroblast kinetic activity is increased during abdominal wall repair compared to dermal fibroblasts', *Wound Repair and Regeneration*, 12(5), pp. 539–545. doi: 10.1111/j.1067-1927.2004.012506.
- Ellis, (2009) *Ephraim Mc Dowell and the first successful elective laparotomy*. Available at : <https://ncbi.nlm.nih.gov/m/pubmed19229153/>.
- Enoch, S., & Leaper, D.J. (2005) Basic Science of wound healing. *Surgery journal*, 26(2), 31–37. DOI : 10.1016/j.mpsur.2007.11.005.
- Faiz, O. and Moffat, D. (2002) *Anatomy at a Glance [e-Book]*. doi: 10.5005/jp/books/10050.
- Fernandez, L. G. (2017) *Abdominal Closure Technique: Closure of Abdomen, Postoperative Care, Complications, Medscape*. Available at: <https://emedicine.medscape.com/article/1961789-technique>.
- Fink, C. *et al.* (2014) 'Incisional hernia rate 3 years after midline laparotomy', *British Journal of Surgery*, 101(2), pp. 51–54. doi: 10.1002/bjs.9364.
- Fortelny, R. H. *et al.* (2015) 'Effect of suture technique on the occurrence of incisional hernia after elective midline abdominal wall closure: Study protocol for a randomized controlled trial', *Trials*, 16(1), pp. 1–8. doi: 10.1186/s13063-015-0572-x.
- Franz, M. G. (2009) 'The Biology of Hernia Formation-2009', 88(1), pp. 1–16. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276402/>.
- Franz, M. G. *et al.* (2001) 'Fasial incisions heal faster than skin: A new model of abdominal wall repair', *Surgery*, 129(2), pp. 203–208. doi: 10.1067/msy.2001.110220.
- Gurtner, G.C. (2007) Wound healing, normal and abnormal. In: Thorne CH, Beasley, R.W., Aston, S.J., Bartlett, S.P., Gurtner, G.C., Spear, S.L. (Eds). *Grabb and Smith's plastic surgery*. 6th ed. Philadelphia: Lippincott Williams and Wilkins; p:15-22.
- Hammamy E, B-Lynch C (2005) *A worldwide review of the uses of the uterinecompression jahitan techniques as alternative to hysterectomy in the management of severe post-partum haemorrhage. J Obstet Gynaecol ;25:143–9.*
- Heinrich, P. C. *et al.* (2003) 'Principles of interleukin (IL)-6-type cytokine signalling and its regulation', *Biochemical Journal*, 374(1), pp. 1–20. doi: 10.1042/bj20030407.
- Hitesh, K., Pratik, V., Nilesh, P., & Jovin, M. (2015) Factors affecting post-operative laparotomy wound complications. *International archives of Integrated Medicine*, 2(1), 71–74.
- Hodgson, N. C. F., Malthaner, R. A. and Østbye, T. (2000) 'The search for an ideal method of abdominal fasial closure: A meta- analysis', *Annals of Surgery*, 231(3), pp. 436–442. doi: 10.1097/00000658-200003000-00018.
- Höer, J. *et al.* (2002) 'Factors influencing the development of incisional hernia. A retrospective study of 2,983 laparotomy patients over a period of 10

- years', *Der Chirurg*, 73(5), pp. 474–480. doi: 10.1007/s00104-002-0425-5.
- Indrawan, I. and Dachlan, I. (2016) 'Perbandingan pengaruh aplikasi aloe vera, madu, saliva dan putih telur terhadap ekspresi interleukin-6 pada proses penyembuhan luka insisi kulit tikus', Universitas Gadjah Mada, pp. 1-14.
- Israelsson, L. A. and Millbourn, D. (2013) 'Prevention of incisional hernias. How to close a midline incision.', *Surgical Clinics of North America*, pp. 1027–1040. doi: 10.1016/j.suc.2013.06.009.
- Janz, A. B. (2013) *Regions and Planes of the Abdomen: Overview, Abdominal Skin, Superficial Fasia*. Available at: <http://emedicine.medscape.com/article/1923166-overview#a1>.
- Johnson, C.M. (2009) Development of abdominal wound dehiscence after colectomy : a nursing challenge. *MedSurg Nursing*, 1–7.
- Keith, L. and Arthur, F. (2012) Ovid: Clinically Oriented Anatomy. Available at: <http://ueu.co/zh/ovid-clinically-oriented-anatomy-2/> (Accessed: 19 January 2018).
- Kenig, J., Richter, P., Lasek, A., Zurawska, S. & Zbierska, K., (2013). Risk factors for wound dehiscence after laparotomy-Clinical control trial. [Pol Przegl Chir.](#);84(11):565-73. doi: 10.2478/v10035-012-0094-0.
- Khorgami, Z., Shoar, B., Laghaie, B., Aminian, A., Araghi, N.H., & Soroush, A. (2012) Prophylactic retention suture in midline laparotomy in high risk for wound dehiscence : a randomized controlled trial. *Journal of Surgical Research*, xxx, E1-E6.
- Klinge, U. *et al.* (2001) 'Collagen I/III and matrix metalloproteinases (MMP) 1 and 13 in the fascia of patients with incisional hernias', *Journal of Investigative Surgery*, pp. 47–54. doi: 10.1080/089419301750072202.
- Kopf, M. *et al.* (1994) 'Impaired immune and acute-phase responses in interleukin-6-deficient mice', *Nature*, 368(6469), pp. 339–342. doi: 10.1038/368339a0.
- Kumar, R. and Hastir, A. (2017) 'Prospective Clinical Study: Mass Closure Versus Layer Closure of Abdominal Wall', *International Journal of Surgery and Medicine*, 3, p. 1. doi: 10.5455/ijism.mass-closure-versus-layer-closure-abdominal-wall.
- Lambertz, A. *et al.* (2015) 'Polyvinylidene fluoride as a suture material: Evaluation of comet tail-like infiltrate and foreign body granuloma', *European Surgical Research*, 55(1–2), pp. 1–11. doi: 10.1159/000371797.
- Lau, F. H. and Pomahac, B. (2014) 'Wound healing in acutely injured fascia', *Wound Rep Reg*, 22, pp. 14–17. doi: 10.1111/wrr.12165.
- Lin, Z.-Q. (2003) 'Essential involvement of IL-6 in the skin wound-healing process as evidenced by delayed wound healing in IL-6-deficient mice', *Journal of Leukocyte Biology*, 73(6), pp. 713–721. doi: 10.1189/jlb.0802397.

- Liptan, G. L. (2010) 'Fasia: A missing link in our understanding of the pathology of fibromyalgia', *Journal of Bodywork and Movement Therapies*. Elsevier Ltd, 14(1), pp. 3–12. doi: 10.1016/j.jbmt.2009.08.003.
- Lock, A. M., Gao, R., Naot, D., Coleman, B., Cornish, J., & Musson, D. S. (2017) Induction of immune gene expression and inflammatory mediator release by commonly used surgical suture materials: an experimental in vitro study. *Patient Safety in Surgery*, 11, 16. <http://doi.org/10.1186/s13037-017-0132-2>.
- Lorentz, H. P., Longaker, M. T (2006) Wound Healing: Repair Biology and Wound and Scar Treatment. In: Mathes, S. J. and Hentz, V. R., (Eds). *Plastic surgery*. 2nd ed. Philadelphia: Saunders Elsevier; 209-234.
- Perdanakusuma, D.S. (2006) Penanganan luka pada luka bakar. In Noer, M.S. (eds) Penanganan luka bakar. Airlangga University Press. Surabaya. p: 83, 89.
- Potter, P.A. and Perry, A.G. (2005) Buku Ajar Fundamental Keperawatan: Konsep, Proses, dan Praktik. Edisi 4, Volume 1, Alih Bahasa, Asih, Y., dkk. EGC, Jakarta.
- Poulose BK, Shelton J, Phillips S, et al. (2012) Epidemiology and cost of ventral hernia repair: making the case for hernia research. *Hernia* ; **16**: 179–83.
- Ramshorst Gabrielle H van, Eker Hasan H, Hop Wim C.J., Jeekel Johannes, L. J. F. (2012) 'Impact of incisional hernia on health-related quality of life and body image : a prospective cohort study', *AJS*. Elsevier Inc., 204(2), pp. 144–150. doi: 10.1016/j.amjsurg.2012.01.012.
- Roberts AB, Sporn MB, Assoian RK, Smith JM, Roche NS, Wakefield LM, Heine UI, Liotta LA, Falanga V, Kehrl JH, Fauci AS (1986) Transforming growth factor- β : rapid induction of fibrosis and angiogenesis in vivo and stimulation of collagen formation in vitro. *Proc Natl Acad Sci USA*. 1986;83:4167–4171. [[PMC free article](#)] [[PubMed](#)].
- Marston WA (2006).: Risk factors associated with healing. *Ostomy Wound Manage* ; 53:26-8, 30, 32.
- Mary, C., Marois, Y. and King, M. W. (1998) 'Comparison of the in vivo behavior of polyvinylidene flouride and polypropylen sutures used in vascular surgery', *ASAIO Journal*, pp. 199–206. Available at: <http://europepmc.org/abstract/med/9617952>.
- Meijer, E. J. *et al.* (2013) 'The principles of abdominal wound closure', *Acta Chirurgica Belgica*, 113(4), pp. 239–244. doi: [http://dx.doi.org/10.1016/0890-4332\(93\)90047-Y](http://dx.doi.org/10.1016/0890-4332(93)90047-Y).
- Mercandetti, A. M., Editor, C. and Molnar, J. A. (2015) 'Wound Healing and Repair: Overview, Types of Wound Healing, Categories of Wound Healing', pp. 1–7. Available at: <http://emedicine.medscape.com/article/1298129-overview>.

- Millbourn, D. (2009) Effect of Stitch Length on Wound Complications After Closure of Midline Incisions A Randomized Controlled Trial. *Archives of Surgery*, 144 (11), 1056–1059. <https://doi.org/10.1001/archsurg.2009.189>.
- Miyzono K, Heldin CH.(1992 Structure, function and possible clinical application of TGF-beta1. *Dermatology*.1992;19:644-647 8.
- Mizell, J. S. (2015) ‘Complications of abdominal surgical incisions’, *UpToDate*, pp. 1–27.
- Muysoms, F. E. *et al.* (2015) ‘European Hernia Society guidelines on the closure of abdominal wall incisions’, *Hernia: the journal of hernias and abdominal wall surgery*, 19(1), pp. 1–24. doi: 10.1007/s10029-014-1342-
- Molloy, Li, Y., R. Moysey, Molloy, A. L. Vuidepot, T. Mahon, E. Baston, S. Dunn, N. Liddy, J. Jacob, B. K. Jakobsen, and J. M. Boulter. 2005. Directed evolution of human T-cell receptors with picomolar affinities by phage display. *Nat. Biotechnol.* 23: 349–354.
- Moore KL, Dalley AF, eds (1999) *Clinically Oriented Anatomy*. 4th ed. Philadelphia: Lippincott Williams & Wilkins :185.
- Naka, T., Nishimoto, N. and Kishimoto, T. (2002) ‘The paradigm of IL-6: from basic science to medicine.’, *Arthritis research*, 4 Suppl 3, pp. S233–S242. doi: 10.1186/ar565.
- Nall AV, Brownlee RE, Colvin CP, Schultz G, Fein D, Cassisi NJ, Nguyen T, Kalra A (1996) Transforming growth factor beta 1 improves wound healing and random flap survival in normal and irradiated rats. *Arch Otolaryngol.*; 122:171–177. [[PubMed](#)].
- Nout, E. *et al.* (2007) ‘Creep Behavior of Commonly Used Suture Materials in Abdominal Wall Surgery’, *Journal of Surgical Research*, 138(1), pp. 51–55. doi: 10.1016/j.jss.2006.06.001.
- Osther, P. J. *et al.* (1995) ‘Randomized comparison of polyglycolic acid and polyglyconate sutures for abdominal fasial closure after laparotomy in patients with suspected impaired wound healing’, *British Journal of Surgery*, 82(8), pp. 1080–1082. doi: 10.1002/bjs.1800820824.
- Popa, F. and Georgescu, A. V (2017) ‘Abdominal Wall Reconstruction after Flap Surgery and the Effect on the Immune System’, *Hindawi*, 2017, pp. 1–10.
- Radu, P. *et al.* (2013) ‘Molecular factors of failure in incisional hernia surgery.’,
- Rahbari, N. N. *et al.* (2009) ‘Current practice of abdominal wall closure in elective surgery? Is there any consensus?’, *BMC Surgery*, 9(1), pp. 1–8. doi: 10.1186/1471-2482-9-8.
- Ramshorst Gabrielle H van, Eker Hasan H, Hop Wim C.J., Jeekel Johannes, L. J. F. (2012) ‘Impact of incisional hernia on health-related quality of life and body image : a prospective cohort study’, *AJS*. Elsevier Inc., 204(2), pp. 144–150. doi: 10.1016/j.amjsurg.2012.01.012.

- Regan, M.C., Barbul, A. (1994) The cellular biology of wound healing. In (eds) Redl, H., Schlag, G. Wound healing. Berlin: Springer-Verlag, 1: 2-13.
- Robson, M. C. *et al.* (2000) 'Wound healing trajectories as predictors of effectiveness of therapeutic agents', *Archives of surgery*. doi: 10.1001/archsurg.135.7.773.
- Rohrich, R.J., Robinson, J.B. (1999) Wound healing. Selected readings in plastic surgery 9 (3): 1-17
- Roses, R. E. and Morris, J. B. (2013) 'Incisions, Closures, And Management of The Abdominal Wound', in Zinner, M. J. and Ashley, S. W. (eds) *Maingot's Abdominal Operations*. 12th editi. New York: Mc Graw Hill Companies, pp. 99–122.
- Satteson, E. S. (2017) *Materials for Wound Closure: Wound Healing and Closure, Suture Characteristics, Suture Materials*, Medscape. Available at: <https://emedicine.medscape.com/article/1127693overview%0Ahttps://emedicine.medscape.com/article/1127693-overview#a3>.
- Stewart, R. J. *et al.* (1981) 'the Wound Fibroblast and Macrophage .1. Wound Cell-Population Changes Observed in Tissue-Culture', *British Journal of Surgery*, 68(2), pp. 125–128.
- Shah, M., Foreman, D.M., Ferguson, M.W., (1995) Neutralisation of TGF-beta 1 and TGF beta 2 or exogenous addition of TGF-beta 3 to cutaneous rat wounds reduces.
- Simon S.R (1994) Anatomy Biology and biomechanics of Tendon, Ligament and Meniscus, *Orthopaedic Basic Science, American Academy of Orthopaedic Surgeons*, p 47-58.
- Sajid, M *et al.*, (2014) Systematic review of absorbable vs non-absorbable sutures used for the closure of surgical incisions. *World J Gastrointest Surg.*; 6(12): 241–247. doi: [10.4240/wjgs.v6.i12.241](https://doi.org/10.4240/wjgs.v6.i12.241)
- Schultz, G.S. (2007) The physiology of wound bed preparation. In (eds) Granick, M.S., Ganelli, R.L. *Surgical wound healing and management*. New York: Informa Healthcare USA Inc., p:1-5.
- Simon Gurzu, Ioan Jung (2017), *Textbook of Iatrogenic Pathology*. Betham Science Publishers, page 281.
- Sivender, A., Ilaiah, M., &Reddy, S. (2015) A Clinical Study on risk factors causing abdominal wound dehiscence and management. *IOSR Journal of Dental and Medical Sciences*, 14(10), 18–23.
- Skandalakis (2014) *Surgical Anatomy and Technique A Pocket Manual Fourth Edition*, Springer New York Heidelberg Dordrecht London.
- Slater, N.J., Bleichrodt R.P., and Goor H.V (2012) Wound dehiscence and incisional hernia. *Surgery (Oxford)* 2012; 30:282–289.

- Smeltzer, S.C., & Bare, B.G. (2010). Post operative nursing wound management. *In Brunner & Suddart's Textbook of medicalsurgical nursing* (12th ed.). Philadelphia: J.B. Lippincott Company.
- Spiliotis, J., T siveriotis, K., Datsis, A.D., Vaxevanidou, A., Zacharis, G., Konstantinos, G. Rogdakis, A. (2009) Wound Dehiscence : is still a problem in the 21th century : a restropective study. *World Journal of Emergency Surgery*, 4(12), 1–5. doi:10.1186/1749-7922-4-12.
- Sporn MB, Roberts AB, Wakefield NM, *et al.* (1987): Some recent advances in the chemistry and biology of TGF beta. *J. Cell. Biol* ; 105:1039-1045.
- Thiruvoth, F. *et al.* (2015) ‘Current concepts in the physiology of adult wound healing’, *Plastic and Aesthetic Research*, 2(5), p. 250. doi: 10.4103/2347-9264.158851.
- Urban, E. *et al.* (1994) ‘Why make monofilament sutures out of polyvinylidene fluoride?’, *ASAIO journal (American Society for Artificial Internal Organs : 1992)*, pp. 145–156. Available at:
http://journals.lww.com/asaiojournal/Abstract/1994/04000/Why_Make_Monofilament_Sutures_Out_of.6.aspx%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/8003751.
- Urvashi Vats, [Pandit Suchitra N](#) . Comparison of Efficacy of Three Suture Materials, i.e., Poliglecaprone 25, Polyglactin 910, Polyamide, as Subcuticular Skin Stitches in Post-Cesarean Women: A Randomized Clinical Trial . [J Obstet Gynaecol India](#). (2013) ;64(1):14-8. doi: 10.1007/s13224-013-0448-5.
- Van Ramshorst, G. H. *et al.* (2012) ‘Impact of incisional hernia on health-related quality of life and body image: A prospective cohort study’, *American Journal of Surgery*. doi: 10.1016/j.amjsurg.2012.01.012.
- Veljkovic, R. *et al.* (2010) ‘Prospective Clinical Trial of Factors Predicting the Early Development of Incisional Hernia after Midline Laparotomy’, *Journal of the American College of Surgeons*. Elsevier Inc., 210(2), pp. 210–219. doi: 10.1016/j.jamcollsurg.2009.10.013.
- Vishy M (2006) *Abdominal Surgery : Anatomy of the anterior abdominal wall and groin*. Royal College of Surgeons of England, London, . United Kingdom. Elsevier Ltd.
- Waqar, S., Malik, Z., Razzaq, A., Abdullah, M., Shaima, A., & Zahid, M. (2005) Frequency and risk factors for wound dehiscence/burst abdomen in midline laparotomies. *Journal Ayub Medical College Abottabad*, 17(4), 70–73.
- Xing, L. *et al.* (2013) ‘Early laparotomy wound failure as the mechanism for incisional hernia formation’, *Journal of Surgical Research*. Elsevier Ltd, 182(1), pp. 1–8. doi: 10.1016/j.jss.2012.09.009.



UNIVERSITAS
GADJAH MADA

Perbandingan Pengaruh Jahitan Antara Benang Polyglecaprone 25 Dengan Teknik Kontinyu Large Stitch Dan Kontinyu Small Stitch Terhadap Transforming Growth Factor Beta Pada Garis Insisi Kulit Abdomen Tikus Albino Galur Wistar (*Rattus norvegicus*)
DHYAS MUNANDAR AS, dr. Imam Sofii Sp.B-KBD , DR. dr. Akhmad Mahmudi Sp.B., Sp.BA (K)
Universitas Gadjah Mada, 2019 | Diunduh dari <http://etd.repository.ugm.ac.id/>