

DAFTAR PUSTAKA

1. Miller M. The Kremlin papers. *Technology Wound Journal*. 2008;1:22–24.
2. Fleischmann, W., Becker, U., Bischoff, M., Hoekstra, H. Vacuum sealing: indications, technique and results. *Eur J Orthop Surg Traumatol*. 1995; 5:37–40.
3. Morykwas, M.J., Argenta, L.C., Shelton-Brown, E.I., McGuirt, W. Vacuum assisted closure: a new method for wound control and treatment: animal studies and basic foundation. *Ann Plast Surg*. 1997; 38 (6): 553–562.
4. Argenta, L.C., Morykwas, M.J. Vacuum-assisted closure: a new method for wound control and treatment: clinical experience. *Ann Plast Surg*. 1997; 38 (6): 563–577.
5. Joseph, E., Hamori, C.A., Berman, S. et al. A prospective randomized trial of vacuum-assisted closure versus standard therapy of chronic nonhealing wounds. *Wounds*. 2000; 12(3). 60–67.
6. McCallon, S.K., Knight, C.A., Valiulus, J.P. et al. Vacuum-assisted closure versus saline-moistened gauze in the healing of postoperative diabetic foot wounds. *Ostomy Wound Manage*. 2000; 46: 8, 28–34.
7. Chariker, M.E., Jeter, K.F., Tintle, T.E., Bottsford, J.E. Effective management of incisional and cutaneous fistulae with closed suction wound drainage. *Contemp Surg*. 1989; 34: 59–63.
8. Bobkiewicz, A., Banasiewicz, T., Ledwosinski, W. Drews, M. Medical terminology associated with Negative Pressure Wound Therapy (NPWT). Understanding and misunderstanding in the field of NPWT. 2014; 1(2): 69–73.
9. Hampton, J. Providing cost-effective treatment of hard-to-heal wounds in the community through use of NPWT. *Br J Community Nurs*. 2015; 20: Sup6, S14–S20.
10. Apelqvist, J., Armstrong, D.G., Lavery, L.A., Boulton, A.J. Resource utilization and economic costs of care based on a randomized trial of vacuum-assisted closure therapy in the treatment of diabetic foot wounds. *Am J Surg*. 2008; 195 (6): 782–788.

11. Enquist I, Adamson R: Collagen syntheses and lysis in healing wounds. *Minnesota Medicine*. 48; 1965. 1695–1698,
12. Gruber RP, Vistnes L, Pardoe R: The effect of commonly used antiseptics on wound healing. *Plastic and Reconstructive Surgery*. 55.4; 1975. 472-476.
13. Peacock E, Madden J. Some studies on the effects of B-aminopropionitrile on collagen in healing wounds. *Surgery* 60; 1969. 7–12.
14. Dingman R. Factors of clinical significance affecting wound healing. *The Laryngoscope* 83.9; 1973. 1540–1544.
15. Moss G, Bierenbaum A, Bova F: Postoperative metabolic patterns following immediate total nutritional support: Hormone levels, DNA syntheses, nitrogen balance, and accelerated wound healing. *J Surg Res*. 1976; 21(6): 383-393.
16. Hunt T, Van Winkle W. *Wound healing: Normal Repair Fundamentals of Wound Management in Surgery*. South Plainfield, NJ, Chirurgecom Inc; 1976.
17. Montadon D, et al. The mechanism of wound contraction and epithelialization. *Clin Plast Surg*. 1977; 4: 325-345.
18. Niinikoshi J. Oxygen and wound healing. *N Am Clin Plast Surg*. 1977; 4: 361.
19. Goss RJ. *The Physiology of Growth*. New York, Academic Press; 1978.
20. Colin JF, et al. The effect of uraemia upon wound healing: An experimental study. *Br J Surg*. 1979; 66(11): 793-797.
21. Pollack, Sheldon V. Wound healing: A review. III. Nutritional factors affecting wound healing. *J Dermatol Surg Oncol*. 1979; 5(8): 615–619.
22. Thakral KK, Goodson WH III, Hunt TK. Stimulation of wound blood vessel growth by wound macrophages. *J Surg Res*. 1979; 26(4): 430–436.
23. Kunt T. Disorders of wound healing. *World J Surg*. 1980; 4(3): 271–277.
24. Linsky CB, Rovee DT, Dow T. Effects of dressings on wound inflammation and scar tissue, in Dineen P, Hildick-Smith G, (eds), *The Surgical Wound*. Philadelphia, Lea and Febiger; 1981, pp. 876–891.

25. Acosta, S., Bjarnason, T., Petersson, U. et al. Multicentre prospective study of fascial closure rate after open abdomen with vacuum and mesh-mediated fascial traction. *Br J Surg.* 2011; 98(5): 735–743.
26. Swan, M., Banwell, P. Topical negative pressure. Advanced management of the open abdomen. Oxford Wound Healing Society. 2003.
27. Fong, K.D., Hu, D., Eichstadt, S. et al. The SNaP system: biomechanical and animal model testing of a novel ultraportable negative-pressure wound therapy system. *Plast Reconstr Surg.* 2010; 125(5): 1362–1371.
28. Marston, W.A., Armstrong, D.G., Reyzelman, A.M., Kirsner, R.S. A multicenter randomized controlled trial comparing treatment of venous leg ulcers using mechanically versus electrically powered negative pressure wound therapy. *Adv Wound Care.* 2015; 4(2): 75–82.
29. Birke-Sorensen, H., Malmsjo, M., Rome, P. et al. Evidence-based recommendations for negative pressure wound therapy: treatment variables (pressure levels, wound filler and contact layer) – Steps towards an international consensus. *J Plast Reconstr Aesthet Surg.* 2011; 64: S1–S16.
30. Torbrand, C., Ingemansson, R., Gustafsson, L. et al. Pressure transduction to the thoracic cavity during topical negative pressure therapy of a sternotomy wound. *Int Wound J.* 2008; 5(4): 579–584.
31. Petzina, R., Ugander, M., Gustafsson, L. et al. Hemodynamic effects of vacuum-assisted closure therapy in cardiac surgery: Assessment using magnetic resonance imaging. *J Thorac Cardiovasc Surg.* 2007; 133(5): 1154–1162.
32. Cheng, B., Fu, X.B., Gu, X.M. et al. The regulation mechanisms of MMP-1,2 and TIMP-1,2 on wound healing after partial thickness scald. *Zhonghua Wai Ke Za Zhi [Chinese journal of surgery].* 2003; 41(10): 766–769.
33. Cook, H., Stephens, P., Davies, K.J. et al. Defective extracellular matrix reorganization by chronic wound fibroblasts is associated with alterations in TIMP-1, TIMP-2, and MMP-2 activity. *J Invest Dermatol* 2000; 115(2): 225–233.
34. Maier, D., Beck, A., Kinzl, L., Bischoff, M. The physics of vacuum therapy. *Zentralbl Chir.* 2005; 130(5): 463–468.
35. Willy, C., von Thun-Hohenstein, H., von Lubken, F. et al. Experimental principles of the V.A.C.-therapy pressure values in superficial soft tissue and the applied foam. *Zentralbl Chir.* 2006; 131: S50–S61.

36. Eisenhardt, S.U., Schmidt, Y., Thiele, J.R. et al. Negative pressure wound therapy reduces the ischaemia/reperfusion-associated inflammatory response in free muscle flaps. *J Plast Reconstr Aesthet Surg.* 2012; 65(5): 640–649
37. Isago, T., Nozaki, M., Kikuchi, Y. et al. Effects of different negative pressures on reduction of wounds in negative pressure dressings. *J Dermatol.* 2003; 30(8): 596–601.
38. Morykwas, M.J., Faler, B.J., Pearce, D.J., Argenta, L.C. Effects of varying levels of subatmospheric pressure on the rate of granulation tissue formation in experimental wounds in swine. *Ann Plast Surg.* 2001; 47(5): 547–551.
39. Borgquist, O., Ingemansson, R., Malmsjo, M. Wound edge microvascular blood flow during negative-pressure wound therapy: examining the effects of pressures from –10 to –175 mmHg. *Plast Reconstr Surg.* 2010; 125(2): 502–509.
40. Nease, C. Using low pressure, NPWT for wound preparation & the management of split-thickness skin grafts in 3 patients with complex wound. *Ostomy/Wound Management.* 2009; 55(6): 32–42.
41. Zhou, M., Yu, A., Wu, G. et al. Role of different negative pressure values in the process of infected wounds treated by vacuum-assisted closure: an experimental study. *Int Wound J.* 2013; 10(5): 508–515.
42. Wackenfors, A., Sjogren, J., Gustafsson, R. et al. Effects of vacuum-assisted closure therapy on inguinal wound edge microvascular blood flow. *Wound Repair Regen.* 2004; 12(6): 600–606.
43. Bollero, D., Carnino, R., Risso, D et al. Acute complex traumas of the lower limbs: a modern reconstructive approach with negative pressure therapy. *Wound Repair Regen.* 2007;15(4): 589–594.
44. Stannard, J.P., Robinson, J.T., Anderson, E.R. et al. Negative pressure wound therapy to treat hematomas and surgical incisions following highenergy trauma. *J Trauma Inj Infect Crit Care.* 2006; 60(6): 1301–1306
45. Lee, K.N., Ben-Nakhi, M., Park, E.J., Hong, J.P. Cyclic negative pressure wound therapy: an alternative mode to intermittent system. *Int Wound J.* 2013; 2(6): 686–92.

46. Malmjsjo, M., Ingemansson, R. Variable, intermittent and continuous negative pressure wound therapy using foam or gauze: the biological effects on the wound bed including, blood flow, micro and macro deformation, granulation tissue quantity, wound bed character. Symposium on Advanced Wound Care and the Wound Healing Society Meeting; 2010 17-20 April; Orlando, Florida.
47. Borgquist, O., Ingemansson, R., Malmjsjo, M. The effect of intermittent and variable negative pressure wound therapy on wound edge microvascular blood flow. *Ostomy Wound Management*. 2010; 56(3): 60–67.
48. Fujiwara, T., Nishimoto, S., Ishise, H. et al. Influence of continuous or intermittent negative pressure on bacterial proliferation potency in vitro. *J Plast Surg Hand Surg*. 2013; 47(3): 180–184.
49. Ahearn, C. Intermittent NPWT and lower negative pressures--exploring the disparity between science and current practice: a review. *Ostomy/Wound Manage*. 2009; 55(6): 22–28.
50. Falabella, A.F., Carson, P., Eaglstein, W.H., Falanga, V. The safety and efficacy of a proteolytic ointment in the treatment of chronic ulcers of the lower extremity. *J Am Acad Dermatol*. 1998; 39(5): 737–740.
51. Morykwas. M.J., Simpson. J., Pungler, K. et al. Vacuum-assisted closure: state of basic research and physiologic foundation. *Plast Reconstr Surg*. 2006; 117(7S): 121S–126S.
52. Kairinos, N., Solomons, M., Hudson, D.A. Negative-pressure wound therapy I: the paradox of negative-pressure wound therapy. *Plast Reconstr Surg*. 2009; 123(2): 589–600.
53. Kairinos, N., Solomons, M., Hudson, D.A. The paradox of negative pressure wound therapy – in vitro studies. *J Plast Reconstr Aesthet Surg*. 2010; 63(1): 174–179.
54. Webb, L.X. New techniques in wound management: vacuum-assisted wound closure. *J Am Acad Orthop Surg*. 2002; 10(5): 303–311.
55. Young, S.R., Hampton, S., Martin, R. Non-invasive assessment of negative pressure wound therapy using high frequency diagnostic ultrasound: oedema reduction and new tissue accumulation. *Int Wound J*. 2013; 10(4): 383–388.
56. Borgquist, O., Gustafsson, L., Ingemansson, R., Malmjsjo, M. Micro- and macromechanical effects on the wound bed of negative pressure wound therapy using gauze and foam. *Ann Plast Surg*. 2010; 64(6): 789–793.

57. Saxena. V., Hwang, C.W., Huang, S. et al. Vacuum-assisted closure: microdeformations of wounds and cell proliferation. *Plast Reconstr Surg.* 2004; 114(5): 1086–1096.
58. Wilkes, R., Zhao, Y., Cunningham, K. et al. 3D strain measurement in soft tissue: Demonstration of a novel inverse finite element model algorithm on MicroCT images of a tissue phantom exposed to negative pressure wound therapy. *J Mech Behav Biomed Mater.* 2009; 2(3): 272–287.
59. Kremers, L., Kearns, M., Hammon, D. et al. Involvement of mitogen activated protein kinases (MAP Kinases) in increased wound healing during sub-atmospheric pressure (SAP) treatment. *Wound Repair and Regeneration.* 2003; 11(5).
60. Sumpio, B.E., Banes, A.J., Levin, L.G., Johnson, G. Jr. Mechanical stress stimulates aortic endothelial cells to proliferate. *J Vasc Surg.* 1987; 6(3): 252–256.
61. Greene, A.K., Puder, M., Roy, R. et al. Microdeformational wound therapy: effects on angiogenesis and matrix metalloproteinases in chronic wounds of 3 debilitated patients. *Ann Plast Surg.* 2006; 56(4): 418–422
62. McNulty, A., Spranger, I., Courage, J. et al. The consistent delivery of negative pressure to wounds using reticulated, open cell foam and regulated pressure feedback. *Wounds.* 2010; 22(5): 114–120.
63. Li, J., Hampton, T., Morgan, J.P., Simon, M. Stretch-induced VEGF expression in the heart. *J Clin Invest.* 1997; 100(1): 18–24.
64. Shiu, Y.T., Weiss, J.A., Hoying, J.B. et al. The role of mechanical stresses in angiogenesis. *Crit Rev Biomed Eng.* 2005; 33(5): 431–510.
65. Ingber, D.E., Prusty, D., Sun, Z. et al. Cell shape, cytoskeletal mechanics, and cell cycle control in angiogenesis. *J Biomec.* 1995; 28(12): 1471–1484. Medline doi:10.1016/0021-9290(95)00095-X.
66. Von Offenbergsweeney, N., Cummins, P.M., Cotter, E.J. et al. Cyclic strain-mediated regulation of vascular endothelial cell migration and tube formation. *Biochem Biophys Res Commun.* 2005; 329(2): 573–582.
67. Ingber, D.E. Tensegrity: the architectural basis of cellular mechanotransduction. *Annu Rev Physiol.* 1997; 59(1): 575–599.
68. Vandenberg, H.H. Mechanical forces and their second messengers in stimulating cell growth in vitro. *Am J Physiol.* 1992; 262(3): R350–R355.

69. Sadoshima J, Takahashi T, Jahn L, Izumo S. Roles of mechano-sensitive ion channels, cytoskeleton, and contractile activity in stretch-induced immediateearly gene expression and hypertrophy of cardiac myocytes. *Proc Natl Acad Sci USA*. 1992; 89(20): 9905–9909
70. Baudouin-Legros, M., Paquet, J.L., Brunelle, G., Meyer, P. Role of nuclear proto-oncogenes in the proliferation of aortic smooth muscle cells in spontaneously hypertensive rats. *J Hypertens*. 1989; 7(6): S114–S115.
71. Folkman, J., Moscona, A. Role of cell shape in growth control. *Nature*. 1978; 273(5661): 345–349.
72. Younan, G., Ogawa, R., Ramirez, M. et al. Analysis of nerve and neuropeptide patterns in vacuum-assisted closure-treated diabetic murine wounds. *Plast Reconstr Surg*. 2010; 126(1): 87–96.
73. Ingber, D.E., Huang, S. The structural and mechanical complexity of cell-growth control. *Nat Cell Biol*. 1999; 1(5): E131–E138.
74. Chen, C.S., Mrksich, M., Huang, S. et al. Geometric control of cell life and death. *Science*. 1997; 276(5317): 1425–1428.
75. Sumpio, B.E., Banes, A.J. Response of porcine aortic smooth muscle cells to cyclic tensional deformation in culture. *J Surg Res*. 1988; 44(6): 696–701
76. Lohman, R.F., Lee, R.C. Discussion: vacuum assisted closure: microdeformations of wounds and cell proliferation. *Plast Reconstr Surg*. 2004; 114(5): 1097–1098.
77. Morykwas, M.J., Argenta, L.C. Nonsurgical modalities to enhance healing and care of soft tissue wounds. *J South Orthop Assoc*. 1997; 6(4): 279–288.
78. Rejzek. A., Weyer, F. The use of the V.A.C.TM system in the therapy of ulcer cruris and diabetic gangrene. *Acta Chir Austriaca. Supplement*. 1998; 150: 12–13.
79. De Lange, M.Y., Nicolai, J.P. The influence of subatmospheric pressure on tissue oxygenation and temperature. *Abstract volume 2nd WUWHS Meeting*. 2004; Paris 2004: 51.
80. Kamolz, L.P., Andel, H., Haslik, W. et al. Use of subatmospheric pressure therapy to prevent burn wound progression in human: first experiences. *Burns*. 2004; 30(3): 253–258.

81. Banwell, P.E., Morykwas, M.J., Jennings, D.A. Dermal microvascular blood flow in experimental partial thickness burns: the effect of topical subatmospheric pressure. *J Burn Care Rehabil.* 2000; 21:161.
82. Chen, S.Z., Li, J., Li, X.Y., Xu, L.S. Effects of vacuum-assisted closure on wound microcirculation: an experimental study. *Asian J Surg.* 2005; 28(3): 211–217.
83. Wackenfors, A., Sjogren, J., Gustafsson, R. et al. Effects of vacuum-assisted closure therapy on inguinal wound edge microvascular blood flow. *Wound Repair Regen.* 2004; 12(6): 600–606.
84. Wackenfors, A., Gustafsson, R., Sjogren, J. et al. Blood flow responses in the peristernal thoracic wall during vacuum-assisted closure therapy. *Ann Thorac Surg.* 2005; 79(5): 1724–1730.
85. Petzina, R., Gustafsson, L., Mokhtari, A. et al. Effect of vacuum-assisted closure on blood flow in the peristernal thoracic wall after internal mammary artery harvesting. *Eur J Cardiothorac Surg.* 2006; 30(1): 85–89.
86. Petzina, R., Ugander, M., Gustafsson, L. et al. Topical negative pressure therapy of a sternotomy wound increases sternal fluid content but does not affect internal thoracic artery blood flow: Assessment using magnetic resonance imaging. *J Thorac Cardiovasc Surg.* 2008; 135(5): 1007–1013.
87. Ichioka, S., Watanabe, H., Sekiya, N. et al. A technique to visualize wound bed microcirculation and the acute effect of negative pressure. *Wound Repair Regen.* 2008; 16(3): 460–465.
88. Horch, R.E., Muchow, S., Dragu, A. Erste Zwischenergebnisse der Perfusionsbeeinflussung durch Prevena: Gewebsperfusion. *Dzf.* 2012;16:1–3.
89. Hudlicka, O., Brown, M., Egginton, S. Angiogenesis in skeletal and cardiac muscle. *Physiol Rev.* 1992; 72(2): 369–417.
90. Ping, A., Zhang, T., Ren, B. et al. Effect of vacuum-assisted closure combined with open bone grafting to promote rabbit bone graft vascularization. *Med Sci Monit.* 2015;21:1200–1206. Medline doi:10.12659/MSM.892939.
91. Genecov, D.G., Schneider, A.M., Morykwas, M.J. et al. A controlled subatmospheric pressure dressing increases the rate of skin graft donor site reepithelialization. *Ann Plast Surg.* 1998;40(3): 219–225. Medline doi:10.1097/00000637-199803000-00004.

92. Erba, P., Ogawa, R., Ackermann, M. et al. Angiogenesis in wounds treated by microdeformational wound therapy. *Ann Surg.* 2011; 253(2): 402–409. Medline doi:10.1097/SLA.0b013e31820563a8.
93. Deva, A.K., Buckland, G.H., Fisher, E. et al. Topical negative pressure in wound management. *Med J Aust.* 2000; 173(3): 128–131.
94. Fleck, T.M., Fleck, M., Moidl, R. et al. The vacuum-assisted closure system for the treatment of deep sternal wound infections after cardiac surgery. *Ann Thorac Surg.* 2002; 74(5): 1596–1600.
95. Moues, C.M., Vos, M.C., Van Den Bemd, G.J. et al. Bacterial load in relation to vacuum-assisted closure wound therapy: A prospective randomized trial. *Wound Repair Regen.* 2004; 12(1): 11–17.
96. Weed, T., Ratliff, C., Drake, D.B. Quantifying bacterial bioburden during negative pressure wound therapy: does the wound VAC enhance bacterial clearance? *Ann Plast Surg.* 2004; 52(3): 276–279.
97. James, G.A., Swogger, E., Wolcott, R. et al. Biofilms in chronic wounds. *Wound Repair Regen.* 2008; 16(1): 37–44.
98. Gouttefangeas, C., Eberle, M., Ruck, P. et al. Functional T lymphocytes infiltrate implanted polyvinyl alcohol foams during surgical wound closure therapy. *Clin Exp Immunol.* 2001; 124(3): 398–405.
99. Adams, T.S., Herrick, S., McGrouther, D.A. VAC-therapy alters the number and distribution of neutrophils in acute dermal wounds. In: Topical negative pressure (TNP) focus group meeting (Proceedings European Tissue Repair Society). TXP Communications: 212; 2004.
100. Buttenschoen, K., Fleischmann, W., Haupt, U. et al. The influence of vacuum-assisted closure on inflammatory tissue reactions in the postoperative course of ankle fractures. *Foot Ankle Surg.* 2001; 7(3): 165–173.
101. Kilpadi, D.V., Bower, C.E., Reade, C.C. et al. Effect of Vacuum Assisted Closure Therapy on early systemic cytokine levels in a swine model. *Wound Repair Regen.* 2006; 14(2): 210–215.
102. Labler, L., Mica, L., Harter, L. et al. Influence of V.A.C.-therapy on cytokines and growth factors in traumatic wounds. *Zentralbl Chir.* 2006;131: S62–S67.

103. Fraccalvieri, M., Scalise, A., Ruka, E. et al. Negative pressure wound therapy using gauze and foam: histological, immunohistochemical, and ultrasonography morphological analysis of granulation and scar tissues. *Eur J Plast Surg*. 2014; 37(8): 411–416.
104. Fraccalvieri, M., Zingarelli, E., Ruka, E. et al. Negative pressure wound therapy using gauze and foam: histological, immunohistochemical and ultrasonography morphological analysis of the granulation tissue and scar tissue. Preliminary report of a clinical study. *Int Wound J*. 2011; 8(4): 355–364.
105. Jeffery, SL. Advanced wound therapies in the management of severe military lower limb trauma: a new perspective. *Eplasty*. 2009; 9.
106. Hurd, T., Chadwick, P., Cote, J. et al. Impact of gauze-based NPWT on the patient and nursing experience in the treatment of challenging wounds. *Int Wound J*. 2010; 7(6): 448–455.
107. Fraccalvieri, M., Ruka, E., Bocchiotti, M.A. et al. Patient's pain feedback using negative pressure wound therapy with foam and gauze. *Int Wound J*. 2011; 8(5): 492–499.
108. Malmsjo, M., Gustafsson, L., Lindstedt, S., Ingemansson, R. Negative pressure wound therapy-associated tissue trauma and pain: a controlled in vivo study comparing foam and gauze dressing removal by immunohistochemistry for substance P and calcitonin gene-related peptide in the wound edge. *Ostomy/ Wound Management*. 2011; 57(12): 30–35.