

DAFTAR PUSTAKA

- Ab Wahab, S. Z. Kadir, A. A., Hussain, N. H. N, Omar, J., Yunus, R., Baie, S., Noor, N. M., Hassan, I. I., Mahmood, W. H. W., Razak, A. A., Yusoff, W. Z. W. 2015. The Effect of *Channa striatus* (Haruan) Extract on Pain and Wound Healing of Post-Lower Segment Caesarean Section Women. *Evidence-based Complementary and Alternative Medicine*. Hindawi Publishing Corporation. p. 16.
- Allen, G. R. 1991. Field guide to the freshwater fishes of New Guinea. *Christensen Research Institute*, Madang, Papua New Guinea. 9, p. 268.
- Bailey, R. 2019. *Genetic Variation Definition, Causes, and Examples* [online]. [Cited 28 September 2019]. <https://www.thoughtco.com/genetic-variation-373457>
- Brox, T. 2014. Maximum Likelihood Estimation. In K. Ikeuchi, ed. *Computer Vision: A Reference Guide*, Boston, MA, Springer US. pp. 481–482.
- Cahyono, B. 2000. *Budidaya Ikan Air Tawar: Ikan Gurami, Ikan Nila, Ikan Mas*. Kanisius. Yogyakarta. p. 9.
- Dawkins, R. 1994. The eye in a twinkling. *Nature*, 368(6473): 690–691.
- Giacomello, E. and Marchini, D. 2006. A male sexually dimorphic trait provides antimicrobials to eggs in blenny fish. *Biology Letters*, 2(3): 330–333.
- Ilmi, W. and Arisuryanti, T. 2018. Composition of Mitochondrial DNA 16S Nucleotide of Dwarf Snakehead (*Channa gachua* Hamilton, 1822) from Keji River, Magelang, Central Java. *Journal of Tropical Biodiversity and Biotechnology*, 3(2): 57.
- de Jong, M.A., Wahlberg, N., van Eijk, M., Brakefield, P.M. & Zwaan, B.J. 2011. Mitochondrial DNA signature for range-wide populations of *Bicyclus anynana* suggests a rapid expansion from recent refugia. *PLoS ONE*, 6(6): 1–5.
- Jamaluddin, J. A. F., Pau, T. M., Siti-Azizah, M. N. 2011. Genetic structure of the snakehead murrel, *Channa striata* (channidae) based on the *cytochrome c oxidase subunit i* gene: Influence of historical and geomorphological factors. *Genetics and Molecular Biology*, 34(1): 152–160.
- Kombong, C.B.S. and Arisuryanti, T. 2019. The 16S and COI Mitochondrial DNA Nucleotide Composition of Stripped Snakehead (*Channa striata* Bloch, 1793) from Lake Sentani, Jayapura, Papua. *Jurnal Perikanan Universitas Gadjah Mada*, 20(2): 57.
- Kumar, S., Stecher, G. and Tamura, K. 2016. MEGA7 : Molecular Evolutionary Genetics Analysis Version 7 . 0 for Bigger Datasets Brief communication. , 33(7): 1870–1874.
- Leigh, J.W. and Bryant, D. 2015. Application POPART: full-feature software for haplotype network construction. *Methods in Ecology and Evolution* 6: 1110–1116
- Li, J., Wang, X., Kong, X., Zhao, K., He, S. and Mayden, R.L. 2008. Variation patterns of the mitochondrial 16S rRNA gene with secondary structure constraints and their application to phylogeny of cyprinine fishes (Teleostei: Cypriniformes). *Molecular Phylogenetics and Evolution*, 47(2): 472–487.
- Librado, P. and Rozas, J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25(11): 1451–1452.

- Listyanto, N. and Andriyanto, S. 2009. IKAN GABUS (*Channa striata*) MANFAAT PENGEMBANGAN DAN ALTERNATIF TEKNIK BUDIDAYANYA. *Media Akuakultur*, 4(1): 18.
- Maddison, W. P. and D.R. Maddison. 2018. Mesquite: a modular system for evolutionary analysis. Version 3.5 <http://www.mesquiteproject.org>.
- Mustafa, A., Widodo, M. A., Kristianto, Y. 2012. Albumin And Zinc Content Of Snakehead Fish (*Channa striata*) Extract And Its Role In Health. *IEESE International Journal Of Science and Technology*. 1(2): 1–8.
- Muthmainnah, D. 2016. Growout of Striped Snakehead (*Channa Striata*) in Swamp Water System Using Fences and Cages. *Romanian Biotechnological Letters*. 21(2): 11298–11303.
- Nei, M. and Li, W.H. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Sciences of the United States of America*, 76(10): 5269–5273.
- Nurdawati, S., Muflikhah, N. and Sunarno, M.T.D. 2017. Sumber Daya Perikanan Perairan Sungai Batang Hari Jambi. *BAWAL Widya Riset Perikanan Tangkap*, 1(1): 1.
- Nishimaki, T. and Sato, K. 2019. An Extension of the Kimura Two-Parameter Model to the Natural Evolutionary Process. *Journal of Molecular Evolution*, 87(1): 60–67.
- Ramadhaniaty, M., Setyobudiandi, I. & Madduppa, H.H. 2018. Morphogenetic and population structure of two species marine bivalve (Ostreidae: *Saccostrea cucullata* and *Crassostrea iredalei*) in aceh, Indonesia. *Biodiversitas*, 19(3): 978–988.
- Makmur S., Rahardjo M. F., and Sutrisno Sukimin. 2003. Biologi reproduksi ikan gabus (*Channa striata* bloch) di daerah banjir sungai musu sumatera selatan. *Jurnal Ikhtiologi Indonesia*. 3(2): 58-62.
- Saitou, N. and Nei, M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4): 406–425.
- Supriatna, J. 2008. *Melestarikan Alam Indonesia*. Yayasan Obor Indonesia. Jakarta. p. 391.
- Yamamoto, M.N. and A.W. Tagawa, 2000. *Hawai'i's native and exotic freshwater animals*. Mutual Publishing, Honolulu, Hawaii. p. 200.
- Van der Laan, R., Fricke, R., and Eschmeyer, W. N. 2019. Eschmeyer's Catalog Of Fishes: Classification. (<http://www.calacademy.org/scientists/catalog-of-fishes-classification/>). *Electronic version accessed* 19 September 2019.