

REFERENCES

- AOAC. (2002). Official Methods of Analysis. In *Association of Official Analytical Chemists* (16 ed.). Washington, DC.
- Ando, H., Adachi, M., Umeda, K., Matsuura, A., Nonaka, M., Uchio, R., *et al.* (1989). Purification and characteristics of a novel transglutaminase derived from microorganisms. *Agricultural and Biological Chemistry*, 53, 2613–2617.
- Antolovich, M., Prenzler, P. D., Patsalides, E., McDonald, S., and Robards, K. (2001). Methods for testing antioxidant activity. *The Analyst*, 127, 183–198.
- Alvarez, C. and Tejada, M. 1997. Influence of texture of suwari gels on kamaboko gels made from sardine (*Sardina pilchardus*) surimi. *Journal of Science and Food Agriculture*, 75: 472-480.
- Arfat, Y. and Benjakul, S. (2012). Impact of zinc salts on heat-induced aggregation of natural actomyosin from yellow stripe trevally. *Food Chemistry*. 135, 2721–2727.
- Bagagli, M. P. (2009). Produção de transglutaminase de *Streptomyces* sp. P20, caracterização e aplicação da enzima bruta. Campinas: Universidade Estadual de Campinas
- Balange, A. and Benjakul, S. (2009). Enhancement of gel strength of bigeye snapper (*Priacanthus tayenus*) surimi using oxidised phenolic compounds. *Food Chemistry*. 113, 61–70.
- Benjakul, S., Chantarasuwan, C. and Visessanguan, W. (2003). Effect of medium temperature setting on gelling characteristics of surimi from some tropical fish. *Food Chemistry*, 82, 567-574.
- Benjakul, S. and Visessanguan, W. (2003). Transglutaminase mediated setting in bigeye snapper surimi. *Food Research International*. 36, 253–266.
- Benjakul, S., Visessanguan, W., and Pecharat, S. (2004). Suwari gel properties as affected by transglutaminase activator and inhibitors. *Food Chemistry*. 85, 91-99.
- Benjakul, S., Visessanguan, W., Thongkaew, C., and Tanaka, M. (2005). Effect of frozen storage on chemical and gel-forming properties of fish commonly used for surimi production in Thailand. *Food Hydrocolloids*, 19, 197–207.

- Benjakul, S., Visessanguan, V., and Srivilai, C. (2007). Gel properties of bigeye snapper (*Priacanthus tayenus*) surimi as affected by setting and plasma proteins. *Journal of Food Quality*, 24, 453-471.
- Benjakul, S., Phatcharat, S., Tammatinna, A., Visessanguan, W., and Kishimura, H. (2008). Improvement of Gelling Properties of Lizardfish Mince as Influenced by Microbial Transglutaminase and Fish Freshness. *Journal of Food Science*, 73, S239–S246.
- Benjakul, S., Yarnpakdee, S., and Phatcharat, S. (2010). Combination Effects Of Whey Protein Concentrate and Calcium Chloride on the Properties of Goatfish Surimi Gel. *Journal of Texture Studies*, 41, 341–357.
- Buamard, N., and Benjakul, S. (2015). Food Hydrocolloids Improvement of gel properties of sardine (*Sardinella albella*) surimi using coconut husk extracts. *Food Hydrocolloids*, 51, 146–155.
- Buamard, N., and Benjakul, S. (2017). Ethanolic coconut husk extract: In vitro antioxidative activity and effect on oxidative stability of shrimp oil emulsion. *European Journal of Lipid Science and Technology*, 119, 1700131.
- Carvalho, O. V., Botelho, C. V., Ferreira, C. G. T., Ferreira, H. C. C., Santos, M. R., Diaz, M. A. N., *et al.* (2013). *In vitro* inhibition of canine distemper virus by flavonoids and phenolic acids: Implications of structural differences for antiviral design. *Research in Veterinary Science*, 95, 717-724.
- Chaijan, M., Benjakul, S., Visessanguan, W., and Faustman, C. (2004). Characteristics and gel properties of muscles from sardine (*Sardinella gibbosa*) and mackerel (*Rastrelliger kanagurta*) caught in Thailand. *Food Research International*, 37, 1021-1030.
- Chalamaiah, M., Dinesh kumar, B., Hemalatha, R., and Jyothirmayi, T. (2012). Fish protein hydrolysates: Proximate composition, amino acid composition, antioxidant activities and applications: A review. *Food Chemistry*, 135(4), 3020–3038.
- Chan, J. K., Gill, T. A. and Paulson, A. T. 1992. Cross-linking of myosin heavy chains from cod, herring and silver hake during thermal setting. *Journal of Food Science* 57, 906-912.

- Chanarat, S., and Benjakul, S. (2013). Impact of microbial transglutaminase on gelling properties of Indian mackerel fish protein isolates. *Food Chemistry*, 136, 929–937.
- Chanarat, S., Benjakul, S. and H-Kittikun, A. (2012). Comparative study on protein cross-linking and gel enhancing effect of microbial transglutaminase on surimi from different fish. *Journal of the Science of Food and Agriculture*. 92, 844–852.
- Chanarat, S., and Benjakul, S. (2013). Impact of microbial transglutaminase on gelling properties of Indian mackerel fish protein isolates. *Food Chemistry*, 136, 929–937.
- Craft, B. D., Kerrihard, A. L., Amarowicz, R., and Pegg, R. B. (2012). Phenol-Based Antioxidants and the in Vitro Methods Used for Their Assessment. *Comprehensive Reviews in Food Science and Food Safety*, 11, 148–173.
- De Jong, G. A. H., and Koppelman, S. J. (2002). Transglutaminase catalyzed reactions: Impact on food applications. *Journal of Food Science*, 67, 2798–2806.
- Dube, M., Schäfer, C., Neidhart, S., and Carle, R. (2007). Texturisation and modification of vegetable proteins for food applications using microbial transglutaminase. *European Food Research and Technology*, 225, 287–299.
- Edenharder, R., and Grünhage, D. (2003). Free radical scavenging abilities of flavonoids as mechanism of protection against mutagenicity induced by tert-butyl hydroperoxide or cumene hydroperoxide in *Salmonella typhimurium* TA102. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 540, 1–18.
- Erwanto, Y., Kawahara, S., Katayama, K., Takenoyama, S., Fujino, H., and Yamauchi, K. (2003). Microbial transglutaminase modifies gel properties of porcine collagen. *Asian-Australasian Journal of Animal Sciences*, 6, 269-276.
- Esfandi, R., Walters, M. E., and Tsopmo, A. (2019). Antioxidant properties and potential mechanisms of hydrolyzed proteins and peptides from cereals. *Heliyon*, 5, e01538.
- Folk, J. E., and Finlayson, J. S. (1977). The ϵ -(γ -glutamyl) lysine cross-link and the catalytic role of transglutaminase. *Advances in Protein Chemistry*, 31, 1–133.
- Gawlik-Dziki, U., Jezyna, M., Swieca, M., Dziki, D., Baraniak, B., and Czyz, J. (2012). Effect of bioaccessibility of phenolic compounds on *in vitro* anticancer activity of broccoli sprouts. *Food Research International*, 49, 469-476.

- Griffin, M., Casadio, R. and Bergamini, C.M. (2002). Transglutaminases: Nature's biological glues. *Biochemical Journal*, 368, 377–396.
- Han, M., Zhang, Y., Fei, Y., Xu, X., and Zhou, G. (2009). Effect of microbial transglutaminase on NMR relaxometry and microstructure of pork myofibrillar protein gel. *European Food Research and Technology*, 228, 665–670.
- Hernández-Briones A., Velázquez, G. Vázquez, M. and Ramírez, J. A. (2009). Effects of adding fish gelatin on Alaska pollock surimi gels. *Food Hydrocolloids*, 23, 2446–2449.
- Hossain, M. I., Itoh, Y., Morioka, K., and Obatake, A. (2001). Contribution of the polymerization of protein by disulfide bonding to increased gel strength of walleye pollack surimi gel with preheating time. *Fisheries Science*, 67, 710-717.
- Hoyos, M. N., Sánchez-Patán, F., Murillo Masis, R., Martín-Álvarez, P., Zamora Ramirez, W., Monagas, M., and Bartolomé, B. (2015). Phenolic Assesment of *Uncaria tomentosa* L. (Cat's Claw): Leaves, Stem, Bark and Wood Extracts. *Molecules*, 20, 22703–22717.
- Hsieh, J.F. Tsai, G.J., and Jiang, S.T. (2006). Microbial transglutaminase and recombinant cystatin effects on improving the quality of mackerel surimi. *Journal of Food Science*, 67, 3120–3125.
- Hu, Y. Liu, W. Yuan, C. Morioka, K. Chen, S. Liu, D. and Ye, X. (2015). Enhancement of the gelation properties of hairtail (*Trichiurus haumela*) muscle protein with curdlan and transglutaminase. *Food Chemistry*. 176, 115–122.
- Jiang, S.T., Hsieh, J.F., Ho, M.L. and Chung, Y.C. (2000). Microbial transglutaminase affects gel properties of golden threadfin bream and pollack surimi. *Journal of Food Science*, 65, 694-699.
- Ju, Z. Y., and Kilara, A. (1998). Aggregation Induced by Calcium Chloride and Subsequent Thermal Gelation of Whey Protein Isolate. *Journal of Dairy Science*, 8, 925–931.
- Kaewudom, P., Benjakul, S., and Kijroongrojana, K. (2013). Properties of surimi gel as influenced by fish gelatin and microbial transglutaminase. *Food Bioscience*, 1, 39-47
- Kallel, F., Driss, D., Chaari, F., Belghith, L., Bouaziz, F., Ghorbel, R., *et al.* (2014). Garlic (*Allium ativum* L.) husk waste as a potential source of phenolic compounds: Influence of extracting solvents on its antimicrobial and antioxidant properties. *Industrial Crops and Products*, 62, 34-41.

- Klomklao, S., Kishimura, H., and Benjakul, S. (2008). Endogenous proteinases in true sardine (*Sardinops melanostictus*). *Food Chemistry*, 107, 213–220.
- Kumazawa Y, Numazawa T, Seguro K, Motoki M. (1995). Suppression of surimi gel setting by transglutaminase inhibitors. *Journal of Food Science*. 60, 715–17
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of head of bacteriophage T4. *Nature*, 227, 680-685.
- Lanier, T.C. (2000). Surimi gelation chemistry. In *Surimi and Surimi Seafood* (J.W. Park, ed.) pp. 237–265, Marcel Dekker, New York.
- Lee EY and Park J. (2003). Microbial transglutaminase induced cross-linking of a selected comminuted muscle system processing conditions for physical properties of restructured meat. *Food Science and Biotechnology*, 12, 365– 70
- Liao, K., and Yin, M. (2000). Individual and Combined Antioxidant Effects of Seven Phenolic Agents in Human Erythrocyte Membrane Ghosts and Phosphatidylcholine Liposome Systems: Importance of the Partition Coefficient. *Journal of Agricultural and Food Chemistry*, 48, 2266–2270.
- Lobbes, J. M., Fitznar, H. P., and Kattner, G. (1999). High-performance liquid chromatography of lignin-derived phenols in environmental samples with diode array detection. *Analytical Chemistry*, 71, 3008-3012.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193, 265-275.
- Macedo, J., and Sato, H. (2005). Propriedades e aplicações da transglutaminase microbiana em alimentos. *Alimentos e Nutrição*, 16, 413–419.
- Martins, N., Barros, L., Santos-Buelga, C., and Ferreira, I. C. F. R. (2016). Antioxidant potential of two Apiaceae plant extracts: A comparative study focused on the phenolic composition. *Industrial Crops and Products*, 79, 188–194.
- Maqsood, S., Benjakul, S., and Balange, A. K. (2012). Effect of tannic acid and kiam wood extract on lipid oxidation and textural properties of fish emulsion sausages during refrigerated storage. *Food Chemistry*, 130, 408-416.

- Maqsood, S., Kittiphattanabawon, P., and Sumpavapol, P. (2015). Antioxidant activity of date (*Phoenix dactylifera* var . Khalas) seed and its preventive effect on lipid oxidation in model systems Antioxidant activity of date (*Phoenix dactylifera* var . Khalas) seed and its preventive effect on lipid oxidation in mod. *International Food Research Journal*, 22(, 1180–1188.
- Meilgaard, M., Civille, G. V., and Carr, B. T. (1999). *Sensory evaluation techniques* (3rd ed.). Florida: CRC Press. 1-464
- Mleko, S., and Foegeding, E. A. (2000). pH Induced Aggregation and Weak Gel Formation of Whey Protein Polymers. *Journal of Food Science*, 65(1), 139–143.
- Motoki, M., and Seguro, K. (1998). Transglutaminase and its use for food processing. *Trends in Food Science and Technology*, 9, 204–210.
- Nakahara, C., Nozawa, H., and Seki, N. (1999). A Comparison of Cross-linking of Fish Myofibrillar by Endogenous and Microbial Transglutaminases. *Fisheries Science*, 65, 138–144.
- O’Connell, J.E. and Fox, P.F. (2001). Significance and applications of phenolic compounds in the production and quality of milk and dairy products. *International Dairy Journal*. 11: 103-120.
- Oteng-Pabi, S. K., Pardin, C., Stoica, M., and Keillor, J. W. (2014). Site-specific protein labelling and immobilization mediated by microbial transglutaminase. *Chem. Commun*, 50, 6604–6606.
- Ozdal, T., Capanoglu, E. and Altay, F. (2013). A review on protein–phenolic interactions and associated changes. *Food Research International*, 51, 954–970.
- Park, J.W., (2010). *Surimi and Surimi Seafood*, Marcel Dekker, Newyork, 2000, pp. 237–265.
- Parkin, K. L. Enzimas. In S. Damodaran, K. L. Parkin, and O. R. Fennema (Eds.) (2010). *Química de alimentos de Fennema* (4th ed., pp. 263–342). Porto Alegre: Artmed.
- Panyakaew, S. and Fotios, S. 2011. New thermal insulation boards made from coconut husk and bagasse. *Energy Buildings*, 43, 1732–1739.

Pereira, D., Valentão, P., Pereira, J., and Andrade, P. (2009). Phenolics: From Chemistry to Biology. *Molecules*, 14, 2202–2211.

Prodpran, T., Benjakul, S., and Phatcharat, S. (2012). Macromolecules Effect of phenolic compounds on protein cross-linking and properties of film from fish myofibrillar protein. *International Journal of Biological Macromolecules*, 51, 774–782.

Ramadan H, Byujing Min, Amit K and Tiwari, G. R. (2015). Antibacterial activity of pomegranate, orange and lemon peel extracts against food-borne pathogens and spoilage bacteria *in vitro* and on poultry skin. *Journal of Poultry Sciences*, 14, 29-39.

Rawdkuen, S. and Benjakul, S. (2008). Whey protein concentrate: Autolysis inhibition and effects on the gel properties of surimi prepared from tropical fish. *Food Chemistry*, 106, 1077–1084.

Rattaya, S., Benjakul, S., and Prodpran, T. (2009). Properties of fish skin gelatin film incorporated with seaweed extract. *Journal of Food Engineering*, 95, 151-157.

Robinson, H. W., and Hogden, C. G. (1940). The biuret reaction in the determination of serum proteins. I. A study of the conditions necessary for the production of a stable color which bears a quantitative relationship to the protein concentration. *Journal of Biological Chemistry*, 135, 707-725.

Rodrigues, S., and Pinto, G. A. S. (2007). Ultrasound extraction of phenolic compounds from coconut (*Cocos nucifera*) shell powder. *Journal of Food Engineering*, 80, 869-872.

Sakamoto, H., Kumazawa, Y., Toiguchi, S., Seguro, K., Soeda, T. and Motoki, M. (2006). Gel strength enhancement by addition of microbial transglutaminase during onshore surimi manufacture. *Journal of Food Science*, 60, 300–304.

Samejima, K., Ishioroshi, M. and Yasui, T. (1981). Relative roles of the head and tail portions of the molecule in the heat-induced gelation of myosin. *Journal of Food Science*. 46, 1412-1418

Sano, T., Ohno, T., Otsuka, F. H., Matsumoto, J. J., and Tsuchiya, T. (1994). Carp natural actomyosin: thermal denaturation mechanism. *Journal of Food Science*, 59, 1002-1008.

Seguro K, Nio N, and Motoki M. (1996). Some characteristics of a microbial protein cross-linking enzyme: Transglutaminase. In *Macromolecular Interactions in Food Technology*,

- Steel, R. G. D., and Torrie, J. H. (1980). Principles and procedures of statistics: A biometrical approach (2nd ed.). New York: McGraw-Hill.
- Singh, A., and Benjakul, S. (2018). Proteolysis and its control using protease inhibitors in fish and fish products: a review, 17, 496-509.
- Turan, H., and Sönmez, G. (2007). Changes in the quality of surimi made from thornback ray (*Raja clavata*, L. 1758) during frozen storage. International Journal of Food Sciences and Nutrition, 58, 557–566.
- Téllez-Luis SJ, González-Cabriaes JJ, Ramírez JA, Vázquez M (2004) Production of transglutaminase by *Streptovorticillium ladakanum* NRRL-3191 grown on media made from hydrolysates of sorghum straw. Food Technology Biotechnology 42:1–4
- Woods, M. N., and Gorbach, S. L. (2001). Influences of fiber on the ecology of the intestinal flora. In G. A. Spiller (Ed.), CRC Handbook of dietary fiber in human nutrition (pp. 257-270). Florida: CRC Press.
- Vazquez-Torres, H., Canch, E-Escamilla, G. and Cruz-Ramos, C.A. (1992). Coconut husk lignin. II. Characterization by infrared and nuclear magnetic resonance spectroscopy. Journal of Applied Polymer Science. 45, 645–653.
- Yang, Z. (1997). Process Development for Microbial Transglutaminase Production. pp 1-7.
- Yokoyama, K., Nio, N., and Kikuchi, Y. (2004). Properties and applications of microbial transglutaminase. Applied Microbiology and Biotechnology, 64, 447–454.
- You, L., Zhao, M., Regenstein, J. M., and Ren, J. (2010). Changes in the antioxidant activity of loach (*Misgurnus anguillicaudatus*) protein hydrolysates during asimulated gastrointestinal digestion. Food Chemistry, 120, 810–816.
- Zhang, F., Fang, L., Wang, C., Shi, L., Chang, T., Yang, H., and Cui, M. (2013). Effects of starches on the textural, rheological, and color properties of surimi–beef gels with microbial tranlglutaminase. Meat Science, 93, 533–537.