

DAFTAR PUSTAKA

Alge, J. L. and Arthur, J. M. (2015) 'Biomarkers of AKI: A review of mechanistic relevance and potential therapeutic implications', *Clinical Journal of the American Society of Nephrology*, 10(1), hal. 147–155. doi: 10.2215/CJN.12191213.

Birn, H. *et al.* (2000) 'Cubilin is an albumin binding protein important for renal tubular albumin reabsorption', *Journal of Clinical Investigation*. The American Society for Clinical Investigation, 105(10), hal. 1353–1361. doi: 10.1172/JCI8862.

Brunvand, L. *et al.* (2017) 'Advanced glycation end products in children with type 1 diabetes and early reduced diastolic heart function'. doi: 10.1186/s12872-017-0551-0.

Christensen, E. I. *et al.* (2012) 'Endocytic receptors in the renal proximal tubule', *Physiology (Bethesda)*, 27, hal. 223–236.

Christensen, E. I. and Birn, H. (2002) 'Megalin and cubilin: Multifunctional endocytic receptors', *Nature Reviews Molecular Cell Biology*, 3(4), hal. 258–268. doi: 10.1038/nrm778.

Corbacioglu, S. K. *et al.* (2017) 'Value of plasma neutrophil gelatinase-associated lipocalin (NGAL) in distinguishing between acute kidney injury (AKI) and chronic kidney disease (CKD)', *Turkish Journal of Emergency Medicine*, 17(3), hal. 85–88. doi: 10.1016/j.tjem.2017.03.002.

De, S., Kuwahara, S. and Saito, A. (2014) 'membranes The Endocytic Receptor Megalin and its Associated Proteins in Proximal Tubule Epithelial Cells', *Membranes*, 4, hal. 333–355. doi: 10.3390/membranes4030333.

Figueira, M. F. *et al.* (2017) 'Diabetic rats present higher urinary loss of proteins and lower renal expression of megalin, cubilin, CIC-5, and CFTR', *Physiological Reports*, 15(13), hal. 1–13. doi: 10.14814/phy2.13335.

Forbes, J. M. (2003) 'Role of Advanced Glycation End Products in Diabetic Nephropathy', *Journal of the American Society of Nephrology*, 14(90003), hal. 254S–258. doi: 10.1097/01.ASN.0000077413.41276.17.

Gilbert, R. E. (2017) 'Proximal Tubulopathy: Prime Mover and Key Therapeutic Target in Diabetic Kidney Disease', *Diabetes*, 66, hal. 791–800. doi: 10.2337/db16-0796.

Hovind, P. *et al.* (2003) 'Decreasing incidence of severe diabetic microangiopathy in type 1 diabetes', *Diabetes Care*, 26(4), hal. 1258–1264. doi: 10.2337/diacare.26.4.1258.

Hvidberg, V. *et al.* (2005) 'The endocytic receptor megalin binds the iron transporting neutrophil-gelatinase-associated lipocalin with high affinity and mediates its cellular uptake', *FEBS Letters*, 579(3), hal. 773–777. doi: 10.1016/j.febslet.2004.12.031.

KEMENKES (2018) ‘Hasil Utama Riskesdas Tentang Prevalensi Diabetes Mellitus di Indonesia 2018’, *Hasil Utama Riskesdas Tentang Prevalensi Diabetes Mellitus di Indonesia 2018*, hal. 8.

Liu, D. *et al.* (2015) ‘Megalin/cubulin-lysosome-mediated albumin reabsorption is involved in the tubular cell activation of NLRP3 inflammasome and tubulointerstitial inflammation’, *Journal of Biological Chemistry*. doi: 10.1074/jbc.M115.662064.

Luo, Q. H. *et al.* (2016) ‘Evaluation of KIM-1 and NGAL as Early Indicators for Assessment of Gentamycin-Induced Nephrotoxicity in Vivo and in Vitro’, *Kidney and Blood Pressure Research*. S. Karger AG, 41(6), hal. 911–918. doi: 10.1159/000452592.

Pieters, T. T. *et al.* (2019) ‘Histological characteristics of Acute Tubular Injury during Delayed Graft Function predict renal function after renal transplantation’, *Physiological Reports*. American Physiological Society, 7(5). doi: 10.14814/phy2.14000.

Saito, A. *et al.* (2010) ‘Molecular Mechanisms of Receptor-Mediated Endocytosis in the Renal Proximal Tubular Epithelium’, *Journal of Biomedicine and Biotechnology*. Hindawi Publishing Corporation, 2010. doi: 10.1155/2010/403272.

Si Nga, H. *et al.* (2015) ‘Sepsis and AKI in Clinical Emergency Room Patients: The Role of Urinary NGAL’, *BioMed Research International*, 2015. doi: 10.1155/2015/413751.

‘WHO | Country and regional data on diabetes’ (2011) WHO. World Health Organization. Available at: http://www.who.int/diabetes/facts/world_figures/en/ (Diakses pada: 4 September 2019).

Wild, S. *et al.* (2004) ‘Global Prevalence of Diabetes: Estimates for the year 2000 and projections for 2030’, *Diabetes Care*, 27(5), hal. 1047–1053. doi: 10.2337/diacare.27.5.1047.

World Health Organisation (2006) *definition and diagnosis of diabetes mellitus and intermediate hyperglycemia Report of a WHO/IDF Consultation*.

World Health Organisation (2016) *GLOBAL REPORT ON DIABETES WHO Library Cataloguing-in-Publication Data Global report on diabetes*. Tersedia di: http://www.who.int/about/licensing/copyright_form/index.html (Diakses pada: 13 September 2019).

World Health Organisation (2018) ‘Global report on diabetes’, *Global Report on Diabetes*, hal. 88.