

## REFERENCES

- Agler, M. T., Wrenn, B. A., Zinder, S. H., & Angenent, L. T. (2011). Waste to bioproduct conversion with undefined mixed cultures: The carboxylate platform. *Trends in Biotechnology*, 29(2), 70–78. <https://doi.org/10.1016/j.tibtech.2010.11.006>
- Agnihotri, V. P. (1967). Role of Trace Elements in the Growth and Morphology of Five Ascosporic *Aspergillus* Species. *Canadian Journal of Botany*, 45(1), 73–79. <https://doi.org/10.1139/b67-005>
- Asadollahzadeh, M., Ghasemian, A., Saraeian, A., Resalati, H., & Taherzadeh, M. J. (2018). Production of Fungal Biomass Protein by Filamentous Fungi Cultivation on Liquid Waste Streams from Pulping Process. *BioResources*, 13(3), 5013–5031.
- Atasoy, M., Owusu-Agyeman, I., Plaza, E., & Cetecioglu, Z. (2018). Bio-based volatile fatty acid production and recovery from waste streams: Current status and future challenges. *Bioresource Technology*, 268(July), 773–786. <https://doi.org/10.1016/j.biortech.2018.07.042>
- Bhatia, S. K., & Yang, Y. H. (2017). Microbial production of volatile fatty acids: current status and future perspectives. *Reviews in Environmental Science and Biotechnology*, 16(2), 327–345. <https://doi.org/10.1007/s11157-017-9431-4>
- Chaves-Lopez, C., Nguyen, H. N., Oliveira, R. C., Nadres, E. T., Paparella, A., & Rodrigues, D. F. (2018). A morphological, enzymatic and metabolic approach to elucidate apoptotic-like cell death in fungi exposed to h- and  $\alpha$ -molybdenum trioxide nanoparticles. *Nanoscale*, 10(44), 20702–20716. <https://doi.org/10.1039/c8nr06470a>
- Deacon, J. W. (2006). Fungal Biology. In *Blackwell Publishing* (Vol. 4, Issue 9). Blackwell. <https://doi.org/10.1017/CBO9781107415324.004>

- Evren, M., Ozgun, H., Kaan, R., & Ozturk, I. (2011). Anaerobic Treatment of Industrial Effluents: An Overview of Applications. *Waste Water - Treatment and Reutilization*, June 2014, 2–28. <https://doi.org/10.5772/16032>
- Fei, Q., Chang, H. N., Shang, L., & Choi, J. D. R. (2011). Exploring low-cost carbon sources for microbial lipids production by fed-batch cultivation of *Cryptococcus albidus*. *Biotechnology and Bioprocess Engineering*, 16(3), 482–487. <https://doi.org/10.1007/s12257-010-0370-y>
- Fei, Q., Chang, H. N., Shang, L., Choi, J. dal rae, Kim, N. J., & Kang, J. W. (2011). The effect of volatile fatty acids as a sole carbon source on lipid accumulation by *Cryptococcus albidus* for biodiesel production. *Bioresource Technology*, 102(3), 2695–2701. <https://doi.org/10.1016/j.biortech.2010.10.141>
- Fei, Q., Fu, R., Shang, L., Brigham, C. J., & Chang, H. N. (2015). Lipid production by microalgae *Chlorella protothecoides* with volatile fatty acids (VFAs) as carbon sources in heterotrophic cultivation and its economic assessment. *Bioprocess and Biosystems Engineering*, 38(4), 691–700. <https://doi.org/10.1007/s00449-014-1308-0>
- Ferreira, J. A., Lennartsson, P. R., Niklasson, C., Lundin, M., Edebo, L., & Taherzadeha, M. J. (2012). Spent sulphite liquor for cultivation of an edible *Rhizopus* sp. *BioResources*, 7(1), 173–188. <https://doi.org/10.15376/biores.7.1.0173-0188>
- Ferreira, J. A., Lennartsson, P. R., & Taherzadeh, M. J. (2014). Production of ethanol and biomass from thin stillage using food-grade Zygomycetes and Ascomycetes filamentous fungi. *Energies*, 7(6), 3872–3885. <https://doi.org/10.3390/en7063872>
- Ferreira, J. A., Lennartsson, P. R., & Taherzadeh, M. J. (2015). Production of ethanol and biomass from thin stillage by *Neurospora intermedia*: A pilot study for process diversification. *Engineering in Life Sciences*, 15(8), 751–759.

<https://doi.org/10.1002/elsc.201400213>

Ferreira, J. A., Mahboubi, A., Lennartsson, P. R., & Taherzadeh, M. J. (2016). Waste biorefineries using filamentous ascomycetes fungi: Present status and future prospects. *Bioresource Technology*, *215*, 334–345.  
<https://doi.org/10.1016/j.biortech.2016.03.018>

Filho, P. F. S., Zamani, A., & Taherzadeh, M. J. (2017). Production of edible fungi from potato protein liquor (PPL) in airlift bioreactor. *Fermentation*, *3*(1).  
<https://doi.org/10.3390/fermentation3010012>

Fontanille, P., Kumar, V., Christophe, G., Nouaille, R., & Larroche, C. (2012). Bioconversion of volatile fatty acids into lipids by the oleaginous yeast *Yarrowia lipolytica*. *Bioresource Technology*, *114*, 443–449.  
<https://doi.org/10.1016/j.biortech.2012.02.091>

Foster, J. W. (1939). *THE BOTANICAL REVIEW THE HEAVY METAL NUTRITION OF FUNGI\** *New Jersey Agricultural Experiment Station*. V(4).  
<https://link.springer.com/content/pdf/10.1007/BF02878490.pdf>

Gerwien, F., Skrahina, V., Kasper, L., Hube, B., & Brunke, S. (2018). Metals in fungal virulence. *FEMS Microbiology Reviews*, *42*(1), 1–21.  
<https://doi.org/10.1093/femsre/fux050>

Gibbs, P. A., Seviour, R. J., & Schmid, F. (2000). Growth of filamentous fungi in submerged culture: Problems and possible solutions. *Critical Reviews in Biotechnology*, *20*(1), 17–48. <https://doi.org/10.1080/07388550091144177>

Griffin, D. H., Winkelmann, G., & Winge, D. R. (1995). Metal Ions in Fungi. In *Mycologia* (Vol. 87, Issue 3). <https://doi.org/10.2307/3760843>

Haenseler, C. M. (1921). The Effect of Salt Proportions and Concentration on the Growth of *Aspergillus niger*. *American Journal of Botany*, *8*(3), 147.

<https://doi.org/10.2307/2435113>

Hynes, M. J., Murray, S. L., Khew, G. S., & Davis, M. A. (2008). Genetic analysis of the role of peroxisomes in the utilization of acetate and fatty acids in *Aspergillus nidulans*. *Genetics*, *178*(3), 1355–1369.  
<https://doi.org/10.1534/genetics.107.085795>

Jacobsen, B. T., Wathne, B., & Lie, S. (1977). *CHELATORS AND METAL BUFFERING IN FERMENTATION*. *83*(I), 139–143.

Jin, B., Van Leeuwen, H. J., Patel, B., & Yu, Q. (1998). Utilisation of starch processing wastewater for production of microbial biomass protein and fungal  $\alpha$ -amylase by *Aspergillus oryzae*. *Bioresource Technology*, *66*(3), 201–206.  
[https://doi.org/10.1016/S0960-8524\(98\)00060-1](https://doi.org/10.1016/S0960-8524(98)00060-1)

Jin, Bo, Van Leeuwen, H. J., Patel, B., Doelle, H. W., & Yu, Q. (1999). Production of fungal protein and glucoamylase by *Rhizopus oligosporus* from starch processing wastewater. *Process Biochemistry*, *34*(1), 59–65. [https://doi.org/10.1016/S0032-9592\(98\)00069-7](https://doi.org/10.1016/S0032-9592(98)00069-7)

Jin, Bo, Van Leeuwen, J., & Patel, B. (1999). Mycelial morphology and fungal protein production from starch processing wastewater in submerged cultures of *Aspergillus oryzae*. *Process Biochemistry*, *34*(4), 335–340.  
[https://doi.org/10.1016/S0032-9592\(98\)00098-3](https://doi.org/10.1016/S0032-9592(98)00098-3)

Karimi, S., Soofiani, N. M., Lundh, T., Mahboubi, A., Kiessling, A., & Taherzadeh, M. J. (2019). Evaluation of filamentous fungal biomass cultivated on vinasse as an alternative nutrient source of fish feed: Protein, lipid, and mineral composition. *Fermentation*, *5*(4). <https://doi.org/10.3390/fermentation5040099>

Kavanagh, K. (2005). *Fungi: Biology and Applications*. John Wiley & Sons Ltd.

Kim, N.-J., Lim, S.-J., & Chang, H. N. (2018). Volatile Fatty Acid Platform: Concept

and Application. *Emerging Areas in Bioengineering*, *i*, 173–190.  
<https://doi.org/10.1002/9783527803293.ch10>

Kinderlerer, J. L. (1993). Fungal strategies for detoxification of medium chain fatty acids. *International Biodeterioration and Biodegradation*, *32*(1–3), 213–224.  
[https://doi.org/10.1016/0964-8305\(93\)90053-5](https://doi.org/10.1016/0964-8305(93)90053-5)

Kisworini, A. D. (2019). *Effect of Individual Volatile Fatty Acids (VFAs) on the Growth of Rhizopus microscopus var. oligosporus CBS 112586 in Batch Fermentation*.

Kleerebezem, R., Joosse, B., Rozendal, R., & Van Loosdrecht, M. C. M. (2015). Anaerobic digestion without biogas? *Reviews in Environmental Science and Biotechnology*, *14*(4), 787–801. <https://doi.org/10.1007/s11157-015-9374-6>

Lubert Stryer Jeremy M. Berg John L. Tymoczko with, & Gregory J. Gatto, J. W. (2012). *Biochemistry* (Vol. 33). <https://doi.org/10.1073/pnas.0703993104>

Moore, D., Robson, G. D., & Trinci, A. P. J. (2011). 21st Century Guidebook to Fungi. In *Cambridge University Press*. Cambridge University Press, New York.

Nair, R. B., & Taherzadeh, M. J. (2016). Valorization of sugar-to-ethanol process waste vinasse: A novel biorefinery approach using edible ascomycetes filamentous fungi. *Bioresource Technology*, *221*, 469–476.  
<https://doi.org/10.1016/j.biortech.2016.09.074>

Nitayavardhana, S., Issarapayup, K., Pavasant, P., & Khanal, S. K. (2013). Production of protein-rich fungal biomass in an airlift bioreactor using vinasse as substrate. *Bioresource Technology*, *133*, 301–306.  
<https://doi.org/10.1016/j.biortech.2013.01.073>

Nitayavardhana, S., & Khanal, S. K. (2010). Innovative biorefinery concept for sugar-based ethanol industries: Production of protein-rich fungal biomass on vinasse as an aquaculture feed ingredient. *Bioresource Technology*, *101*(23), 9078–9085.

<https://doi.org/10.1016/j.biortech.2010.07.048>

Nout, M. J. R., & Aidoo, K. E. (2002). Asian Fungal Fermented Food. *Industrial Applications*, 23–47. [https://doi.org/10.1007/978-3-662-10378-4\\_2](https://doi.org/10.1007/978-3-662-10378-4_2)

Papargyropoulou, E., Lozano, R., K. Steinberger, J., Wright, N., & Ujang, Z. Bin. (2014). The food waste hierarchy as a framework for the management of food surplus and food waste. *Journal of Cleaner Production*, 76, 106–115. <https://doi.org/10.1016/j.jclepro.2014.04.020>

Robbins, W. J. ., & Kavanagh, V. (1942). Vitamin Deficiencies of the Filamentous Fungi. *New York Botanical Garden Press*, 8(7), 411–471. <http://www.jstor.org/stable/4353273>

Ruhil, S., Kumar, V., Balhara, M., Malik, M., Dhankhar, S., Kumar, M., & Kumar Chhillar, A. (2014). In vitro evaluation of combination of polyenes with EDTA against *Aspergillus* spp. by different methods (FICI and CI Model). *Journal of Applied Microbiology*, 117(3), 643–653. <https://doi.org/10.1111/jam.12576>

Safaei, Z., Karimi, K., & Zamani, A. (2016). Impact of phosphate, potassium, yeast extract, and trace metals on chitosan and metabolite production by *Mucor indicus*. *International Journal of Molecular Sciences*, 17(9). <https://doi.org/10.3390/ijms17091429>

Souza Filho, P. F. (2018). *Fungi-based biorefinery model for food industry waste Progress toward a circular economy*. <http://urn.kb.se/resolve?urn=urn:nbn:se:hb:diva-14888>

Souza Filho, P. F., Nair, R. B., Andersson, D., Lennartsson, P. R., & Taherzadeh, M. J. (2018). Vegan-mycoprotein concentrate from pea-processing industry byproduct using edible filamentous fungi. In *Fungal Biology and Biotechnology* (Vol. 5, Issue 1). <https://doi.org/10.1186/s40694-018-0050-9>

- Souza Filho, P. F., Zamani, A., & Taherzadeh, M. J. (2019). Edible Protein Production by Filamentous Fungi using Starch Plant Wastewater. *Waste and Biomass Valorization*, 10(9), 2487–2496. <https://doi.org/10.1007/s12649-018-0265-2>
- Steinberg, R. A. (1919). a Study of Some Factors in the Chemical Stimulation of the Growth of *Aspergillus Niger*. *American Journal of Botany*, 6(8), 330–356. <https://doi.org/10.1002/j.1537-2197.1919.tb05547.x>
- Stratford, M., Nebe-von-Caron, G., Steels, H., Novodvorska, M., Ueckert, J., & Archer, D. B. (2013). Weak-acid preservatives: PH and proton movements in the yeast *Saccharomyces cerevisiae*. *International Journal of Food Microbiology*, 161(3), 164–171. <https://doi.org/10.1016/j.ijfoodmicro.2012.12.013>
- Stratford, M., Plumridge, A., Nebe-von-Caron, G., & Archer, D. B. (2009). Inhibition of spoilage mould conidia by acetic acid and sorbic acid involves different modes of action, requiring modification of the classical weak-acid theory. *International Journal of Food Microbiology*, 136(1), 37–43. <https://doi.org/10.1016/j.ijfoodmicro.2009.09.025>
- Strazzera, G., Battista, F., Garcia, N. H., Frison, N., & Bolzonella, D. (2018). Volatile fatty acids production from food wastes for biorefinery platforms: A review. *Journal of Environmental Management*, 226(August), 278–288. <https://doi.org/10.1016/j.jenvman.2018.08.039>
- Sues, A., Millati, R., Edebo, L., & Taherzadeh, M. J. (2005). Ethanol production from hexoses, pentoses, and dilute-acid hydrolyzate by *Mucor indicus*. *FEMS Yeast Research*, 5(6–7), 669–676. <https://doi.org/10.1016/j.femsyr.2004.10.013>
- Taherzadeh, M. J., Fox, M., Hjorth, H., & Edebo, L. (2003). Production of mycelium biomass and ethanol from paper pulp sulfite liquor by *Rhizopus oryzae*. *Bioresource Technology*, 88(3), 167–177. [https://doi.org/10.1016/S0960-8524\(03\)00010-5](https://doi.org/10.1016/S0960-8524(03)00010-5)

- Wainaina, S., Kisworini, A. D., Fanani, M., Wikandari, R., Millati, R., Niklasson, C., & Taherzadeh, M. J. (2020). Utilization of food waste-derived volatile fatty acids for production of edible *Rhizopus oligosporus* fungal biomass. *Bioresource Technology*, *310*, 123444. <https://doi.org/10.1016/j.biortech.2020.123444>
- Wainaina, S., Lukitawesa, Kumar Awasthi, M., & Taherzadeh, M. J. (2019). Bioengineering of anaerobic digestion for volatile fatty acids, hydrogen or methane production: A critical review. *Bioengineered*, *10*(1), 437–458. <https://doi.org/10.1080/21655979.2019.1673937>
- Wainaina, S., Parchami, M., Mahboubi, A., Horváth, I. S., & Taherzadeh, M. J. (2019). Food waste-derived volatile fatty acids platform using an immersed membrane bioreactor. *Bioresource Technology*, *274*, 329–334. <https://doi.org/10.1016/j.biortech.2018.11.104>
- Xu, F., Li, Y., Ge, X., Yang, L., & Li, Y. (2018). Anaerobic digestion of food waste – Challenges and opportunities. *Bioresource Technology*, *247*(September), 1047–1058. <https://doi.org/10.1016/j.biortech.2017.09.020>
- Yun, J. H., Sawant, S. S., & Kim, B. S. (2013). Production of polyhydroxyalkanoates by *Ralstonia eutropha* from volatile fatty acids. *Korean Journal of Chemical Engineering*, *30*(12), 2223–2227. <https://doi.org/10.1007/s11814-013-0190-9>
- Zhang, J., & Zhang, J. (2016). The filamentous fungal pellet and forces driving its formation. *Critical Reviews in Biotechnology*, *36*(6), 1066–1077. <https://doi.org/10.3109/07388551.2015.1084262>