

DAFTAR PUSTAKA

- Aissa, A.F., Bianchi, M.L.P., Ribeiro, J.C., Hernandez, L.C.H., de Faria, A.F., Mercadante, A.Z., Antunes, L.M.G., 2012, Comparative study of β -carotene and microencapsulated β -carotene: Evaluation of their genotoxic and antigenotoxic effects. *Food Chem. Toxicol.*, 50, 1418–1424.
- Alcázar-Alay, S.C. and Meireles, M.A.A., 2015, Physicochemical properties, modifications and applications of starches from different botanical sources. *Food Sci. Technol.*, 35, 215–236.
- Anonim, 2002, *Starch* (CAS reg no: 9005-25-8, 9005-84-9), Health-based Reassessment of Administrative Occupational Exposure Limits Committee on Updating of Occupational Exposure Limits, A committee of the Health Council of the Netherlands.
- Anonim, 2011, *Provisional peer-reviewed toxicity values for inorganic phosphates (orthophosphoric acid and inorganic phosphate compounds, including ortho- and condensed phosphates) (various CASRNs included in the text)*, Superfund Health Risk Technical Support Center, National Center for Environmental Assessment, Office of Research and Development, U.S. Environmental Protection Agency.
- Anonim, 2017, *NTP Technical report on the toxicity study of (CASRN 9012-76-4) chitosan administered in feed to Sprague dawley [CRL:CD(SD)] rats*, National Toxicology Program, Public Health Service, U.S. Department of Health and Human Services.
- Anonim, 2020a, Beta-Carotene. *Natl. Cent. Biotechnol. Information. PubChem Database.* *beta-Carotene*, CID=5280489, <https://pubchem.ncbi.nlm.nih.gov/compound/beta-Carotene> (accessed Mar. 5, 2020).
- Anonim, 2020b, Corn starch (Compound summary). <https://pubchem.ncbi.nlm.nih.gov/compound/24836924> (accessed Juli. 22, 2020).
- Ariviani, S., Anggrahini, S., Naruki, S. and Raharjo, S., 2015, Characterization and chemical stability evaluation of β -carotene microemulsions prepared by spontaneous emulsification method using VCO and palm oil as oil phase. *Int. Food Res. J.*, 22, 2432–2439.
- Arunkumar, R., Prashanth, K.V.H., and Baskaran, V., 2013, Promising interaction between nanoencapsulated lutein with low molecular weight chitosan: Characterization and bioavailability of lutein in vitro and in vivo. *Food Chem.*, 141, 327–337.

- El Asbahani, A., Miladi, K., Badri, W., Sala, M., Addi, E.H.A., Casabianca, H., et al., 2015, Essential oils : From extraction to encapsulation. *Int. J. Pharm.*, 483, 220–243.
- Bajer, D. and Kaczmarek, H., 2010, Study of the Influence OV UV Radiation on Biodegradable Blends Based on Chitosan and Starch. *Prog. Chem. Appl. Chitin Its Deriv.*, 15, 17–24.
- Belščak-Cvitanovic, A., Bušić, A., Barišić, L., Vrsaljko, D., Karlović, S., Špoljarić, I., et al., 2016, Emulsion templated microencapsulation of dandelion (*Taraxacum officinale* L.) polyphenols and β -carotene by ionotropic gelation of alginate and pectin. *Food Hydrocoll.*, 57, 139–152.
- Blanchette, F., Messio, L., and Bush, J.W.M., 2009, The influence of surface tension gradients on drop coalescence. *Phys. Fluids*, 21, 1–10.
- Chang, Y., Yan, X., Wang, Q., Ren, L., Tong, J., and Zhou, J., 2017, High efficiency and low cost preparation of size controlled starch nanoparticles through ultrasonic treatment and precipitation. *Food Chem.*, 227, 369–375.
- Chavarri, M., Maranon, I., and Carmen, M., 2012, Encapsulation Technology to Protect Probiotic Bacteria. In, Rigobelo, E.C. (ed), *Probiotics.*, Intech, pp.501–540.
- Chen, J., Shi, J., Macnaughton, L., Kakuda, Y., Xue, S.J., Ma, Y., Zhang, M., Jiang, Y., 2009, Lycopene, β -carotene, vitamin E, and vitamin C on the free radical 2,2-diphenyl-1-picrylhydrazyl (DPPH). *J. Food Biochem.*, 33, 232–245.
- Chen, L., McClements, D.J., Zhang, H., Zhang, Z., Jin, Z., and Tian, Y., 2019, Impact of amylose content on structural changes and oil absorption of fried maize starches. *Food Chem.*, 287, 28–37.
- Chu, B.S., Ichikawa, S., Kanafusa, S., and Nakajima, M., 2007, Preparation and characterization of β -carotene nanodispersions prepared by solvent displacement technique. *J. Agric. Food Chem.*, 55, 6754–6760.
- Coates, J., 2000, Interpretation of Infrared Spectra, A Practical Approach. In, Meyers, R.A. (ed), *Encyclopedia of Analytical Chemistry*. John Wiley & Sons Ltd, Chichester, pp. 10815–10837.
- Coronel-Aguilera, C.P. and Martín-González, 2015, Encapsulation of spray dried β -carotene emulsion by fluidized bed coating technology. *LWT - Food Sci. Technol.*, 62, 187–193.
- Dash, S., Murthy, P.N., Nath, L., and Chowdhury, P., 2010, Kinetic modeling on drug release from controlled drug delivery systems. *Acta Pol. Pharm.*, 67,

217–23.

Donhowe, E.G., Flores, F.P., Kerr, W.L., Wicker, L., and Kong, F., 2014, LWT - Food Science and Technology Characterization and in vitro bioavailability of β -carotene : Effects of microencapsulation method and food matrix. *LWT - Food Sci. Technol.*, 57, 42–48.

Donhowe, E.G. and Kong, F., 2014, Beta-carotene : Digestion, Microencapsulation, and In Vitro Bioavailability. *Food Bioprocess Technol*, 7, 338–354.

El-Naggar, M.E., El-Rafie, M.H., El-sheikh, M.A., El-Feky, G.S., and Hebeish, A., 2015, Synthesis, characterization, release kinetics and toxicity profile of drug-loaded starch nanoparticles. *Int. J. Biol. Macromol.*, 81, 718–729.

Fathi, M., Martín, Á., and McClements, D.J., 2014, Fathi, M., Martín, Á., and McClements, D.J., 2014, Nanoencapsulation of food ingredients using carbohydrate based delivery systems. *Trends Food Sci. Technol.* ,39, 18–39. Nanoencapsulation of food ingredients using carbohydrate based delivery systems. *Trends Food Sci. Technol.*, 39, 18–39.

Ferranti, V., Marchais, H., Chabenat, C., Orecchioni, A.M., and Lafont, O., 1999, Primidone-loaded poly- ϵ -caprolactone nanocapsules: Incorporation efficiency and in vitro release profiles. *Int. J. Pharm.* ,193, 107–111.

Fessi, H., Puisieux, F., Devissaguet, J.P., Ammoury, N., and Benita, S., 1989, Nanocapsule formation by interfacial polymer deposition following solvent displacement. *Int. J. Pharm.*, 55, 1–4.

Fouladi, E. and Nafchi, A.M., 2014, Effects of acid-hydrolysis and hydroxypropylation on functional properties of sago starch. *Int. J. Biol. Macromol.*, 68, 251–257.

Gul, K., Tak, A., Singh, A.K., Singh, P., Yousuf, B., and Wani, A.A., 2015, Chemistry, encapsulation, and health benefits of β -carotene - A review. *Cogent Food Agric.*, 1, 1–12.

Guti, F.J., Silvia, M., Casas-sanz, E., Garc, A., Gat, P., Gonz, C., et al., 2013, Methods for the nanoencapsulation of β -carotene in the food sector. *Trends Food Sci. Technol.*, 32, 73–83.

Hart, A.Y., 2012, Effects of resistant starch and soluble fiber on the bioaccessibility of dietary carotenoids from spinach and carrot using simulated in vitro digestion, *Thesis*, Graduate Program in Food Science and Technology, Ohio State University.

Hasani, S., Ojagh, S.M., and Ghorbani, M., 2018, Nanoencapsulation of lemon

essential oil in Chitosan-Hicap system. Part 1: Study on its physical and structural characteristics. *Int. J. Biol. Macromol.*, 115, 143–151.

Higuchi, T., 1963, Mechanism of sustained-action medication. Theoretical analysis of rate of release of solid drugs dispersed in solid matrices. *J. Pharm. Sci.*, 52, 1145–1149.

Honary, S. and Zahir, F., 2013, Effect of Zeta Potential on the Properties of Nano-Drug Delivery Systems -A Review (Part 2). *Trop J Pharm Res.*, 12, 2, 265–273

Hoover, R., 2007, Acid-Treated Starches. *Food Rev. Int.*, 16, 37–41.

Hou, Z., Gao, Y., Yuan, F., Liu, Y., Li, C., and Xu, D., 2010, Investigation into the physicochemical stability and rheological properties of β -carotene emulsion stabilized by soybean soluble polysaccharides and chitosan. *J. Agric. Food Chem.*, 58, 8604–8611.

Hsin, W.L., Sheng, Y.J., Lin, S.Y., and Tsao, H.K., 2004, Surface tension increment due to solute addition. *Phys. Rev. E*, 69, 1–8.

Joye, I.J., Davidov-Pardo, G., and McClements, D.J., 2014, Nanotechnology for increased micronutrient bioavailability. *Trends Food Sci. Technol.*, 40, 168–182.

Kim, J.Y. and Huber, K.C., 2016, Preparation and characterization of corn starch- β -carotene composites. *Carbohydr. Polym.*, 136, 394–401.

Kong, L., Bhosale, R., and Ziegler, G., 2018, Encapsulation and stabilization of β -carotene by amylose inclusion complexes. *Food. Res. Int.*, 105, 446–452.

Korsmeyer, R.W., Gurny, R., Doelker, E., Buri, P., and Peppas, N.A., 1983, Mechanisms of solute release from porous hydrophilic polymers. *Int. J. Pharm.*, 15, 25–35.

Lawal, O.S. and Adebawale, K.O., 2005, Physicochemical characteristics and thermal properties of chemically modified jack bean (*Canavalia ensiformis*) starch. *Carbohydr. Polym.*, 60, 331–341.

Lin, Q., Liang, R., Williams, P.A., and Zhong, F., 2017, Factors affecting the bioaccessibility of β -carotene in lipid-based microcapsules: digestive conditions, the composition, structure and physical state of microcapsules. *Food Hydrocoll.*, xxx, 1–7.

Liu, F., Qin, B., He, L., and Song, R., 2009, Novel starch/chitosan blending membrane: Antibacterial, permeable and mechanical properties.

Carbohydr. Polym., 78, 146–150.

- Loksuwan, J., 2007, Characteristics of microencapsulated β -carotene formed by spray drying with modified tapioca starch, native tapioca starch and maltodextrin. *Food Hydrocoll.*, 21, 928–935.
- Loutfy, S.A., El-Din, H.M.A., Elberry, M.H., Allam, N.G., Hasanin, M.T.M., and Abdellah, A.M., 2016, Synthesis, characterization and cytotoxic evaluation of chitosan nanoparticles: In vitro liver cancer model. *Adv. Nat. Sci. Nanosci. Nanotechnol.*, 7, 035008- 035017.
- Lu, D.R., Xiao, C.M., and Xu, S.J., 2009, Starch-based completely biodegradable polymer materials. *Express Polym. Lett.*, 3, 366–375.
- Mao, Y., Dubot, M., Xiao, H., and McClements, D.J., 2013, Interfacial Engineering Using Mixed Protein Systems: Emulsion- Based Delivery Systems for Encapsulation and Stabilization of β -Carotene. *J. Agric. Food Chem. Mix.*, 61, 5163–5169.
- Ma, Y., Hou, C.J., Fa, H.B., Huo, D.Q., Yang, M., 2016, Synthesis and antioxidant property of hydroxycinnamoyl maltodextrin derivatives. *Ifst*, 51, 11, 2450-2459.
- Mishra, K., Ojha, H., and Chaudhury, N.K., 2012, Estimation of antiradical properties of antioxidants using DPPH - assay: A critical review and results. *Food Chem.*, 130, 1036–1043.
- Moon, J. and Shibamoto, T., 2009, Antioxidant Assays for Plant and Food Components Antioxidant Assays for Plant and Food Components. *J. Agric. Food Chem.*, 57, 1655–1666.
- Mun, S., Park, S., Kim, Y., and Julian, D., 2016, Influence of methylcellulose on attributes of β -carotene fortified starch-based filled hydrogels: Optical, rheological, structural, digestibility, and bioaccessibility properties. *Food Res. Int.*, 87, 18–24.
- Muxika, A., Etxabide, A., Uranga, J., Guerrero, P., and de la Caba, K., 2017, Chitosan as a bioactive polymer: Processing, properties and applications. *Int. J. Biol. Macromol.*, 105, 1358–1368.
- Namitha, K.K. and Negi, P.S., 2010, Chemistry and Biotechnology of Carotenoids. *Crit. Rev. Food Sci. Nutr.*, 50, 728–760.
- Nilsen-Nygaard, J., Strand, S.P., Vårum, K.M., Draget, K.I., and Nordgård, C.T., 2015, Chitosan: Gels and interfacial properties. *Polymers (Basel)*, 7, 552–579.

- Nimse, S.B. and Pal, D., 2015, Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Adv.*, 5, 27986–28006.
- Obayemi, J.D., Danyuo, Y., Dozie-Nwachukwu, S., Odusanya, O.S., Anuku, N., Malatesta, K., et al., 2016, PLGA-based microparticles loaded with bacterial-synthesized prodigiosin for anticancer drug release: Effects of particle size on drug release kinetics and cell viability. *Mater. Sci. Eng. C*, 66, 51–65.
- Ozcelik, B., Karadag, A., and Ersen, S., 2009, Bioencapsulation of Beta-Carotene in Three Different Methods. In, *VIIth International Conference on Bioencapsulation*. Groningen, Netherlands, pp. 2–3.
- Parize, A.L., Stulzer, H.K., Laranjeira, M.C.M., Da Costa Brighente, I.M., and De Souza, T.C.R., 2012, Evaluation of chitosan microparticles containing curcumin and crosslinked with sodium tripolyphosphate produced by spray drying. *Quim. Nova*, 35, 1127–1132.
- Peinado, I., Mason, M., Romano, A., Biasioli, F., Scampicchio, M., 2016, Stability of β -carotene in polyethylene oxide electrospun nanofibers. *Appl. Surf. Sci.*, 370, 111–116.
- Peng, H., Xiong, H., Li, J., Xie, M., Liu, Y., Bai, C., and Chen, L., 2010, Vanillin cross-linked chitosan microspheres for controlled release of resveratrol. *Food Chem.*, 121, 23–28.
- Perevyazko, I.Y., Vollrath, A., Pietsch, C., Schubert, S., Pavlov, G.M., and Schubert, U.S., 2012, Nanoprecipitation of poly(methyl methacrylate)-based nanoparticles: Effect of the molar mass and polymer behavior. *J. Polym. Sci. Part A Polym. Chem.*, 50, 2906–2913.
- Perez, J.J. and Francois, N.J., 2016, Chitosan-starch beads prepared by ionotropic gelation as potential matrices for controlled release of fertilizers. *Carbohydr. Polym.*, 148, 134–142.
- Priamo, W.L., De Cezaro, A.M., Ferreira, S.R.S., and Oliveira, J.V., 2010, Precipitation and encapsulation of β -carotene in PHBV using carbon dioxide as anti-solvent. *J. Supercrit. Fluids*, 54, 103–109.
- Putseys, J.A., Lamberts, L., and Delcour, J.A., 2010, Amylose-inclusion complexes: Formation, identity and physico-chemical properties. *J. Cereal Sci.*, 51, 238–247.
- Pyrzyska, K. and Pękal, A., 2013, Application of free radical diphenylpicrylhydrazyl (DPPH) to estimate the antioxidant capacity of food samples. *Anal. Methods*, 5, 4288–4295.

- Raguzzoni, J.C., Delgadillo, I., and Lopes Da Silva, J.A., 2016, Influence of a cationic polysaccharide on starch functionality. *Carbohydr. Polym.*, 150, 369–377.
- Rampino, A., Borgogna, M., Bellich, B., Blasi, P., Virgilio, F., and Cesàro, A., 2016, Chitosan-pectin hybrid nanoparticles prepared by coating and blending techniques. *Eur. J. Pharm. Sci.*, 84, 37–45.
- Ray, S., Raychaudhuri, U., and Chakraborty, R., 2016, Food Bioscience An overview of encapsulation of active compounds used in food products by drying technology. *Food Biosci.*, 13, 76–83.
- Rivas, C.J.M., Tarhini, M., Badri, W., Miladi, K., Greige-Gerges, H., Nazari, Q.A., et al., 2017, Nanoprecipitation process: From encapsulation to drug delivery. *Int. J. Pharm.*, 532, 66–81.
- Roohinejad, S., Middendorf, D., Burritt, D.J., Bindrich, U., Everett, D.W., and Oey, I., 2014, Capacity of natural β -carotene loaded microemulsion to protect Caco-2 cells from oxidative damage caused by exposure to H₂O₂. *Frin*, 66, 469–477.
- Ruiz, R.B. and Hernández, P.S., 2016, Cancer chemoprevention by dietary phytochemicals: epidemiological evidence. *Maturitas*, 94, 13–19.
- Rutz, J.K., Borges, C.D., Zambiasi, R.C., Cleonice, G., and Médelin, M., 2016, Elaboration of microparticles of carotenoids from natural and synthetic sources for applications in food. *Food Chem.*, 202, 324–333.
- Saini, R.K., Nile, S.H., and Park, S.W., 2015, Carotenoids from fruits and vegetables: Chemistry, analysis, occurrence, bioavailability and biological activities. *Food Res. Int.*, 76, 735–750.
- Science, O.F., 2016, *Alcohol Hand Sanitizer With Improved Dermal Compatibility And Feel*, US Patent, Pub. No .: US 2016/0271610 A1.
- Senanayake, S., Gunaratne, A., Ranaweera, K., and Bamunuarachchi, A., 2013, Effect of heat moisture treatment conditions on swelling power and water soluble index of different cultivars of sweet potato (Ipomea batatas (L). Lam) Starch . *ISRN Agron.*, 2013, 1–4.
- Singh, B.B., Shakil, N.A., Kumar, J., Walia, S., Kar, A., 2015, Development of slow release formulations of β -carotene employing amphiphilic polymers and their release kinetics study in water and different pH conditions. *J. Food. Sci. Technol.*, 52, 12, 8068–8076.
- Singh, H., Sodhi, S.N., and Singh, N., 2009, Structure and functional properties of

- acid thinned sorghum starch. *Int J Food Prop.*, 12, 4, 713-725.
- Singh, S.K., Singh, N., and Lim, S.T., 2007, A comparison of native and acid thinned normal and waxy corn starches: Physicochemical, thermal, morphological and pasting properties. *LWT - Food Sci. Technol.*, 40, 1527–1536.
- Soukoulis, C., Tsevdou, M., Andre, C.M., Cambier, S., Yonekura, L., Taoukis, P.S., and Hoffmann, L., 2017, Modulation of chemical stability and in vitro bioaccessibility of beta-carotene loaded in kappa-carrageenan oil-in-gel emulsions. *Food Chem.*, 220, 208–218.
- Spada, J.C., Marczak, D.L.F., Tessaro, I.C., and Noreña, C.P.Z., 2012a, Microencapsulation of β -carotene using native pinhão starch, modified pinhão starch and gelatin by freeze-drying. *Int. J. Food Sci. Technol.*, 47, 186–194.
- Spada, J.C., Noreña, C.P.Z., Marczak, L.D.F., and Tessaro, I.C., 2012b, Study on the stability of β -carotene microencapsulated with pinhão (*Araucaria angustifolia* seeds) starch. *Carbohydr. Polym.*, 89, 1166–1173.
- Subramanian, S.B., Francis, A.P., and Devasena, T., 2014, Chitosan-starch nanocomposite particles as a drug carrier for the delivery of bis-desmethoxy curcumin analog. *Carbohydr. Polym.*, 114, 170–178.
- Sun, Q., Zhu, X., Si, F., and Xiong, L., 2015, Effect of acid hydrolysis combined with heat moisture treatment on structure and physicochemical properties of corn starch. *J. Food Sci. Technol.*, 52, 375–382.
- Talón, E., Trifkovic, K.T., Nedovic, V.A., Bugarski, B.M., Vargas, M., Chiralt, A., and González-Martínez, C., 2017, Antioxidant edible films based on chitosan and starch containing polyphenols from thyme extracts. *Carbohydr. Polym.*, 157, 1153–1161.
- Tapola, N.S., Lyyra, M.L., Kolehmainen, R.M., Sarkkinen, E.S., and Schauss, A.G., 2008, Safety aspects and cholesterol-lowering efficacy of chitosan tablets, *J. Am. Coll. Nutr.*, 27, 1, 22–30.
- Thamaket, P. and Raviyan, P., 2015, Preparation and physical properties of carotenoids encapsulated in chitosan cross-linked tripolyphosphate nanoparticles. *Food Appl. Biosci. J.*, 3, 69–84.
- Ujhelyi, Z., Fenyvesi, F., Váradi, J., Fehér, P., Kiss, T., Veszélka, S., et al., 2012, Evaluation of cytotoxicity of surfactants used in self-micro emulsifying drug delivery systems and their effects on paracellular transport in Caco-2 cell monolayer. *Eur. J. Pharm. Sci.*, 47, 564–573.

- Ulbrich, M., Beresnewa-Seekamp, T., Walther, W., and Flöter, E., 2016, Acid-thinned corn starch-impact of modification parameters on molecular characteristics and functional properties. *Starch/Stärke*, 68, 399–409.
- de Vos, P., Faas, M.M., Spasojevic, M., and Sikkema, J., 2010, Encapsulation for preservation of functionality and targeted delivery of bioactive food components. *Int. Dairy J.*, 20, 292–302.
- Wang, Q., Zhang, N., Hu, X., Yang, J., and Du, Y., 2007, Chitosan/starch fibers and their properties for drug controlled release. *Eur. J. Pharm. Biopharm.*, 66, 398–404.
- Wang, S and Copeland, L., 2012, New insights into loss of swelling power and pasting profiles of acid hydrolyzed starch granules. *Starch/Stärke*, 00, 1-7.
- Wang, Y.J., Truong, V.D., and Wang, L., 2003, Structures and rheological properties of corn starch as affected by acid hydrolysis. *Carbohydr. Polym.*, 52, 327–333.
- Worthen, A.J., Irving, K.S., and Lapitsky, Y., 2019, Supramolecular Strategy Effects on Chitosan Bead Stability in Acidic Media: A Comparative Study. *Gels.*, 5, 11, 1-12.
- Yahia, Y., Villén, F.G., Djelad, A., Belaroui, L.S., Espejo, R.S., Sassi, M., Galindo, A.L., Viseras, C., 2019, Crosslinked palygorskite-chitosan beads as diclofenac carriers. *Appl. Clay Sci.*, 180, 105169, 1-10.
- Yang, J., Han, S., Zheng, H., Dong, H., and Liu, J., 2015, Preparation and application of micro/nanoparticles based on natural polysaccharides. *Carbohydr. Polym.*, 123, 53–66.
- Yen, M.T., Yang, J.H., and Mau, J.L., 2008, Antioxidant properties of chitosan from crab shells. *Carbohydr. Polym.*, 74, 840–844.
- Yi, J., Lam, T.I., Yokoyama, W., Cheng, L.W., and Zhong, F., 2015, Beta-carotene encapsulated in food protein nanoparticles reduces peroxy radical oxidation in Caco-2 cells. *Food Hydrocoll.*, 43, 31–40.
- Yousif, E.I., Gadallah, M.G.E., and Sorour, A.M., 2012, Physico-chemical and rheological properties of modified corn starches and its effect on noodle quality. *Ann. Agric. Sci.*, 57, 19–27.
- Yuan, Y., Gao, Y., Zhao, J., and Mao, L., 2008, Characterization and stability evaluation of β -carotene nanoemulsions prepared by high pressure homogenization under various emulsifying conditions. *Food Res. Int.*, 41, 61–68.

Zhang, Z., Zhang, R., Julian, D., and McClements, D.J., 2016, Encapsulation of β -carotene in alginate-based hydrogel beads: Impact on physicochemical stability and bioaccessibility. *Food Hydrocoll.*, 61, 1–10.

Zhu, F., 2017, Encapsulation and delivery of food ingredients using starch based systems. *Food Chem.*, 229, 542–552.