

DAFTAR PUSTAKA

- Aaberg, M. L. *et al.* (2008) 'Gender differences in the onset of diabetic neuropathy', 22, pp. 83–87. doi: 10.1016/j.jdiacomp.2007.06.009.
- Abbott, C. A. *et al.* (2011) 'Prevalence and characteristics of painful diabetic neuropathy in a large community-based diabetic population in the U.K.', *Diabetes Care*. doi: 10.2337/dc11-1108.
- Allredge, B. K. *et al.* (2013) *Koda-Kimble and Young's applied therapeutics: The clinical use of drugs, Koda-Kimble and Young's Applied Therapeutics: The Clinical Use of Drugs*.
- Alvarez, M. *et al.* (2019) 'Vitamin B12 deficiency and diabetic neuropathy in patients taking metformin: A cross-sectional study', *Endocrine Connections*, 8(10), pp. 1324–1329. doi: 10.1530/EC-19-0382.
- American Diabetes Association. (2019) 'Standars of Medical Care in Diabetes 2019.' *Diabetes Care*
- Andres, E. and Serraj (2012) 'Optimal management of pernicious anemia', *Journal of Blood Medicine*, p. 97. doi: 10.2147/jbm.s25620.
- Aroda, V. R. *et al.* (2016) 'Long-term metformin use and vitamin B12 deficiency in the diabetes prevention program outcomes study', *Journal of Clinical Endocrinology and Metabolism*, 101(4), pp. 1754–1761. doi: 10.1210/jc.2015-3754.
- Asri DN. (2006) 'Kualitas Hidup Penderita Diabetes melitus Tipe 2 Ditinjau Dari Efikasi Diri, Persepsi, Dukungan Sosial Dan Strategi mengatasi masalah Aktif.' *Thesis Universitas Gajah Mada Yogyakarta*.
- Bc, C. *et al.* (2012) 'Enhanced glucose control for preventing and treating diabetic neuropathy (Review)', (6).
- Berhanu, P., Perez, A. and Yu, S. (2007) 'Effect of pioglitazone in combination with insulin therapy on glycaemic control , insulin dose requirement and lipid profile in patients with type 2 diabetes previously poorly controlled with combination therapy', pp. 512–520. doi: 10.1111/j.1463-1326.2006.00633.x.
- Booya, F. *et al.* (2005) 'Potential risk factors for diabetic neuropathy: A case control study', *BMC Neurology*. doi: 10.1186/1471-2377-5-24.
- Bril, V. (2004) 'Aldose Reductase Inhibition by AS-3201 in Sural Nerve From

Patients With', *Diabetes care*, 27(10).

Bril, V. *et al.* (2009) 'Reliability and validity of the modified Toronto Clinical Neuropathy Score in diabetic sensorimotor polyneuropathy', *Diabetic Medicine*, 26(3), pp. 240–246. doi: 10.1111/j.1464-5491.2009.02667.x.

Bril, V. and Perkins, B. A. (2002) 'Validation of the Toronto clinical scoring system for diabetic polyneuropathy', *Diabetes Care*, 25(11), pp. 2048–2052. doi: 10.2337/diacare.25.11.2048.

Brunton, Laurence; Chabner, Bruce; Knollman, B. (2013) *Goodman and Gilman's The Pharmacological Basis of Therapeutics, Twelfth Edition, Journal of Chemical Information and Modeling*. doi: 10.1017/CBO9781107415324.004.

Brunton, L. (2006) 'Goodman & Gilman' s The Pharmacological Basis of Therapeutics', *Goodman & Gilman' s The Pharmacological Basis of Therapeutics*. doi: 10.1017/CBO9781107415324.004.

Chaturvedi, N. *et al.* (2002) 'Risk of diabetes-related amputation in South Asians vs. Europeans in the UK', *Diabetic Medicine*. doi: 10.1046/j.1464-5491.2002.00583.x.

Chisholm-Burns. M., Wells, B., Schwinghammer, T., Malone, P.M., Kolesar, J.M., Dipiro, J.T., (2013) 'Pharmacotherapy Principles and Practice. 3rd edition.' *McGraw-Hill Medical Publishing Division*

Cipolle, R. ., Strand, L. M. & and Morley, P. C. (1998) 'Cipolle, R.J, Strand, L.M. & Morley, P.C., 1998, Pharmaceutical Care Practice', *Cipolle, R.J, Strand, L.M. & Morley, P.C., 1998, Pharmaceutical Care Practice* , hal : 75, 82 - 83, 96 - 101, 116, *Mc Graw Hill Company, New York*. doi: 10.1155/2012/796590.

Cipolle, R. J., Strand, L. M. and Morley, P. C. (2012) *Pharmaceutical Care Practice* :, *Pharmaceutical Care Practice: The Patient-Centered Approach to Medication Management Services. McGraw-Hill Medical Publishing Division*.

Corbett, C. F. (2005) 'Practical management of patients with painful diabetic neuropathy', *Diabetes Educator*. doi: 10.1177/0145721705278800.

Dahlan M. (2013) 'Beser Sampel Dan Cara Pengambilan Sampel Dalam Penelitian Kedokteran dan Kesehatan, 3rd ed.' *Salemba Medika, Jakarta*.

Dina M, Murti T and Rahmawati F. (2016) 'Kajian Drug Related Problems Penggunaan Obat Antihipertensi pada Pasien Diabetes Melitus Tipe 2 di

RSUD Temanggung.' *Thesis Fakultas Farmasi Univesitas Gadjah Mada Yogyakarta.*

Dirga D, Nugroho A, Pramantara D. (2019) 'Faktor-Faktor yang Memengaruhi Clinical Outcome Nyeri pada Pasien Diabetes Neuropati di Poliklinik Penyakit Dalam RSUD Kota Yogyakarta.' *Jurnal Kefarmasian Indonesia*. 30 Aug 2019 ; 9 (2):106-13.

Dormandy, J. A. *et al.* (2005) 'Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAZone Clinical Trial in macroVascular Events): A randomised controlled trial', *Lancet*, 366(9493), pp. 1279–1289. doi: 10.1016/S0140-6736(05)67528-9.

Draznin, B. *et al.* (2013) 'Pathways to quality inpatient management of hyperglycemia and diabetes: A call to action', *Diabetes Care*, 36(7), pp. 1807–1814. doi: 10.2337/dc12-2508.

Dworkin, R. H. *et al.* (2008) 'Interpreting the Clinical Importance of Treatment Outcomes in Chronic Pain Clinical Trials: IMMPACT Recommendations', *Journal of Pain*. doi: 10.1016/j.jpain.2007.09.005.

Dyck, R. *et al.* (2002) 'A comparison of rates , risk factors , and outcomes of gestational ...', *Diabetes Care*.

Fowler MJ (2008) 'Diabetes Foundation: Microvascular and Macrovascular Complications of Diabetes', *Clin Diab*, 26(3), pp. 77–82.

Handelsman, Y. *et al.* (2015) 'American Association of Clinical Endocrinologists and American College of Endocrinology - Clinical practice guidelines for developing a diabetes mellitus comprehensive care plan - 2015', *Endocrine Practice*, 21(4), pp. 1–87. doi: 10.4158/EP15672.GLSUPPL.

Hannibal, L. *et al.* (2016) 'Biomarkers and algorithms for the diagnosis of vitamin B 12 deficiency', *Frontiers in Molecular Biosciences*, 3(JUN). doi: 10.3389/fmolb.2016.00027.

Henry RR, Mudaliar S. (2012) 'Obesity and Type II Diabetes Mellitus.' In: Eckel RH, ed 'Obesity, Mechanism and Clinical Management.' *Lippincott Williams and Wilkins, Philadelphia*, 2003 (38): 1-3.

Hicks, C. W., Therapy, E. and Selvin, E. (2019) 'HHS Public Access', 19(10). doi: 10.1007/s11892-019-1212-8.Epidemiology.

Huri, H. Z. and Ling, L. C. (2013) 'Drug-related problems in type 2 diabetes mellitus patients with dyslipidemia'.

- Hvas, A. M. and Nexo, E. (2006) 'Diagnosis and treatment of vitamin B12 deficiency. An update', *Haematologica*, 91(11), pp. 1506–1512.
- Ismail-Beigi, F. *et al.* (2010) 'Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: An analysis of the ACCORD randomised trial', *The Lancet*. Elsevier Ltd, 376(9739), pp. 419–430. doi: 10.1016/S0140-6736(10)60576-4.
- De Jager, J. *et al.* (2010) 'Long term treatment with metformin in patients with type 2 diabetes and risk of vitamin B-12 deficiency: Randomised placebo controlled trial', *BMJ (Online)*, 340(7757), p. 1177. doi: 10.1136/bmj.c2181.
- Jambart, S. *et al.* (2011) 'Prevalence of painful diabetic peripheral neuropathy among patients with diabetes mellitus in the Middle East region', *Journal of International Medical Research*. doi: 10.1177/147323001103900204.
- Javed, A. *et al.* (2014) 'Gender based differences in diabetic peripheral neuropathy', 9(4).
- Juarez, D. T. *et al.* (2012) 'Factors Associated With Poor Glycemic Control or Wide Glycemic Variability Among Diabetes Patients in Hawaii , 2006 – 2009', 9, pp. 1–10.
- Juster-Switlyk, K. and Smith, A. G. (2016) 'Updates in diabetic peripheral neuropathy', *F1000Research*, 5(0), pp. 1–7. doi: 10.12688/f1000research.7898.1.
- Katzung, B. G. (2012) *Basic & Clinical Pharmacology, Basic and clinical Pharmacology*.
- Kemenkes.RI (2014) 'Prevalensi Penderita Diabetes', *Infodatin*. doi: 10.1002/ajmg.a.35913.
- Khawaja, N. *et al.* (2018) 'The prevalence and risk factors of peripheral neuropathy among patients with type 2 diabetes mellitus; The case of Jordan', *Diabetology and Metabolic Syndrome*. BioMed Central, 10(1), pp. 1–10. doi: 10.1186/s13098-018-0309-6.
- Kibirige, D. and Mwebaze, R. (2013) 'Vitamin B12 deficiency among patients with diabetes mellitus: Is routine screening and supplementation justified?', *Journal of Diabetes and Metabolic Disorders*, 12(1), pp. 2–7. doi: 10.1186/2251-6581-12-17.
- Kirby, M. (2003) 'Painful diabetic neuropathy - Current understanding and management for the primary care team', *British Journal of Diabetes and Vascular Disease*. doi: 10.1177/14746514030030021001.

- Ko, Sun-hye *et al.* (2014) 'Association of Vitamin B 12 Deficiency and Metformin Use in Patients with Type 2 Diabetes', pp. 965–972.
- Lacy, C.F, Amstrong, L.L, Goldman, M.P and Lance, L.L. (2014) 'Drug Information Handbook 23rd Edition.' *American Pharmacist Assosiation, Lexicomp.*
- Li, L. *et al.* (2015) 'Prevalence and risk factors of diabetic peripheral neuropathy in Type 2 diabetes mellitus patients with overweight/obese in Guangdong province, China', *Primary Care Diabetes*, 9(3), pp. 191–195. doi: 10.1016/j.pcd.2014.07.006.
- Moghissi, E. S. *et al.* (2009) 'American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control', *Diabetes Care*, 32(6), pp. 1119–1131. doi: 10.2337/dc09-9029.
- Moon, M. K. *et al.* (2017) 'Combination therapy of oral hypoglycemic agents in patients with type 2 diabetes mellitus', pp. 974–983.
- Nam Han Cho (chair) *et al.* (2017) *Eighth edition 2017, IDF Diabetes Atlas, 8th edition.* doi: [http://dx.doi.org/10.1016/S0140-6736\(16\)31679-8](http://dx.doi.org/10.1016/S0140-6736(16)31679-8).
- Out, M. *et al.* (2018) 'Long-term treatment with metformin in type 2 diabetes and methylmalonic acid: Post hoc analysis of a randomized controlled 4.3 year trial', *Journal of Diabetes and its Complications.* Elsevier Inc., 32(2), pp. 171–178. doi: 10.1016/j.jdiacomp.2017.11.001.
- PERKENI (2015) *Indonesia, P. E. (2015). Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia. PB. PERKENI.*
- Perkins, B. A. *et al.* (2001) 'Simple screening test for peripheral neuropathy in the diabetes clinic', *Diabetes Care*, 24(2), pp. 250–256. doi: 10.2337/diacare.24.2.250.
- Pharmaceutical care network Europe foundation (2017) 'Classification for Drug related problems V 8.01', *PCNE Classification*, pp. 1–10. Available at: https://www.pcne.org/upload/files/215_PCNE_classification_V8-01.pdf.
- Pop-busui, R. *et al.* (2017) 'Diabetic Neuropathy : A Position Statement by the American Diabetes Association', 40(January), pp. 136–154. doi: 10.2337/dc16-2042.
- Pop-Busui, R. *et al.* (2009) 'Prevalence of diabetic peripheral neuropathy and relation to glycemic control therapies at baseline in the BARI 2D cohort', *Journal of the Peripheral Nervous System.* doi: 10.1111/j.1529-

8027.2009.00200.x.

- Porcellati, F. *et al.* (2015) 'Pharmacokinetics and pharmacodynamics of insulin glargine given in the evening as compared with in the morning in type 2 diabetes', *Diabetes Care*, 38(3), pp. 503–512. doi: 10.2337/dc14-0649.
- Prasetyo GA. (2011) 'Lama Menderita Diabetes Melitus Tipe 2 Sebagai Faktor Risiko Nyeri Neuropati Diabetik.' *Thesis Fakultas Kedokteran Universitas Gadjah Mada Yogyakarta.*
- Riset Kesehatan Dasar. (2013) 'Riset Kesehatan Dasar.' *Badan Penelitian Dan Pengembangan Kesehatan Kementerian Indonesia.*
- Reinstatler, L. *et al.* (2012) 'Association of biochemical B 12 deficiency with metformin therapy and vitamin B 12 supplements: The National Health and Nutrition Examination Survey, 1999-2006', *Diabetes Care*, 35(2), pp. 327–333. doi: 10.2337/dc11-1582.
- Rodbard, H. W. *et al.* (2014) 'Treatment intensification with stepwise addition of prandial insulin aspart boluses compared with full basal-bolus therapy (FullSTEP Study): A randomised, treat-to-target clinical trial', *The Lancet Diabetes and Endocrinology*. Elsevier Ltd, 2(1), pp. 30–37. doi: 10.1016/S2213-8587(13)70090-1.
- Sabanayagam, C. *et al.* (2009) 'Relationship between glycated haemoglobin and microvascular complications: Is there a natural cut-off point for the diagnosis of diabetes?', *Diabetologia*, 52(7), pp. 1279–1289. doi: 10.1007/s00125-009-1360-5.
- Sadeli HA. (2008) 'Nyeri Neuropati Diabetika.' Dalam : Meliala L, Suryamiharja, Wirawan, Sadeli HA, Amir D, editor. 'Nyeri Neuropatik.' *Medigama Press Yogyakarta 2008 : 77-90.*
- Saltiel, A. R. and Olefsky, J. M. (1996) 'Thiazolidinediones in the treatment of insulin resistance and type II diabetes', *Diabetes*, 45(12), pp. 1661–1669. doi: 10.2337/diab.45.12.1661.
- Schreiber, A. K. (2015) 'Diabetic neuropathic pain: Physiopathology and treatment', *World Journal of Diabetes*, 6(3), p. 432. doi: 10.4239/wjd.v6.i3.432.
- Silver, B. *et al.* (2018) 'EADSG Guidelines: Insulin Therapy in Diabetes EXECUTIVE SUMMARY AND', *Diabetes Therapy*. Springer Healthcare, 9(2), pp. 449–492. doi: 10.1007/s13300-018-0384-6.
- Sjöström, L. *et al.* (2004) 'Lifestyle, diabetes, and cardiovascular risk factors 10

years after bariatric surgery', *New England Journal of Medicine*. doi: 10.1056/NEJMoa035622.

Soelistijo, S. *et al.* (2015) *Konsensus Pengelolaan Dan Pencegahan Diabetes Melitus Tipe 2 Di Indonesia 2015, Perkeni*. Available at: <https://www.google.com/url?sa=t&source=web&rct=j&url=https://pbperkeni.or.id/wp-content/uploads/2019/01/4.-Konsensus-Pengelolaan-dan-Pencegahan-Diabetes-melitus-tipe-2-di-Indonesia-PERKENI-2015.pdf&ved=2ahUKEwjy8KOs8cfoAhXCb30KHQb1Ck0QFjADegQIBhAB&usg=AOv>.

Strowig, S. M. and Raskin, P. (2005) 'Combination therapy using metformin or thiazolidinediones and insulin in the treatment of diabetes mellitus', pp. 633–641. doi: 10.1111/j.1462.

Su, J. bin *et al.* (2018) 'HbA1c variability and diabetic peripheral neuropathy in type 2 diabetic patients', *Cardiovascular Diabetology*. BioMed Central, 17(1), pp. 1–9. doi: 10.1186/s12933-018-0693-0.

Tarigan, TJE. *et al.* (2015) 'Profile and analysis of diabetes chronic complications in Outpatient Diabetes Clinic of Cipto Mangunkusumo Hospital, Jakarta.' *Medical Journal of Indonesia*, 24(3), 156. doi:10.13181/mji.v24i3.1249

Tesfaye, S. *et al.* (2010) 'Diabetic neuropathies: Update on definitions, diagnostic criteria, estimation of severity, and treatments', in *Diabetes Care*. doi: 10.2337/dc10-1303.

Treede, R. (2008) 'Neuropathic Pain: Redefinition and a Grading System for Clinical and Research Purposes', *Neurology*, 72.

Triplitt, C.L, Reasner, C.A and Isley, W.L. (2005) 'Diabetes Melitus,' in Dipiro, J.T, Talber, R.L, Yee, G.C, Matzke, G.R, Wells, B.G, Posey, L.M. 'Endocrinologic Disorders : Diabetes mellitus Pharmacotherapy A Pathophysiologic Approach, Sixth Edition,' *McGraw-Hill Companies* , pp. 1333-1363.

UK Prospective Diabetes Study Group (1998) 'Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). UK Prospective Diabetes Study (UKPDS) Group.', *Lancet*.

Wang, Z. *et al.* (2010) 'Dose-response effects of insulin glargine in type 2 diabetes', *Diabetes Care*, 33(7), pp. 1555–1560. doi: 10.2337/dc09-2011.

Wells, B.G, DiPiro, C.V, DiPiro, J.T and Schwinghammer, T.L. (2014) 'Endocrinologic Disorder: Diabetes Mellitus. Pharmacotherapy Handbook 9th

Edition.' *McGraw-Hill Education*. (19):161-175.

Whalen, K, Finkel, R, Panavelil, TA. (2015) 'Lippincott Illustrated Reviews: Pharmacology 6th edition.' *Wolters Kluwer*.

Yamanouchi, T. and Press, D. (2010) 'Concomitant therapy with pioglitazone and insulin for the treatment of type 2 diabetes', pp. 189–197.

Yigazu, D. M. and Desse, T. A. (2017) 'Glycemic control and associated factors among type 2 diabetic patients at Shanan Gibe Hospital, Southwest Ethiopia', *BMC Research Notes*. BioMed Central, 10(1), pp. 1–6. doi: 10.1186/s13104-017-2924-y.

Zaman Huri, H. and Fun Wee, H. (2013) 'Drug related problems in type 2 diabetes patients with hypertension: A cross-sectional retrospective study', *BMC Endocrine Disorders*. doi: 10.1186/1472-6823-13-2.

Zazuli, Z., Rohaya, A. and Adnyana, K. (2017) 'Drug-related problems in Type 2 diabetic patients with hypertension in Cimahi, West Java, Indonesia: A prospective study', 2017(2), pp. 298–304.