

DAFTAR PUSTAKA

- Afrozeh, F., Askarizadeh, G., & Towhidi, A. (2019). The Effectiveness of Cognitive-Behavioral Stress Management Training on Chronic Fatigue and Death Anxiety in Women Undergoing Hemodialysis. *Iranian Journal of Psychiatric Nursing (IJPN)*, 7(4). <https://doi.org/10.21859/ijpn-07408>
- Afshar, M., Mohsenzadeh, A., Gilasi, H., & Sadeghi-Gandomani, H. (2018). The effects of guided imagery on state and trait anxiety and sleep quality among patients receiving hemodialysis: A randomized controlled trial. *Complementary Therapies in Medicine*, 40, 37–41. <https://doi.org/10.1016/j.ctim.2018.07.006>
- Aggarwal, H. K., Jain, D., Dabas, G., & Yadav, R. K. (2017). Prevalence of Depression, Anxiety and Insomnia in Chronic Kidney Disease Patients and their Co-Relation with the Demographic Variables. *Prilozi (Makedonska Akademija Na Naukite I Umetnostite. Oddelenie Za Medicinski Nauki)*, 38(2), 35–44. <https://doi.org/10.1515/prilozi-2017-0020>
- Aisara, S., Azmi, S., & Yanni, M. (2018). Gambaran Klinis Penderita Penyakit Ginjal Kronik yang Menjalani Hemodialisis di RSUP Dr. M. Djamil Padang. *Jurnal Kesehatan Andalas*, 7(1), 42. <https://doi.org/10.25077/jka.v7.i1.p42-50.2018>
- Ait-Daoud, N., Hamby, A. S., Sharma, S., & Blevins, D. (2018). A Review of Alprazolam Use, Misuse, and Withdrawal: *Journal of Addiction Medicine*, 12(1), 4–10. <https://doi.org/10.1097/ADM.0000000000000350>
- Alirudin, C., Hendellyn, A., Sumada, I. K., Yuliani, D., & Sunaka, I. W. (2020). Profil Gangguan Fungsi Kognitif pada Pasien Hemodialisis di Rumah Sakit Umum Daerah Wangaya. *Callosum Neurology - Jurnal Berkala Neurologi Bali*, 6(2), 42–46. <https://doi.org/10.29342/cnj.v3i2/97>
- Alshraifeen, A., Alnuaimi, K., Al-Rawashdeh, S., Ashour, A., Al-Ghabeesh, S., & Al-Smadi, A. (2020). Spirituality, Anxiety and Depression Among People Receiving Hemodialysis Treatment in Jordan: A Cross-Sectional Study. *Journal of Religion and Health*. <https://doi.org/10.1007/s10943-020-00988-8>
- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR*. <https://doi.org/10.1176/appi.books.9780890423349.5847>
- American Psychological Association. (2017). *Beyond worry: How psychologists help with anxiety disorders*. <https://www.apa.org/helpcenter/anxiety>
- Aprilia, S. (2016). *Faktor-faktor yang Berhubungan dengan Tingkat Kecemasan Pasien Gagal Ginjal Kronik yang Menjalani Hemodialisis di PKU Muhammadiyah 2 Yogyakarta [Unpublished Undergraduate Thesis]*. Universitas Muhammadiyah Yogyakarta.
- Ariani, T. A. (2017). The Potential of Psychosocial Health Indicators That Effects to Chronic Renal Failure (CRF) Patient With Cognitive Behavior Therapy (CBT) Based. *Jurnal Kesehatan Hesti Wira Sakti*, 5(1), 60–67.
- Arikunto, S. (2006). *Prosedur Penelitian Suatu Pendekatan Praktik*. Jakarta: Rineka Cipta.

- Arjadi, R. (2012). *Terapi Kognitif-Perilaku untuk Menangani Depresi pada Lanjut Usia (Tesis)*. Fakultas Psikologo, Program Magister Profesi Psikologi Klinis Dewasa, Universitas Indonesia.
- Arsa, P. S. A., Sudiana, I. K., & Hidayati, L. (2018). *Description of Factors Affecting Anxiety on End Stage Renal Disease Patients Which The Hemodialysis at Lavalette Hospital in Malang*. The 9th International Nursing Conference 2018.
- Astuti, P., Herawati, T., & Kariasa, I. M. (2016). *Faktor-Faktor Yang Berhubungan Dengan Self Management Pada Pasien Hemodialisis di Kota Bekasi*. 12.
- Australian Institute of Health and Welfare (Ed.). (2005). *Chronic kidney disease in Australia, 2005*. Australian Institute of Health and Welfare.
- Avramovic, M., & Stefanovic, V. (2012). Health-Related Quality of Life in Different Stages of Renal Failure. *Artificial Organs*, 36(7), 581–589. <https://doi.org/10.1111/j.1525-1594.2011.01429.x>
- Avşar, U., Avşar, U. Z., Cansever, Z., Yucel, A., Cankaya, E., Certez, H., Keles, M., Aydınli, B., & Yucelf, N. (2015). Caregiver Burden, Anxiety, Depression, and Sleep Quality Differences in Caregivers of Hemodialysis Patients Compared With Renal Transplant Patients. *Transplantation Proceedings*, 47(5), 1388–1391. <https://doi.org/10.1016/j.transproceed.2015.04.054>
- Banks, S. J., Eddy, K. T., Angstadt, M., Nathan, P. J., & Phan, K. L. (2007). Amygdala–frontal connectivity during emotion regulation. *Social Cognitive and Affective Neuroscience*, 2(4), 303–312. <https://doi.org/10.1093/scan/nsm029>
- Barlow, D. H. (2004). *Anxiety and its disorders. The nature and treatment of anxiety and panic (2nd ed.)*. New York: The Guilford Press.
- Bieber, B., Qian, J., Anand, S., Yan, Y., Chen, N., Wang, M., Wang, M., Zuo, L., Hou, F. F., Pisoni, R. L., Robinson, B. M., & Ramirez, S. P. B. (2014). Two-times weekly hemodialysis in China: Frequency, associated patient and treatment characteristics and Quality of Life in the China Dialysis Outcomes and Practice Patterns study. *Nephrology, Dialysis, Transplantation: Official Publication of the European Dialysis and Transplant Association - European Renal Association*, 29(9), 1770–1777. <https://doi.org/10.1093/ndt/gft472>
- Bjelland, I., Dahl, A. A., Haug, T. T., & Neckelmann, D. (2002). The validity of the Hospital Anxiety and Depression Scale: An updated literature review. *Journal of Psychosomatic Research*, 52(2), 69–77. [https://doi.org/10.1016/S0022-3999\(01\)00296-3](https://doi.org/10.1016/S0022-3999(01)00296-3)
- Bond, F. W., & Dryden. (2002). *Handbook of cognitive behaviour therapy*. Chichester: John Wiley & Sons, Ltd.
- Bossola, M., Di Stasio, E., Giungi, S., Rosa, F., & Tazza, L. (2015). Fatigue Is Associated With Serum Interleukin-6 Levels and Symptoms of Depression in Patients on Chronic Hemodialysis. *Journal of Pain and Symptom Management*, 49(3), 578–585. <https://doi.org/10.1016/j.jpainsymman.2014.07.009>
- Bura, U. D. K., Hasmono, D., Surdijati, S., & Hartono, R. (2017). Studi Penggunaan Antihipertensi Golongan Angiotensin Receptor Blockers

- Terhadap Pasien Gagal Ginjal Kronik Stadium V di Rumah Sakit Bhayangkara Surabaya. *J PHARM SCI*, 4(2), 8.
- Butler, A., Chapman, J., Forman, E., & Beck, A. (2006). The empirical status of cognitive-behavioral therapy: A review of meta-analyses. *Clinical Psychology Review*, 26(1), 17–31. <https://doi.org/10.1016/j.cpr.2005.07.003>
- Cahyani, N. D., Tyaswati, J. E., Rachmawati, D. A., & Kalimantan, J. (2016). *Hubungan antara Tingkat Kecemasan dengan Kualitas Hidup pada Pasien Chronic Kidney Disease (CKD) yang Menjalani Hemodialisis di RSD dr. Soebandi Jember*. 4, 8.
- Caninsti, R. (2013). Kecemasan dan Depresi pada Pasien Gagal Ginjal Kronis yang Menjalani Terapi Hemodialisis. *Jurnal Psikologi Ulayat*, 1(2). <https://doi.org/10.24854/jpu22013-22>
- CDC. (2014). National Chronic Kidney Disease Fact Sheet, 2014. *Centers for Disease Control and Prevention, US*, 1–3.
- Chen, H.-Y., Cheng, I.-C., Pan, Y.-J., Chiu, Y.-L., Hsu, S.-P., Pai, M.-F., Yang, J.-Y., Peng, Y.-S., Tsai, T.-J., & Wu, K.-D. (2011). Cognitive-behavioral therapy for sleep disturbance decreases inflammatory cytokines and oxidative stress in hemodialysis patients. *Kidney International*, 80(4), 415–422. <https://doi.org/10.1038/ki.2011.151>
- Cukor, D., Coplan, J., Brown, C., Friedman, S., Newville, H., Safier, M., Spielman, L. A., Peterson, R. A., & Kimmel, P. L. (2008). Anxiety Disorders in Adults Treated by Hemodialysis: A Single-Center Study. *American Journal of Kidney Diseases*, 52(1), 128–136. <https://doi.org/10.1053/j.ajkd.2008.02.300>
- Dagnan, D., Chadwick, P., & Trower, P. (2000). Psychometric properties of the Hospital Anxiety and Depression Scale with a population of members of a depression self-help group. *The British Journal of Medical Psychology*, 73 (Pt 1), 129–137. <https://doi.org/10.1348/000711200160255>
- Dani, R., & Utami, G. T. (2015). Hubungan Motivasi, Harapan, dan Dukungan Petugas Kesehatan terhadap Kepatuhan Pasien Gagal Ginjal Kronik untuk Menjalani Hemodialisis. *Jurnal Online Mahasiswa*, 2(2), 10.
- Daugirdas, J. T., Depner, T. A., Inrig, J., Mehrotra, R., Rocco, M. V., Suri, R. S., Weiner, D. E., Greer, N., Ishani, A., MacDonald, R., Olson, C., Rutks, I., Slinin, Y., Wilt, T. J., Rocco, M., Kramer, H., Choi, M. J., Samaniego-Picota, M., Scheel, P. J., ... Brereton, L. (2015). KDOQI Clinical Practice Guideline for Hemodialysis Adequacy: 2015 Update. *American Journal of Kidney Diseases*, 66(5), 884–930. <https://doi.org/10.1053/j.ajkd.2015.07.015>
- Davidson, J. R. T. (2009). First-Line Pharmacotherapy Approaches for Generalized Anxiety Disorder. *The Journal of Clinical Psychiatry*, 70(suppl 2), 25–31. <https://doi.org/10.4088/JCP.s.7002.05>
- Dehkordi, L. M., & Shahgholian, N. (2013). An investigation of coping styles of hemodialysis patients. *Iranian Journal of Nursing and Midwifery Research*, 18(1), 42–46.
- Doehrmann, O., Ghosh, S. S., Polli, F. E., Reynolds, G. O., Horn, F., Keshavan, A., Triantafyllou, C., Saygin, Z. M., Whitfield-Gabrieli, S., Hofmann, S. G., Pollack, M., & Gabrieli, J. D. (2013). Predicting Treatment Response

- in Social Anxiety Disorder From Functional Magnetic Resonance Imaging. *JAMA Psychiatry*, 70(1), 87–97. <https://doi.org/10.1001/2013.jamapsychiatry.5>
- Dwi, A. (2018). Self Management terhadap Psychosocial Adjustment Pasien Penyakit Ginjal Kronis dengan Hemodialisa. *Jurnal Keperawatan Soedirman*, 12(3), 181. <https://doi.org/10.20884/1.jks.2017.12.3.698>
- Dziubek, W., Kowalska, J., Kuztal, M., Rogowski, Ł., Gołębiowski, T., Nikifur, M., Szczepańska-Gieracha, J., Zembroń-Łacny, A., Klinger, M., & Woźniewski, M. (2016). The Level of Anxiety and Depression in Dialysis Patients Undertaking Regular Physical Exercise Training—A Preliminary Study. *Kidney and Blood Pressure Research*, 41(1), 86–98. <https://doi.org/10.1159/000368548>
- Ediz, B., Ozcakil, A., & Bilgel, N. (2017). Depression and anxiety among medical students: Examining scores of the beck depression and anxiety inventory and the depression anxiety and stress scale with student characteristics. *Cogent Psychology*, 4(1). <https://doi.org/10.1080/23311908.2017.1283829>
- Erwina, I. (2010). *Pengaruh Cognitive Behavior Therapy Terhadap Post-Traumatic Stress Disorder pada Penduduk Pasca Gempa di kelurahan Air Tawar Barat Kecamatan Padang Utara Propinsi Sumatera Barat (Tesis)*. Fakultas Ilmu Keperawatan, Program Magister Ilmu Keperawatan, Kekhususan keperawatan Jiwa, Universitas Indonesia.
- Etkin, A., & Wager, T. D. (2007). Functional Neuroimaging of Anxiety: A Meta-Analysis of Emotional Processing in PTSD, Social Anxiety Disorder, and Specific Phobia. *Am J Psychiatry*, 13.
- Fadlilah, S. (2019). Faktor-Faktor yang Berhubungan dengan Kualitas Hidup Pasien Hemodialisis. *Jurnal Kesehatan*, 10(2), 284. <https://doi.org/10.26630/jk.v10i2.1454>
- Ferrey, A. J., Choi, G., Hanna, R. M., Chang, Y., Tantisattamo, E., Ivaturi, K., Park, E., Nguyen, L., Wang, B., Tonthat, S., Rhee, C. M., Reddy, U., Lau, W. L., Huang, S. S., Gohil, S., Amin, A. N., Hsieh, L., Cheng, T. T., Lee, R. A., & Kalantar-Zadeh, K. (2020). A Case of Novel Coronavirus Disease 19 in a Chronic Hemodialysis Patient Presenting with Gastroenteritis and Developing Severe Pulmonary Disease. *American Journal of Nephrology*, 51(5), 337–342. <https://doi.org/10.1159/000507417>
- FIK-UI. (2015). *Draft Modul Cognitive Behavior Therapy (CBT)*. Universitas Indonesia.
- Foji, S., Tadayonfar, M. Alr., Mohsenpour, M., & Rakhshani, M. H. (2015). The study of the effect of guided imagery on pain, anxiety and some other hemodynamic factors in patients undergoing coronary angiography. *Complementary Therapies in Clinical Practice*, 21(2), 119–123. <https://doi.org/10.1016/j.ctcp.2015.02.001>
- Forwaty, E., Malini, H., & Oktarina, E. (2019). Pengaruh Intradialytic Range of Motion (ROM) Exercise terhadap Depresi, Insomnia dan Asupan Nutrisi pada Pasien Hemodialisis. *Jurnal Kesehatan Andalas*, 8(3), 529–536. <https://doi.org/10.25077/jka.v8i3.1038>
- Geglorian, T. R., & Arif, M. S. (2016). Pengaruh Cognitive Therapy Terhadap Tingkat Ansietas Pada Pasien Yang Akan Menjalani Hemodialisa Di RS

- Panti Wilasa Citarum Semarang. *Jurnal Ilmu Keperawatan dan Kebidanan*, 5(1), 1–11.
- Gela, D., & Mengistu, D. (2018). Self-management and associated factors among patients with end-stage renal disease undergoing hemodialysis at health facilities in Addis Ababa, Ethiopia. *International Journal of Nephrology and Renovascular Disease*, Volume 11, 329–336. <https://doi.org/10.2147/IJNRD.S184671>
- Gerasimoula, K., Lefkothea, L., Maria, L., Victoria, A., Paraskevi, T., & Maria, P. (2015). Quality of Life in Hemodialysis Patients. *Materia Socio Medica*, 27(5), 305. <https://doi.org/10.5455/msm.2015.27.305-309>
- Gerogianni, G., Babatsikou, F., Polikandrioti, M., & Grapsa, E. (2019). Management of anxiety and depression in haemodialysis patients: The role of non-pharmacological methods. *International Urology and Nephrology*, 51(1), 113–118. <https://doi.org/10.1007/s11255-018-2022-7>
- Gerontoukou, E.-I., Michaelidou, S., Rekleiti, M., Saridi, M., & Souliotis, K. (2015). Investigation of Anxiety and Depression in Patients with Chronic Diseases. *Health Psychology Research*, 3(2). <https://doi.org/10.4081/hpr.2015.2123>
- Ghonemy, T., Farag, S., Soliman, S., El-okely, A., & El-hendy, Y. (2016). Epidemiology and risk factors of chronic kidney disease in the El-Sharkia Governorate, Egypt. *Saudi Journal of Kidney Diseases and Transplantation*, 27(1), 111. <https://doi.org/10.4103/1319-2442.174137>
- Giusti, L., Ussorio, D., Salza, A., Malavolta, M., Aggio, A., Bianchini, V., Casacchia, M., & Roncone, R. (2018). Preliminary study of effects on paranoia ideation and jumping to conclusions in the context of group treatment of anxiety disorders in young people. *Early Intervention in Psychiatry*, 12(6), 1072–1080. <https://doi.org/10.1111/eip.12415>
- Golper, T. A., Fissell, R., Fissell, W. H., Hartle, P. M., Sanders, M. L., & Schulman, G. (2014). Hemodialysis: Core Curriculum 2014. *American Journal of Kidney Diseases*, 63(1), 153–163. <https://doi.org/10.1053/j.ajkd.2013.07.028>
- Griva, K., Lam, K. F. Y., Nandakumar, M., Ng, J. H., McBain, H., & Newman, S. P. (2018). The effect of brief self-management intervention for hemodialysis patients (HED-SMART) on trajectories of depressive and anxious symptoms. *Journal of Psychosomatic Research*, 113, 37–44. <https://doi.org/10.1016/j.jpsychores.2018.07.012>
- Gunarti, C. S. I., & Pohan, L. D. (2018). Cognitive Behavioral Therapy for Decreasing Anxiety in an Elderly Female Patient Undergoing Hemodialysis: A Single-case Study. *Proceedings of the 1st International Conference on Intervention and Applied Psychology (ICIAP 2017)*. 1st International Conference on Intervention and Applied Psychology (ICIAP 2017), Depok, Indonesia. <https://doi.org/10.2991/iciap-17.2018.32>
- Hare, J., Clark-Carter, D., & Forshaw, M. (2014). A randomized controlled trial to evaluate the effectiveness of a cognitive behavioural group approach to improve patient adherence to peritoneal dialysis fluid restrictions: A pilot study. *Nephrology Dialysis Transplantation*, 29(3), 555–564. <https://doi.org/10.1093/ndt/gft477>

- Hathaway, W. R., & Newton, B. W. (2020). Neuroanatomy, Prefrontal Cortex. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK499919/>
- Hill, N. R., Fatoba, S. T., Oke, J. L., Hirst, J. A., O'Callaghan, C. A., Lasserson, D. S., & Hobbs, F. D. R. (2016). Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis. *PLOS ONE*, *11*(7), e0158765. <https://doi.org/10.1371/journal.pone.0158765>
- Hofmann, S. G., Asmundson, G. J. G., & Beck, A. T. (2013). The Science of Cognitive Therapy. *Behavior Therapy*, *44*(2), 199–212. <https://doi.org/10.1016/j.beth.2009.01.007>
- Hou, Y., Li, X., Yang, L., Liu, C., Wu, H., Xu, Y., Yang, F., & Du, Y. (2014). Factors associated with depression and anxiety in patients with end-stage renal disease receiving maintenance hemodialysis. *International Urology and Nephrology*, *46*(8), 1645–1649. <https://doi.org/10.1007/s11255-014-0685-2>
- Ikizler, T. A., & Klinger, A. S. (2020). Minimizing the risk of COVID-19 among patients on dialysis. *Nature Reviews Nephrology*, *16*(6), 311–313. <https://doi.org/10.1038/s41581-020-0280-y>
- Imelda, F., Susalit, E., Marbun, M. B. M., & Rumende, C. M. (2017). Gambaran Klinis dan Kualitas Hidup Pasien Penyakit Ginjal Tahap Akhir yang Menjalani Hemodialisis Dua Kali Dibandingkan Tiga Kali Seminggu. *Jurnal Penyakit Dalam Indonesia*, *4*(3), 128. <https://doi.org/10.7454/jpdi.v4i3.139>
- Institute for Quality and Efficiency in Health Care (IQWiG). (2018). *Dialysis in chronic kidney disease*. Institute for Quality and Efficiency in Health Care (IQWiG). <https://www.ncbi.nlm.nih.gov/books/NBK492979/>
- Irfan, Wawomeo, A., & Kambuno, N. T. (2019). Hepatitis B Virus Infection in Hemodialysis patient at Prof. DR. W.Z. Johannes Kupang Hospital, East Nusa Tenggara. *Jurnal Kesehatan Primer*, *4*(1), 63–69. <https://doi.org/10.31965/jkp>
- Ismoyowati, T. W. (2018). Pengaruh Acceptance and Commitment Therapy terhadap Perubahan Kecemasan dan Selfefficacy pada Pasien dengan Hemodialisa Di RS Sint Carolus Jakarta. *Jurnal Kesehatan*, *5*(2), 44–58. <https://doi.org/10.35913/jk.v5i2.95>
- Julian, L. J. (2011). Measures of anxiety: State-Trait Anxiety Inventory (STAI), Beck Anxiety Inventory (BAI), and Hospital Anxiety and Depression Scale-Anxiety (HADS-A). *Arthritis Care & Research*, *63*(S11), S467–S472. <https://doi.org/10.1002/acr.20561>
- Julianty, S. A. H., Yustina, I., & Ardinata, D. (2015). Faktor-faktor yang Berhubungan dengan Tingkat Kecemasan Pasien Hemodialisis di RSUD Dr. Pirngadi Medan. *Idea Nursing Journal*, *VI*(3).
- Kaczurkin, A. N., & Foa, E. B. (2015). Cognitive-behavioral therapy for anxiety disorders: An update on the empirical evidence. *Dialogues in Clinical Neuroscience*, *17*(3), 337–346.
- Kaplan, H. I., Sadock, B. J., & Grebb, J. A. (2010). *Sinopsis Psikiatri Jilid 2*. Terjemahan Widjaja Kusuma. Jakarta: Binarupa Aksara.
- Karkar, A. (2013). Advances in Hemodialysis Techniques. In H. Suzuki (Ed.), *Hemodialysis*. InTech. <https://doi.org/10.5772/52444>

- Kementrian Kesehatan RI. (2013). *Riset Kesehatan Dasar*.
- Kidney Disease: Improving Global Outcomes (KDIGO). (2012). *KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease*. *Kidney Int*; 3 (suppl): 1-150.
- Kidney Disease: Improving Global Outcomes (KDIGO). (2013). Chapter 1: Definition and classification of CKD. *Kidney International Supplements*, 3(1), 19–62. <https://doi.org/10.1038/kisup.2012.64>
- Kidney Disease: Improving Global Outcomes (KDIGO). (2018). KDIGO 2018 Clinical Practice Guideline for the Prevention, Diagnosis, Evaluation, and Treatment of hepatitis C in Chronic Kidney Disease. *Official Journal of the International Society of Nephrology*, 8(3).
- Klumpp, H., Fitzgerald, D. A., Angstadt, M., Post, D., & Phan, K. L. (2014). Neural response during attentional control and emotion processing predicts improvement after cognitive behavioral therapy in generalized social anxiety disorder. *Psychological Medicine*, 44(14), 3109–3121. <https://doi.org/10.1017/S0033291714000567>
- Klumpp, Heide, Keutmann, M. K., Fitzgerald, D. A., Shankman, S. A., & Phan, K. L. (2014). Resting state amygdala-prefrontal connectivity predicts symptom change after cognitive behavioral therapy in generalized social anxiety disorder. *Biology of Mood & Anxiety Disorders*, 4(1), 14. <https://doi.org/10.1186/s13587-014-0014-5>
- Kumar, V., Khandelia, V., & Garg, A. (2018). Depression and anxiety in patients with chronic kidney disease undergoing hemodialysis. *Annals of Indian Psychiatry*, 2(2), 115. https://doi.org/10.4103/aip.aip_12_18
- Lee, S., Kang, S., Joo, Y. S., Lee, C., Nam, K. H., Yun, H.-R., Park, J. T., Chang, T. I., Yoo, T.-H., Kim, S. W., Oh, K.-H., Kim, Y. H., Park, S. K., Kang, S.-W., Choi, K. H., Ahn, C., & Han, S. H. (2020). Smoking, Smoking Cessation, and Progression of Chronic Kidney Disease: Results From KNOW-CKD Study. *Nicotine & Tobacco Research*, ntaa071. <https://doi.org/10.1093/ntr/ntaa071>
- Lerma, A., Perez-Grovas, H., Bermudez, L., Peralta-Pedrero, M. L., Robles-García, R., & Lerma, C. (2017). Brief cognitive behavioural intervention for depression and anxiety symptoms improves quality of life in chronic haemodialysis patients. *Psychology and Psychotherapy*, 90(1), 105–123. <https://doi.org/10.1111/papt.12098>
- Lestari, I. G., & Isnaini, N. (2018). Pengaruh Self Management terhadap Tekanan Darah Lansia yang Mengalami Hipertensi. *Indonesian Journal for Health Sciences*, 13.
- Levey, A. S., Atkins, R., Coresh, J., Cohen, E. P., Collins, A. J., Eckardt, K.-U., Nahas, M. E., Jaber, B. L., Jadoul, M., Levin, A., Powe, N. R., Rossert, J., Wheeler, D. C., Lameire, N., & Eknoyan, G. (2007). Chronic kidney disease as a global public health problem: Approaches and initiatives – a position statement from Kidney Disease Improving Global Outcomes. *Kidney International*, 72(3), 247–259. <https://doi.org/10.1038/sj.ki.5002343>
- Lin, M.-Y., Liu, M. F., Hsu, L.-F., & Tsai, P.-S. (2017). Effects of self-management on chronic kidney disease: A meta-analysis. *International*

- Journal of Nursing Studies*, 74, 128–137.
<https://doi.org/10.1016/j.ijnurstu.2017.06.008>
- Ma, T. K.-W., & Li, P. K.-T. (2016). Depression in dialysis patients. *Nephrology*, 21(8), 639–646. <https://doi.org/10.1111/nep.12742>
- Maesaroh, Waluyo, A., & Jumaiyah, W. (2020). Faktor-faktor yang Berhubungan dengan Terjadinya Fatigue pada Pasien Hemodialisis. *Syntax Literate: Jurnal Ilmiah Indonesia*, 5(4), 110–120.
- Maguire, P. N., Clark, G. I., & Wootton, B. M. (2018). The efficacy of cognitive behavior therapy for the treatment of perinatal anxiety symptoms: A preliminary meta-analysis. *Journal of Anxiety Disorders*, 60, 26–34. <https://doi.org/10.1016/j.janxdis.2018.10.002>
- Maidah, D. (2013). Self Injury pada Mahasiswa (Study Kasus pada Mahasiswa Pelaku Self Injury). *Developmental and Clinical Psychology*, 2(1), 6–13.
- Mailani, F., & Andriani, R. F. (2017). Hubungan Dukungan Keluarga dengan Kepatuhan Diet pada Pasien Gagal Ginjal Kronik yang Menjalani Hemodialisis. *Jurnal Endurance*, 2(3), 416. <https://doi.org/10.22216/jen.v2i3.2379>
- Manafe, R. P. (2018). Efektivitas Cognitive Behavioral Therapy Untuk Menurunkan Distres Akibat Proses Hemodialisis. *Jurnal Ilmiah Mahasiswa Universitas Surabaya*, 7(1), 2277–2293.
- Manthey, L., van Veen, T., Giltay, E. J., Stoop, J. E., Neven, A. K., Penninx, B. W. J. H., & Zitman, F. G. (2011). Correlates of (inappropriate) benzodiazepine use: The Netherlands Study of Depression and Anxiety (NESDA): Correlates of benzodiazepine use. *British Journal of Clinical Pharmacology*, 71(2), 263–272. <https://doi.org/10.1111/j.1365-2125.2010.03818.x>
- Mardhatillah, Arsin, A., Syafar, M., & Hardianti, A. (2020). Ketahanan Hidup Pasien Penyakit Ginjal Kronik yang Menjalani Hemodialisis di RSUP Dr. Wahidin Sudirohusodo Makassar. *Jurnal Kesehatan Masyarakat Maritim*, 3(1), 21–33.
- Marthoenis, M., Syukri, M., Abdullah, A., Tandi, T. M. R., Putra, N., Laura, H., Setiawan, A., Sofyan, H., & Schouler-Ocak, M. (2020). Quality of life, depression, and anxiety of patients undergoing hemodialysis: Significant role of acceptance of the illness. *The International Journal of Psychiatry in Medicine*, 009121742091338. <https://doi.org/10.1177/0091217420913382>
- Mitema, D., & Jaar, B. G. (2016). How Can We Improve the Quality of Life of Dialysis Patients? *Seminars in Dialysis*, 29(2), 93–102. <https://doi.org/10.1111/sdi.12467>
- Moledina, D. G., & Wilson, F. P. (2015). Pharmacologic Treatment of Common Symptoms in Dialysis Patients: A Narrative Review. *Seminars in Dialysis*, 28(4), 377–383. <https://doi.org/10.1111/sdi.12378>
- Mosleh, H., Alenezi, M., Al johani, S., Alsani, A., Fairaq, G., & Bedaiwi, R. (2020). Prevalence and Factors of Anxiety and Depression in Chronic Kidney Disease Patients Undergoing Hemodialysis: A Cross-sectional Single-Center Study in Saudi Arabia. *Cureus*, 12(1). <https://doi.org/10.7759/cureus.6668>

- Ng, C. Z., Tang, S. C., Chan, M., Tran, B. X., Ho, C. S., Tam, W. W., & Ho, R. C. (2019). A systematic review and meta-analysis of randomized controlled trials of cognitive behavioral therapy for hemodialysis patients with depression. *Journal of Psychosomatic Research*, *126*, 109834. <https://doi.org/10.1016/j.jpsychores.2019.109834>
- Ng, H. J., Tan, W. J., Mooppil, N., Newman, S., & Griva, K. (2015). Prevalence and patterns of depression and anxiety in hemodialysis patients: A 12-month prospective study on incident and prevalent populations. *British Journal of Health Psychology*, *20*(2), 374–395. <https://doi.org/10.1111/bjhp.12106>
- NICE. (2011). *Generalised Anxiety Disorder and Panic Disorder in Adults: Management*. National Institute for Health and Care Excellence Clinical Guideline. <https://www.nice.org.uk/guidance/cg113>
- Nie, J., Zhu, S., Cui, W., Wu, H., Cai, Y., Wen, X., & Li, Q. (2020). *Psychological status and intervention strategy of maintenance hemodialysis patients in COVID-19 epidemic* [Preprint]. In Review. <https://doi.org/10.21203/rs.3.rs-38725/v1>
- Nugraha, A., Rahmah, G. U., & Budhiaji, P. (2020). Effect of Family Support Towards Elderly Hemodialysis Patients' Anxiety Level. *Basic and Applied Nursing Research Journal*, *1*(1), 5. <https://doi.org/x.doi.org/10.11594/banj.01.01.05>
- Nunnally, J. C., & Bernstein, I. H. (1994). *Psychometric Theory* (3rd ed). New York: McGraw-Hill.
- Nurjanah, D. E., & Elly, N. (2014). Pengaruh Terapi Kognitif Terhadap Tingkat Kecemasan pada Pasien Gagal Ginjal Kronik yang Menjalani Hemodialisa. *Jurnal Media Kesehatan*, *7*.
- Oesmarjoedi, A. K. (2003). *Pendekatan Cognitive Behavior Dalam Psikoterapi*.
- Orgeta, V., Qazi, A., Spector, A., & Orrell, M. (2015). Psychological treatments for depression and anxiety in dementia and mild cognitive impairment: Systematic review and meta-analysis. *British Journal of Psychiatry*, *207*(4), 293–298. <https://doi.org/10.1192/bjp.bp.114.148130>
- Pascoe, M. C., Thompson, D. R., Castle, D. J., McEvedy, S. M., & Ski, C. F. (2017). Psychosocial Interventions for Depressive and Anxiety Symptoms in Individuals with Chronic Kidney Disease: Systematic Review and Meta-Analysis. *Frontiers in Psychology*, *8*, 992. <https://doi.org/10.3389/fpsyg.2017.00992>
- Paulus, E. S. (2015). “Irrational Beliefs” dalam Konteks Kehidupan Seminars. *Jurnal Teologi*, *4*(1), 25–40. <https://doi.org/10.24071/jt.v4i1.468>
- PERNEFRI. (2015). *8th report of Indonesian Renal Registry* (p. 45).
- Phelps, E. A., & LeDoux, J. E. (2005). Contributions of the Amygdala to Emotion Processing: From Animal Models to Human Behavior. *Neuron*, *48*(2), 175–187. <https://doi.org/10.1016/j.neuron.2005.09.025>
- Pranandari, R., & Supadmi, W. (2015). Faktor Risiko Gagal Ginjal Kronik di Unit Hemodialisis RSUD Wates Kulon Progo. *Majalah Farmaseutik*, *11*(2), 316–320.
- Prater, K. E., Hosanagar, A., Klumpp, H., Angstadt, M., & Phan, K. L. (2013). Aberrant Amygdala–Frontal Cortex Connectivity During Perception of

- Fearful Faces and at Rest in Generalized Social Anxiety Disorder. *Depression and Anxiety*, 30(3), 234–241. <https://doi.org/10.1002/da.22014>
- Preljevic, V. T., Østhus, T. B. H., Sandvik, L., Opjordsmoen, S., Nordhus, I. H., Os, I., & Dammen, T. (2012). Screening for anxiety and depression in dialysis patients: Comparison of the Hospital Anxiety and Depression Scale and the Beck Depression Inventory. *Journal of Psychosomatic Research*, 73(2), 139–144. <https://doi.org/10.1016/j.jpsychores.2012.04.015>
- Price, S. A., & Wilson, L. M. (2006). *Pathofisiologi Konsep Klinik Proses-Proses Penyakit*. Jakarta: EGC.
- Puspitosari, W. A. (2009). Terapi Kognitif dan Perilaku pada Gangguan Obsesif Kompulsif. *Mutiara Medika*, 9(2), 7.
- Ramadhan, H. F., Hermansyah, Y., Wisudanti, D. D., & Suryono, S. (2018). Hemodialysis Effect on Systolic Left Ventricular Function in Stage V Chronic Kidney Disease Patients. *Journal of Agromedicine and Medical Sciences*, 4(3), 128. <https://doi.org/10.19184/ams.v4i3.6543>
- Rector, N., A. (2010). *Cognitive-Behavioural Therapy An Information Guide*. Centre for Addiction and Mental Health.
- Reza, I. F. (2016). Implementasi Coping Religious dalam Mengatasi Gangguan Fisik-Psikis-Sosial-Spiritual pada Pasien Gagal Ginjal Kronik. *Intizar*, 22(2), 243. <https://doi.org/10.19109/intizar.v22i2.940>
- Ruiz de Alegría Fernández de Retana, B., Ruiz de Alegría-Fernández de Retana, B., Basabe Barañano, N., Basabe-Barañano, N., Saracho Rotaache, R., & Saracho-Rotaache, R. (2013). Coping mechanisms as a predictor for quality of life in patients on dialysis: A longitudinal and multi-centre study. *Nefrología (English Edition)*, 33(3), 342–354. <https://doi.org/10.3265/Nefrologia.pre2013.Feb.11771>
- Sarah, & Hasanat, N., UI. (2010). Kajian Teoritis Pengaruh ARt Therapy dalam Mengurangi Kecemasan pada Penderita Kanker. *Buletin Psikologi*, 18(1), 29–35.
- Schena, F. P. (2011). Management of patients with chronic kidney disease. *Internal and Emergency Medicine*, 6 Suppl 1, 77–83. <https://doi.org/10.1007/s11739-011-0688-2>
- Semaan, V., Nouredine, S., & Farhood, L. (2018). Prevalence of depression and anxiety in end-stage renal disease: A survey of patients undergoing hemodialysis. *Applied Nursing Research*, 43, 80–85. <https://doi.org/10.1016/j.apnr.2018.07.009>
- Setyaningsih, T., Mustikasari, M., & Nuraini, T. (2011). Peningkatkan Harga Diri Pada Klien Gagal Ginjal Kronik Melalui Cognitive Behavior Therapy (CBT). *Jurnal Keperawatan Indonesia*, 14(3), 165–170. <https://doi.org/10.7454/jki.v14i3.63>
- Sohn, B. K., Oh, Y. K., Choi, J.-S., Song, J., Lim, A., Lee, J. P., An, J. N., Choi, H.-J., Hwang, J. Y., Jung, H.-Y., Lee, J.-Y., & Lim, C. S. (2018). Effectiveness of group cognitive behavioral therapy with mindfulness in end-stage renal disease hemodialysis patients. *Kidney Research and Clinical Practice*, 37(1), 77–84. <https://doi.org/10.23876/j.krcp.2018.37.1.77>

- Sompie, E. M., Kaunang, T. M. D., & Munayang, H. (2015). Hubungan antara Lama Menjalani Hemodialisis dengan Depresi pada Pasien dengan Penyakit Ginjal Kronik di RSUP. Prof. Dr. R. D. Kandou Manado. *Jurnal E-Clinic (ECL)*, 3(1). <https://doi.org/10.35790/ecl.3.1.2015.6834>
- Sopha, R. F., Wardhani, I. Y., & Fakultas Ilmu Keperawatan Universitas Indonesia, Depok - Indonesia. (2016). Stres dan Tingkat Kecemasan saat Ditetapkan Perlu Hemodialisis Berhubungan dengan Karakteristik Pasien. *Jurnal Keperawatan Indonesia*, 19(1), 55–61. <https://doi.org/10.7454/jki.v19i1.431>
- Stallard, P. (2002). *Think good—Feel good: A cognitive behavior therapy workbook for children and young people*. Chicester: John Wiley & Sons.
- Stuart, G. W. (2006). *Buku Saku Keperawatan Jiwa*. Alih Bahasa: Ramona P. Kapoh & Egi Komara Yudha. Jakarta: EGC.
- Sudoyo, A. W., Sutiyohadi, B., Alwi, I., Simadibrata, M., & Setiati, S. (2007). *Buku ajar Ilmu Penyakit Dalam (IV)*. Jakarta: Pusat Penerbitan Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Indonesia.
- Sulistini, R., Damanik, H. DL., & Azinora, D. V. (2018). Kualitas Hidup Pasien yang Menjalani Hemodialisis. *Proceeding Seminar Nasional Keperawatan "tren Perawatan Paliatif sebagai Peluang Praktik Keperawatan Mandiri"*. <http://www.conference.unsri.ac.id/index.php/SNK/article/view/1212/609>
- Sumah, D. F. (2020). Kecerdasan Spiritual Berkorelasi dengan Tingkat Kecemasan Pasien Gagal Ginjal Kronik yang Menjalani Hemodialisa di Ruang Hemodialisa RSUD dr. M. Haulussy Ambon. *Jurnal Biosainstek*, 2(1), 87–92.
- Suseł, J., Batycka-Baran, A., Reich, A., & Szepletowski, J. (2014). Uraemic Pruritus Markedly Affects the Quality of Life and Depressive Symptoms in Haemodialysis Patients with End-stage Renal Disease. *Acta Dermato Venereologica*, 94(3), 276–281. <https://doi.org/10.2340/00015555-1749>
- Syed-Ahmed, M., & Narayanan, M. (2019). Immune Dysfunction and Risk of Infection in Chronic Kidney Disease. *Advances in Chronic Kidney Disease*, 26(1), 8–15. <https://doi.org/10.1053/j.ackd.2019.01.004>
- Terkawi, A. S., Tsang, S., AlKahtani, G. J., Al-Mousa, S. H., Al Musaed, S., AlZoraigi, U. S., Alasfar, E. M., Doais, K. S., Abdulrahman, A., & Altirkawi, K. A. (2017). Development and validation of Arabic version of the Hospital Anxiety and Depression Scale. *Saudi Journal of Anaesthesia*, 11(Suppl 1), S11–S18. https://doi.org/10.4103/sja.SJA_43_17
- Theofilou, P. (2011). Depression and Anxiety in Patients with Chronic Renal Failure: The Effect of Sociodemographic Characteristics. *International Journal of Nephrology*, 2011. <https://doi.org/10.4061/2011/514070>
- Tolin, D. F. (2010). Is cognitive–behavioral therapy more effective than other therapies? A meta-analytic review. *Clinical Psychology Review*, 30(6), 710–720. <https://doi.org/10.1016/j.cpr.2010.05.003>
- Turkistani, I., Nuqali, A., Badawi, M., Taibah, O., Alserihy, O., Morad, M., & Kalantan, E. (2014). The prevalence of anxiety and depression among end-stage renal disease patients on hemodialysis in Saudi Arabia. *Renal Failure*, 36(10), 1510–1515. <https://doi.org/10.3109/0886022X.2014.949761>

- USRDS. (2015). *USRDS Annual Data Report Volume 2: Epidemiology of Kidney Disease in the United States*. United States Renal Data System 2.
- Valsaraj, B. P. (2016). Cognitive Behaviour Therapy for Anxiety and Depression among People Undergoing Haemodialysis: A Randomized Control Trial. *Journal of Clinical and Diagnostic Research*. <https://doi.org/10.7860/JCDR/2016/18959.8383>
- Vasilopoulou, C., Bourtsi, E., Giaple, S., Koutelekos, I., Theofilou, P., & Polikandrioti, M. (2015). The Impact of Anxiety and Depression on the Quality of Life of Hemodialysis Patients. *Global Journal of Health Science*, 8(1), 45. <https://doi.org/10.5539/gjhs.v8n1p45>
- Wahyuni, A., Kartika, I. R., & Asrul, I. F. (2019). Korelasi Lama Hemodialisa dengan Fungsi Kognitif. *Real in Nursing Journal*, 2(1), 1. <https://doi.org/10.32883/rnj.v2i1.328>
- Wang, R., Liao, C., He, H., Hu, C., Wei, Z., Hong, Z., Zhang, C., Liao, M., & Shui, H. (2020). COVID-19 in Hemodialysis Patients: A Report of 5 Cases. *American Journal of Kidney Diseases*, 76(1), 141–143. <https://doi.org/10.1053/j.ajkd.2020.03.009>
- Wibisono, D. (2014). *Deteksi Dini Menjaga Kualitas dan Fungsi Ginjal* (11th ed.). Majalah Rumah Sakit Mitra Keluarga.
- Wright, J., H., Brown, G., K., Thase, M., E., & Basco, M., Ramirez. (2017). *Learning Cognitive-Behavior Therapy An Illustrated Guide* (Second). American Psychiatric Association.
- Wurara, Y. G. V., Kanine, E., & Wowiling, F. (2013). Mekanisme Koping pada Pasien Penyakit Ginjal Kronik yang Menjalani Terapi Hemodialisis di Rumah Sakit Prof. Dr.R.D Kandou Manado. *E Journal Keperawatan (e-Kp)*, 1(1), 7.
- Yang, Y., Ding, R., Hu, D., Zhang, F., & Sheng, L. (2014). Reliability and validity of a Chinese version of the HADS for screening depression and anxiety in psycho-cardiological outpatients. *Comprehensive Psychiatry*, 55(1), 215–220. <https://doi.org/10.1016/j.comppsy.2013.08.012>
- Yanti, E. K. Y., & Miswadi. (2018). Faktor-faktor yang Berhubungan dengan Kecemasan pada Pasien Gagal Ginjal Kronik yang Akan Menjalani Terapi Hemodialisis di Ruang Hemodialisis RSUD Bengkalis Tahun 2016. *Jurnal Ners Universitas Pahlawan*, 2(1), 28–40.
- Zhang, M., Kim, J. C., Li, Y., Shapiro, B. B., Porszasz, J., Bross, R., Feroze, U., Upreti, R., Martin, D., Kalantar-Zadeh, K., & Kopple, J. D. (2014). Relation Between Anxiety, Depression, and Physical Activity and Performance in Maintenance Hemodialysis Patients. *Journal of Renal Nutrition*, 24(4), 252–260. <https://doi.org/10.1053/j.jrn.2014.03.002>
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., Niu, P., Zhan, F., Ma, X., Wang, D., Xu, W., Wu, G., Gao, G. F., & Tan, W. (2020). A Novel Coronavirus from Patients with Pneumonia in China, 2019. *New England Journal of Medicine*, 382(8), 727–733. <https://doi.org/10.1056/NEJMoa2001017>
- Zigmond, A. S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67(6), 361–370. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x>

Zurmeli, Bayhakki, & Utami, G. T. (2015). Hubungan Dukungan Keluarga dengan Kualitas Hidup Pasien Gagal Ginjal Kronik yang Menjalani Terapi Hemodialisis di RSUD Arifin Achmad Pekanbaru. *Jurnal Online Mahasiswa*, 2(1), 670–681.