

## DAFTAR PUSTAKA

- Ajenifuja, K. O., Oyetunji, I. O., Orji, E. O., Adepiti, C. A., Loto, O. M., Tijani, M. A., & Dare, F. O. (2013). Post-partum urinary retention in a teaching hospital in southwestern Nigeria. *Journal of Obstetrics and Gynaecology Research*, 39(8), 1308–1313. <https://doi.org/10.1111/jog.12069>
- Anugerah, I., Iswari, W. A., Pardede, T. U., & Darus, F. (2017). Tatalaksana Retensio Urin Pascapersalinan. *CDK-255*, 44(8), 531–536.
- Bayu, B., Lengkong, R., & Wantania, J. (2019). Postpartum Urinary Retention after Vaginal Delivery. *Indones J Obstet Gynecol*, 7(2), 141–145.
- Beaumont, T. (2019). Prevalence and outcome of postpartum urinary retention at an Australian hospital. *Midwifery*, 70, 92–99. <https://doi.org/10.1016/j.midw.2018.12.013>
- Buchanan, J., & Beckmann, M. (2014). Postpartum voiding dysfunction : Identifying the risk factors. *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 54, 41–45. <https://doi.org/10.1111/ajo.12130>
- Banyumas, RSUD. (2019). Pelayanan Obstetrik Neonatal Emergensi Konprehensif (PONEK) RSUD Banyumas diakses dari <http://rsudbms.banyumaskab.go.id/news/29836/pelayanan-obstetrik-neonatal-emergensi-komprehensif-ponek-rsud-banyumas>
- Butcher, H.K., Bulechek, G.M., Dochterman, J.M., Wagner, C.M (2018) Nursing Intervention Classification (NIC) edisi ketujuh terjemah oleh Intansari Nurjannah. Yogyakarta: Mocomedia.
- Cavkaytar, S., Kokanalı, M. K., Baylas, A., Topçu, H. O., Laleli, B., & Taşçı, Y. (2014). Postpartum urinary retention after vaginal delivery : Assessment of risk factors in a case-control study. *J Turk Ger Gynecol Assoc*, 15, 140–143. <https://doi.org/10.5152/jtgga.2014.13102>
- Chai, A. H., Wong, T., & Mak, H. J. (2008). Prevalence and associated risk factors of retention of urine after caesarean section. *Int Urogynecol J*, 19, 537–542. <https://doi.org/10.1007/s00192-007-0470-1>
- Chaliha, C. (2006). Postpartum bladder dysfunction. *Gynaecological and Perinatal Practice*, 6, 133–139. <https://doi.org/10.1016/j.rigapp.2006.05.007>

- Choe, W. S., Ng, K., Atan, I. K., & Lim, P. S. (2018). Acceptable Postvoid Residual Urine Volume after Vaginal Delivery and Its Association with Various Obstetric Parameters. *Obstetrics and Gynecology International*, 2018, 7. <https://doi.org/doi.org/10.1155/2018/5971795>
- Cunningham, F. G., Leveno, K. J., Bloom, S. L., Hauth, J. C., Rouse, D. J., Spong, C. Y. (2012). *Obstetri Williams* (23rd ed.). EGC.
- Djusad, S. (2020). Manajemen Retensio Urin Pasca Persalinan Pervaginam. *EJournal Kedokteran Indonesia*, 8(1), 2–8. <https://doi.org/10.23886/ejki.8.11690>.
- Drake, R.L., Vogl, A. W., Mitchell, A. W. (2014). *Gray Dasar-Dasar Anatomi*. Singapura: Elsevier.
- Emilia, F.W., Yuliana, F., Noviantina, R. (2017). The correlation between episiotomy, perineal laceration and type of labor toward the incidence of postpartum urinary retensio at dr. H. Soemarno Sosroatmodjo Hospital of Kuala Kapuas in 2015-2016. *Advances in Health Science Research*, 6, 558–565.
- Ganong, W. F. (2008). *Buku Ajar Fisiologi Kedokteran edisi 22*. Jakarta: EGC.
- Gaudernack, L. C., Frøslie, K. F., Michelsen, T. M., Voldner, N., & Lukasse, M. (2018). De-medicalization of birth by reducing the use of oxytocin for augmentation among first-time mothers – a prospective intervention study. *BMC Pregnancy and Childbirth*, 1–9. <https://doi.org/doi.org/10.1186/s12884-018-1706-4>
- Glavind, K., & Bj, J. (2003). Incidence and treatment of urinary retention postpartum. *Int Urogynecol J*, 14, 119–121. <https://doi.org/10.1007/s00192-002-1014-3>
- Hall, J. E. (2009). *Buku Saku Fisiologi Kedokteran Gayton & Hall edisi 11*. Jakarta: EGC.
- Hanretty, K.P. (2014). *Ilustrasi Obstetri (edisi ketujuh)*. Churchill Livingston Elsevier.
- Herdman, T.H., Kamitsuru, S (2017) *NANDA International Nursing Diagnoses: Definitions and Classification 2018-2020 eleventh Edition*. New York: Medical Publisher
- Hidalgo, R., Adair, S., Avondstondt, A., & Salamon, C. (2018). Intrapartum Risk Factors for Overt Postpartum Urinary Retention. *Journal of Pelvic Medicine and Surgery, October*, 1–2. <https://doi.org/10.13140/RG.2.2.18877.72169>
- Humburg, J., Troeger, C., Holzgreve, W., & Hoesli, I. (2011). Risk factors in prolonged

- postpartum urinary retention : an analysis of six cases. *Arch Gynecol Obstet*, 283, 179–183. <https://doi.org/10.1007/s00404-009-1320-9>
- Johansson, R., Malmvall, B., Andersson-ga, B., Larsson, B., Erlandsson, I., & Christensson, L. (2012). Guidelines for preventing urinary retention and bladder damage during hospital care. *Journal of Clinical Nursing*, 22, 347–355. <https://doi.org/10.1111/j.1365-2702.2012.04229.x>
- K. Lamb; R. Sanders. (2016). Bladder care in the context of motherhood : Ensuring holistic midwifery practice. *British Journal of Midwifery*, 24(6), 415–421.
- Kekre, A. N., Vijayanand, S., Dasgupta, R., & Kekre, N. (2011). Postpartum urinary retention after vaginal delivery. *International Journal of Gynecology and Obstetrics*, 112(2), 112–115. <https://doi.org/10.1016/j.ijgo.2010.08.014>
- Kemkes Republik Indonesia. (2013). *Buku Saku Pelayanan Kesehatan Ibu di Fasilitas Kesehatan Dasar dan Rujukan* (pertama).
- KMKP. (2018). Kumpulan Standar Prosedur Operasional RSUD Banyumas,
- Kurniarum, A. (2016). Asuhan Kebidanan Persalinan Dan Bayi Baru Lahir (I). Jakarta. Kemenkes RI.
- Lim, J. L. (2010). Post-partum voiding dysfunction and urinary retention. *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 50(April), 502–505. <https://doi.org/10.1111/j.1479-828X.2010.01237.x>
- Lowdermilk, D.L., Perry, S.E., Cashion, K. (2014). *Maternity Nursing* (8 th edition). Mosby Elsevier
- Mehta, S., & Anger, J. (2012). Evaluation and Management of Postpartum Urinary Retention. *Curr Bladder Dysfunct Rep*, 7, 260–263. <https://doi.org/10.1007/s11884-012-0162-7>
- Mulder, F. E. M., Hakvoort, R. A., de Bruin, J. P., Janszen, E. W., van der Post, J. A. M., & Roovers, J. P. W. R. (2018). Long-term micturition problems of asymptomatic postpartum urinary retention: a prospective case–control study. *International Urogynecology Journal*, 29(4), 481–488. <https://doi.org/10.1007/s00192-017-3457-6>
- Mulder, F. E. M., Rengerink, K. O., & Post, J. A. M. Van Der. (2016). Delivery-related risk factors for covert postpartum urinary retention after vaginal delivery. *Int Urogynecol J*, 27, 55–60. <https://doi.org/10.1007/s00192-015-2768-8>

- Mulder FE, Hakvoort RA, Schoffemeer MA, Limpens J, Van der Post JA, R. J. (2014). Postpartum urinary retention : a systematic review of adverse effects and management. *Int Urogynecol J*, 25, 1605–1612. <https://doi.org/10.1007/s00192-014-2418-6>
- Notoatmodjo, S. (2014). Metodologi Penelitian Kesehatan. Jakarta: Rineka Cipta.
- Oh, J. J., Kim, S. H., Shin, J. S., & Shin, S. J. (2015). Risk factors for acute postpartum urinary retention after vaginal delivery : focus on episiotomy direction. *The Journal of Maternal-Fetal & Neonatal Medicine*, 00(00), 1–4. <https://doi.org/10.3109/14767058.2014.1002395>
- Pribakti, (2011). Dasar-dasar Uroginekologi. Jakarta: Sagung Seto.
- Prawirohardjo, S. (2010). Ilmu Kebidanan edisi 4. Jakarta: P.T Bina Pustaka Sarwono Prawirohardjo.
- Petrana, N. H., Emilia, O., & Pradjatmo, H. (2016). Perbandingan kejadian retensi urin antara persalinan dengan vakum ekstraksi dan persalinan normal. *Jurnal Kesehatan Reproduksi*, 3(3), 188–193.
- Pifarotti, P., Gargasole, C., Folcini, C., Gattei, U., Nieddu, E., Sofi, G., Buonaguidi, A., & Meschia, M. (2014). Acute post-partum urinary retention : analysis of risk factors , a case – control study. *Arch Gynecol Obstet*, 289, 1249–1253. <https://doi.org/10.1007/s00404-014-3144-5>
- Pulatoğlu, Ç., Doğan, O., & Kılıççı, Ç. (2018). Postpartum urinary retention : Evaluation of risk factors Postpartum üriner retansiyon : Risk faktörlerinin değerlendirilmesi. *Turk J Obstet Gynecol*, 15, 70–74. <https://doi.org/10.4274/tjod.43931>
- Rahmanou, P., Caudwell-Hall, J., Kamisan Atan, I., & Dietz, H. P. (2016). The association between maternal age at first delivery and risk of obstetric trauma. *American Journal of Obstetrics and Gynecology*, 215(4), 451.e1-451.e7. <https://doi.org/10.1016/j.ajog.2016.04.032>
- Rarung, M. (2008). Perbandingan Pemasangan Kateter Menetap selama 12 dan 24 jam Pasca Seksio Sesarea pada Pencegahan Retensio Urin dan Resiko Infeksi Saluran Kemih. *JKM.*, Vol. 8(No. 1), 45–52.
- Richens, Y. (2016). Urinary catheterisation: Indications and complications. *British Journal of Midwifery*, vol 24(No 3), 164–168.

- Rizvi, R. M., & Rizvi, J. (2006). Management of postpartum urinary retention. *Gynaecological and Perinatal Practice*, 6, 140–144. <https://doi.org/10.1016/j.rigapp.2006.02.003>
- Rogers, R. G., & Leeman, L. L. (2007). Postpartum Genitourinary Changes. *Urol Clin N Am*, 34(2007), 13–21. <https://doi.org/10.1016/j.ucl.2006.10.005>
- Sauer, M. V. (2015). Reproduction at an advanced maternal age and maternal health. *Fertility and Sterility*, 103(5), 1136–1143. <https://doi.org/10.1016/j.fertnstert.2015.03.004>
- Sastroasmoro, S., Ismael, S. (2016). Dasar-dasar Metodologi Penelitian Klinis. Jakarta: Sagung Seto.
- Siswosudarmo, R., Emilia, O., (2008). Obstetri Fisiologi. Yogyakarta: Pustaka Cendekia.
- Sofian, A., (2013). Rustam Mochtar Sinopsis Obstetri - Obstetri Fisiologi Obstetri Patologi. Jakarta: EGC.
- Teo, R., Punter, J., Abrams, K., Mayne, C., & Tincello, D. (2007). Clinically overt postpartum urinary retention after vaginal delivery : a retrospective case-control study. *Int Urogynecol J*, 18, 521–524. <https://doi.org/10.1007/s00192-006-0183-x>
- Tiberon, A., Carbonnel, M., Vidart, A., Halima, M. Ben, Deffieux, X., & Ayoubi, J. (2018). Risk factors and management of persistent postpartum urinary retention. *Journal of Gynecology Obstetrics and Human Reproduction*, 47(9), 437–441. <https://doi.org/10.1016/j.jogoh.2018.08.002>
- Youssef, A., Salsi, G., Cataneo, I., Pacella, G., & Azzarone, C. (2019). Fundal pressure in second stage of labor ( Kristeller maneuver ) is associated with increased risk of levator ani muscle avulsion. *Ultrasound Obstet Gynecol*, 53, 95–100. <https://doi.org/10.1002/uog.19085>