

DAFTAR PUSTAKA

- Alemu, A., Abageda, M., Assefa, B. & Melaku, G. (2019) Low birth weight: prevalence and associated factors among newborns at hospitals in Kambata-Tembaro zone, southern Ethiopia 2018. *The Pan African Medical Journal*, 34.
- Bahrami, H. R., Mosa Farkhani. E., Beygi. B., Gholian-Aval. M., and, T. A. & Hoseini. S.J. (2020) Risk Factors of Low Birth Weight Infants: A Population-Based Cross-Sectional Study. *International Journal of Pediatrics.*, 8(1)8.
- Banerjee, A., Singh, A. K. & Chaurasia, H. (2020) An exploratory spatial analysis of low birth weight and its determinants in India. *Clinical Epidemiology and Global Health*.
- Barbuscia, A., Martikainen, P., Myrskylä, M., Remes, H., Somigliana, E., Klemetti, R. & Goisis, A. (2020) Maternal age and risk of low birth weight and premature birth in children conceived through medically assisted reproduction. Evidence from Finnish population registers. *Human Reproduction*, 35(1): 212-220.
- Bekele, A., Seyoum, G., Tesfaye, K. & Fantahun, Y. (2019) The effects of maternal age and parity on the birth weight of newborns among mothers with singleton pregnancies and at term deliveries. *Ethiopian Journal of Health Development*, 33(3).
- BPS (2017) Laporan Survei Demografi Kesehatan Indonesia 2017. Jakarta: Kementerian Kesehatan RI.
- Darwis, A., Abdullah, A., Adamy, A. & Septiani, R. (Year) The Relationship Between Service Quality Antenatal Care and Low Birth Weight in Indonesia: IDHS in 2017. In: 4th International Symposium on Health Research (ISHR 2019), 2020. Atlantis Press, 337-340.
- Desta, M., Tadese, M., Kassie, B. & Gedefaw, M. (2019) Determinants and adverse perinatal outcomes of low birth weight newborns delivered in Hawassa University Comprehensive Specialized Hospital, Ethiopia: a cohort study. *BMC research notes*, 12(1): 118.
- Eshete, A., Alemu, A. & Zerfu, T. A. (2019) Magnitude and Risk of Dying among Low Birth Weight Neonates in Rural Ethiopia: A Community-Based Cross-Sectional Study. *International journal of pediatrics*, 2019.
- Figueiredo, A. C. M. G., Gomes-Filho, I. S., Batista, J. E. T., Orrico, G. S., Porto, E. C. L., Pimenta, R. M. C., dos Santos Conceição, S., Brito, S. M., Ramos, M. d. S. X. & Sena, M. C. F. (2019) Maternal anemia and birth weight: A prospective cohort study. *PloS one*, 14(3): e0212817.
- Gebresilassie, B., Belete, T., Tilahun, W., Berhane, B. & Gebresilassie, S. (2019) Timing of first antenatal care attendance and associated factors among pregnant women in public health institutions of Axum town, Tigray,

- Ethiopia, 2017: a mixed design study. *BMC pregnancy and childbirth*, 19(1): 340.
- Handayani, T. R. (2018) Hubungan Paparan Asap Rokok Dan Konsumsi Kafein Terhadap Berat Badan Bayi Lahir. *Babul Ilmi Jurnal Ilmiah Multi Science Kesehatan*, 8.
- Hartini, L. & Salam, M. N. (Year) Age Is A High Risk Of Low Birth Weight In The Working Area Of Seluma District. In: 1st International Conference on Inter-Professional Health Collaboration (ICIHC 2018), 2019. Atlantis Press.
- Helliyana, H., Arintonang, E. Y. & Sanusi, S. R. (2019) The Associations between Maternal Education, Chronic Energy Deficit, and Anemia in Pregnant Women: An Evidence from Lhokseumawe, Indonesia. *Journal of Maternal and Child Health*, 4(5): 302-306.
- Huang, I., Mak, D., Cheung, P., Abraham, M., Clemens, T. & Turner, S. (2019) A systematic review of associations between maternal exposures during pregnancy other than smoking and antenatal fetal measurements. *Environmental research*, 173528-538.
- Indriyani, D. (2013) *Keperawatan Maternitas Pada Area Perawatan Antenatal*, Yogyakarta:Graha Ilmu.
- Kabir, A., Rashid, M. M., Hossain, K., Khan, A., Sikder, S. S. & Gidding, H. F. (2020) Women's empowerment is associated with maternal nutrition and low birth weight: evidence from Bangladesh Demographic Health Survey. *BMC Women's Health*, 201-12.
- Kemenkes (2015) RENCANA STRATEGIS KEMENTERIAN KESEHATAN 2015-2019 Jakarta.
- Khan, N., Mozumdar, A. & Kaur, S. (2020) Determinants of low birth weight in India: an investigation from the national family health survey. *American Journal of Human Biology*, 32(3): e23355.
- Khatun, S. & Rahman, M. (2008) Quality of antenatal care and its dose–response relationship with birth weight in a maternal and child health training institute in Bangladesh. *Journal of biosocial science*, 40(3): 321-337.
- Koren, G., Ornoy, A. & Berkovitch, M. (2018) Hyperemesis gravidarum—Is it a cause of abnormal fetal brain development? *Reproductive Toxicology*, 7984-88.
- Kramer, M. S. (1987) Determinants of low birth weight: methodological assessment and meta-analysis. *Bulletin of the world health organization*, 65(5): 663.
- Lujan, L. M. B., Santana, P. S., García, L. E. G., Hernández, J. A. G. & Majem, L. S. (2015) Sociodemographic and health characteristics associated with low birth weight in the canary islands. *Nutricion hospitalaria*, 32(4): 1541-1547.
- Maryunani, A. P., E (2013) *Asuhan Bayi Dengan Berat Badan Lahir Rendah*, Jakarta:Trans Info Medika.
- Noelke, C., Chen, Y.-H., Osypuk, T. L. & Acevedo-Garcia, D. (2019) Economic Downturns and Inequities in Birth Outcomes: Evidence From 149 Million US Births. *American journal of epidemiology*, 188(6): 1092-1100.

- Nyamasege, C., Kimani-Murage, E., Wanjohi, M., Kaindi, D., Ma, E., Fukushige, M. & Wagatsuma, Y. (2019) Determinants of low birth weight in the context of maternal nutrition education in urban informal settlements, Kenya. *Journal of developmental origins of health and disease*, 10(2): 237-245.
- Olusanya, B. O. & Ofovwe, G. E. (2010) Predictors of preterm births and low birthweight in an inner-city hospital in sub-Saharan Africa. *Maternal and child health journal*, 14(6): 978-986.
- Pantiawati (2010) *Bayi dengan Berat Badan Lahir Rendah*, Jakarta: Nuha Offset.
- Paul, P., Zaveri, A. & Chouhan, P. (2019) Assessing the impact of antenatal care utilization on low birthweight in India: Analysis of the 2015–2016 National Family Health Survey. *Children and Youth Services Review*, 106104459.
- Petry, C. J., Ong, K. K., Beardsall, K., Hughes, I. A., Acerini, C. L. & Dunger, D. B. (2018) Vomiting in pregnancy is associated with a higher risk of low birth weight: a cohort study. *BMC pregnancy and childbirth*, 18(1): 133.
- Pinzón-Rondón, Á. M., Gutiérrez-Pinzon, V., Madriñan-Navia, H., Amin, J., Aguilera-Otalvaro, P. & Hoyos-Martínez, A. (2015) Low birth weight and prenatal care in Colombia: a cross-sectional study. *BMC pregnancy and childbirth*, 15(1): 118.
- Prawirohardjo, S. (2009) *Buku Acuan Nasional Pelayanan Kesehatan Maternal dan Neonatal*, Jakarta: Bina Pustaka.
- Prawirohardjo, S. (2011) *Ilmu Kebidanan*, Jakarta: Bina Pustaka.
- Proverawati, A. & Ismawati, C. (2010) *Berat Badan Lahir Rendah (BBLR)*, Yogyakarta: Nuha Medika.
- Purwaningsih, W. & Fatmawati, S. (2010) *Asuhan Keperawatan Maternitas*, Yogyakarta: Nuha Medika.
- Rahmati, S., Azami, M., Badfar, G., Parizad, N. & Sayehmiri, K. (2019) The relationship between maternal anemia during pregnancy with preterm birth: a systematic review and meta-analysis. *The Journal of Maternal-Fetal & Neonatal Medicine*, 1-11.
- Ramadani, M. & Utomo, B. (2019) Prenatal Tobacco Exposure and Neonate Birth Weight. *Indian Journal of Public Health Research & Development*, 10(1).
- Ramraj, C., Pulver, A., O'Campo, P., Urquia, M. L., Hildebrand, V. & Siddiqi, A. (2020) A Scoping Review of Socioeconomic Inequalities in Distributions of Birth Outcomes: Through a Conceptual and Methodological Lens. *Maternal and Child Health Journal*, 1-9.
- Servan, M. E., Sosa, R. S. G., Najera, L. E. & Darney, B. G. (2016) Timeliness, frequency and content of antenatal care: which is most important to reducing indigenous disparities in birth weight in Mexico? *Health policy and planning*, 31(4): 444-453.
- Sukarni & Wahyu (2013) *Buku Ajar Keperawatan Maternitas*, Yogyakarta: Nuha Medika.
- Tabrizi, J. S., Asghari, A., Pourali, F., Kousha, H. & Nikniaz, L. (2019) Effects of Food Supplementation During Pregnancy on Maternal Weight Gain,

- Hemoglobin Levels and Pregnancy Outcomes in Iran. *Maternal and child health journal*, 23(2): 258-264.
- Tafere, T. E., Afework, M. F. & Yalew, A. W. (2018) Providers adherence to essential contents of antenatal care services increases birth weight in Bahir Dar City Administration, north West Ethiopia: a prospective follow up study. *Reproductive health*, 15(1): 163.
- Takito, M. Y. & Benício, M. H. D. A. (2010) Physical activity during pregnancy and fetal outcomes: a case-control study. *Revista de saude publica*, 44:90-101.
- Tiran, D. (2009) *Mual dan Muntah Kehamilan*, Yogyakarta:EGCUGM.
- UNICEF (2015) *Nearly 15 per cent of babies worldwide are born with low birthweight* [Online]. Available: <https://data.unicef.org/topic/nutrition/low-birthweight/> [Accessed 20 November 2019].
- UNICEF. (2019) Monitoring the situation of children and women: There are wide disparities in antenatal care across countries.
- WHO (2013) *Guideline: Calcium supplementation in pregnant women*:World Health Organization.
- WHO (2014) Global Nutrition Targets 2025: Low birth weight policy brief. World Health Organization.
- WHO (2016a) Standards for improving quality of maternal and newborn care in health facilities.
- WHO (2016b) *WHO recommendations on antenatal care for a positive pregnancy experience*.
- Zhou, H., Wang, A., Huang, X., Guo, S., Yang, Y., Martin, K., Tian, X., Josephs-Spaulding, J., Ma, C. & Scherpbier, R. W. (2019) Quality antenatal care protects against low birth weight in 42 poor counties of Western China. *PloS one*, 14(1): e0210393.