

Daftar Pustaka

- Abdolahi, F.S., Babaei, H., Zarringhalam, M., Ashrafi, J. (2015) 'Wound healing effect of topical phenytoin on rat palatal mucosa', *Int J Curr Res Aca Rev.* 3(6). pp. 463-76.
- Albertsmeier, M. *et al.* (2012) 'Evaluation of the safety and efficacy of MonoMax® suture material for abdominal wall closure after primary midline laparotomy - A controlled prospective multicentre trial: ISSAAC [NCT005725079]', *Langenbeck's Archives of Surgery*, 397(3), pp. 363–371. doi: 10.1007/s00423-011-0884-6.
- Barrientos, S., Stojadinovic, O., Golinko, M.S., Brem, H., Tomic-Canic, M. (2008) 'Growth factors and cytokines in wound healing', *Wound Repair Regen.* 16(5). pp. 585-601. doi: 10.1111/j.1524-475X.2008.00410.x. [PubMed: 19128254].
- Bevis, P. M., Windhaber, R. A., Lear, P. A., Poskitt, K. R., Earnshaw, J. J., Mitchell, D. C. (2010) 'Randomized clinical trial of mesh versus sutured wound closure after open abdominal aortic aneurysm surgery', *Br J Surg.* 97. pp. 1497-1502.
- Bloemen, A., van Dooren, P., Huizinga, B.F., Hoofwijk, A.G., (2011) Randomized clinical trial comparing polypropylene or polydioxanone for midline abdominal wall closure. *Br J Surg*, 98(5):633-639.
- Bosanquet, D. C. *et al.* (2015) 'Systematic review and meta-regression of factors affecting midline Incisional hernia rates: Analysis of 14 618 Patients', *PLoS ONE*, 10(9), pp. 1–18. doi: 10.1371/journal.pone.0138745.
- Brown, R.L., Ormsby, I., Doetschman, T.C., Greenhalgh, D.G. (1995) 'Wound healing in transforming growth factor-beta-deficient mouse', *Wound Repair Regen.* 3(1). pp. 25-36. doi: 10.1046/j.1524-475X.1995.30108.x. [PubMed: 17168860].
- Cengiz, Y., Blomquist, P., Israelsson, L.A., (2001). Small Tissue Bites and Wound Strength. *Arch Surg* (136). 272-275

- Cengiz, Y., Gislason, H., Svanes, K., & Israelsson, L. A. (2001). Mass Closure Technique : An Experimental Study on Separation of Wound Edge, 0–3.
- Ceydeli, A., Rucinski, J. and Wise, L. (2007) ‘Finding the best abdominal closure - An evidence-based overview of the literature’, *Recurrent Hernia: Prevention and Treatment*, pp. 117–122. doi: 10.1007/978-3-540-68988-1_14.
- Chalya, P. L. *et al.* (2015) ‘Abdominal fascia closure following elective midline laparotomy: A surgical experience at a tertiary care hospital in Tanzania’, *BMC Research Notes*. BioMed Central, 8(1), pp. 1–9. doi: 10.1186/s13104-015-1243-4.
- Corman, M. L., Veidenheimer, M. C., Coller, J. A. (1981) ‘Controlled clinical trial of three suture materials for abdominal wall closure after bowel operations’, *Am J Surg*, 141, pp. 510–513.
- D’Souza, R.. and Novell, R. (2013) ‘Laparotomy: Elective and Emergency’, in Novell, R., Baker, D. M., and Goddard, N. (eds) *Kirk’s General Surgical Operations*. Sixth Ed. Edinburgh: Churchill Livingstone Elsevier, pp. 38–56.
- Dahlan, S. (2011) ‘*Statistik Untuk Kedokteran dan Kesehatan*’, Edisi 5, Jakarta: Salemba Medika.
- Dart, A. J. and Dart, C. M. (2011) ‘Suture Material: Conventional and Stimuli Responsive’, *Comprehensive Biomaterials*, pp. 573–587. doi: 10.1016/B978-0-08-055294-1.00245-2.
- Deerenberg, E. B. *et al.* (2015) ‘Small bites versus large bites for closure of abdominal midline incisions (STITCH): A double-blind, multicentre, randomised controlled trial’, *The Lancet*. Elsevier Ltd, 386(10000), pp. 1254–1260. doi: 10.1016/S0140-6736(15)60459-7.
- Dovi, J.V., He, L.K., DiPietro, L.A. (2003) ‘Accelerated wound closure in neutrophil-depleted mice’, *J Leukoc Biol*. 73(4). pp. 448–55. doi: 10.1189/jlb.0802406. [PubMed: 12660219].
- Dubay, D. A. and Franz, M. G. (2003) ‘Acute wound healing: The biology of acute wound failure’, *Surgical Clinics of North America*, pp. 463–481. doi:

10.1016/S0039-6109(02)00196-2.

Dubay, D. A. *et al.* (2004) 'Fascial fibroblast kinetic activity is increased during abdominal wall repair compared to dermal fibroblasts', *Wound Repair and Regeneration*, 12(5), pp. 539–545. doi: 10.1111/j.1067-1927.2004.012506.x.

El Gazaerly, H., Elbardisey, D.M., Eltokhy, H.M., Teaama, D. (2013) 'Effect of transforming growth factor Beta 1 on wound healing in induced diabetic rats', *Int J Health Sci (Qassim)*. 7(2). pp. 160-72. doi: 10.12816/0006040. [PubMed: 24421745].

Faiz, O. and Moffat, D. (2002) *Anatomy at a Glance [e-Book]*. doi: 10.5005/jp/books/10050.

Falanga, V., (2003) *Mechanisms of Cutaneous Wound Repair*, Dalam: Freedberg, I.M., Wolff, K., Eisen, A.Z., et al, *Fitzpatrick's Dermatology In General Medicine*. Edisi ke-6. New York: Graw-Hill.

Fernandez, L. G. (2017) *Abdominal Closure Technique: Closure of Abdomen, Postoperative Care, Complications*, Medscape. Available at: <https://emedicine.medscape.com/article/1961789-technique>.

Fortelny, R. H. *et al.* (2015) 'Effect of suture technique on the occurrence of incisional hernia after elective midline abdominal wall closure: Study protocol for a randomized controlled trial', *Trials*, 16(1), pp. 1–8. doi: 10.1186/s13063-015-0572-x.

Franz, M. G. (2009) 'The Biology of Hernia Formation-2009', 88(1), pp. 1–16. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276402/>.

Franz, M. G. *et al.* (2001) 'Fasial incisions heal faster than skin: A new model of abdominal wall repair', *Surgery*, 129(2), pp. 203–208. doi: 10.1067/msy.2001.110220.

Harding, K.G., Morris, H.L., Patel, G.K. (2002) 'Healing Chronic Wounds', *BMJ*. 321. pp. 160-3.

Harlaar, J. J., van Ramshorst, G. H., Nieuwenhuizen, J., Ten Brinke, J. G., Hop W. C., Kleinrensink, G. J. et al. (2009), 'Small stitches with small suture

- distances increase laparotomy closure strength', *Am J Surg*, 198. pp. 392-395.
- Heinrich, P. C. *et al.* (2003) 'Principles of interleukin (IL)-6-type cytokine signalling and its regulation', *Biochemical Journal*, 374(1), pp. 1–20. doi: 10.1042/bj20030407.
- Hodgson, N. C. F., Malthaner, R. A. and Østbye, T. (2000) 'The search for an ideal method of abdominal fasial closure: A meta- analysis', *Annals of Surgery*, 231(3), pp. 436–442. doi: 10.1097/00000658-200003000-00018.
- Hodgson, N. C., Malthaner, R. A., Ostbye, T. (2000) 'The search for an ideal method of abdominal fascial closure : a meta-analysis', *Ann Surg*. 231. pp. 436-442.
- Hodgson, N. C., Malthaner, R. A., Ostbye, T. (2000) 'The search for an ideal method of abdominal fascial closure : a meta-analysis'. *Ann Surg*, 231.pp. 436-442.
- Höer, J. *et al.* (2002) 'Factors influencing the development of incisional hernia. A retrospective study of 2,983 laparotomy patients over a period of 10 years', *Der Chirurg*, 73(5), pp. 474–480. doi: 10.1007/s00104-002-0425-5.
- Hoer, J. J., Junge, K., Schachtrupp, A., Klinge, U., Schumpelick, V., (2002). Influence of laparotomy closure technique on collagen synthesis in the incisional region, *Hernia* (6) ;93–98. <https://doi.org/10.1007/s10029-002-0070-4>
- Hogstrom, H., Haglund, U., Zederfeldt, B., (1985). Suture Technique and Early Breaking Strength of Intestinal Anastomoses and Laparotomy Wounds. *Acta Chir Scand*. 151:441-443.
- Indrawan, I. and Dachlan, I. (2016) 'Perbandingan pengaruh aplikasi aloe vera, madu, saliva dan putih telur terhadap ekspresi interleukin-6 pada proses penyembuhan luka insisi kulit tikus', Universitas Gadjah Mada, pp. 1-14.
- ink, C. *et al.* (2014) 'Incisional hernia rate 3 years after midline laparotomy', *British Journal of Surgery*, 101(2), pp. 51–54. doi: 10.1002/bjs.9364.
- Israelsson, L. A. (1998), 'The surgeon as a risk factor for complications of midline incisions'. *Eur J Surg*, 164. pp. 353-359.

- Israelsson, L. A. and Millbourn, D. (2013) 'Prevention of incisional hernias. How to close a midline incision.', *Surgical Clinics of North America*, pp. 1027–1040. doi: 10.1016/j.suc.2013.06.009.
- Israelsson, L. A., Jonsson, T. (1993) 'Suture length to wound length ratio and healing of midline laparotomy incisions', *Br J Surg*, 80. pp.1284-1286.
- Israelsson, L. A., Jonsson, T. (1996) 'Incisional hernia after midline laparotomy : a prospective study', *Eur J Surg*, 162.pp. 125-129.
- Israelsson, L. A., Jonsson, T. (1997) 'Overweight and healing of midline incisions : the importance of suture technique', *Eur J Surg*, 163. pp. 175-180.
- Israelsson, L. A., Jonsson, T., Knutsson, A. (1996) 'Suture technique and wound healing in midline laparotomy incisions', *Eur J Surg*, 162. pp. 605-609.
- Janz, A. B. (2013) *Regions and Planes of the Abdomen: Overview, Abdominal Skin, Superficial Fascia*. Available at: <http://emedicine.medscape.com/article/1923166-overview#a1>.
- Jeekel, J. (2002) 'Meta-analysis of techniques for closure of midline abdominal incisions', *Br J Surg*, 89. pp. 1350-1356.
- Jenkins, T. P. (1976) 'The burst abdominal wound : a mechanical approach', *Br J Surg*, 63. pp. 873-876.
- Keith, L. and Arthur, F. (2012) *Ovid: Clinically Oriented Anatomy*. Available at: <http://ueu.co/zh/ovid-clinically-oriented-anatomy-2/> (Accessed: 19 January 2018).
- Klinge, U. *et al.* (2001) 'Collagen I/III and matrix metalloproteinases (MMP) 1 and 13 in the fascia of patients with incisional hernias', *Journal of Investigative Surgery*, pp. 47–54. doi: 10.1080/089419301750072202.
- Kopf, M. *et al.* (1994) 'Impaired immune and acute-phase responses in interleukin-6-deficient mice', *Nature*, 368(6469), pp. 339–342. doi: 10.1038/368339a0.
- Kumar, C. D., Rao, T., & Kishore, N. (2016). Effect of Stitch Length on the Rate of Postoperative Wound Complications in Midline Incisions, *I5(5)*, 37–48. <https://doi.org/10.9790/0853-1505013748>

- Kumar, C.D., Thatha, R.V., Kishore, N.B. (2016) 'Effect of stitch length on the rate of postoperative wound complications in midline incisions', *IOSR Journal of Dental and Medical Sciences*. 15(5) Ver I. pp. 37-48. doi: 10.9790/0853-1505013748.
- Kumar, P., Kumar, S., Udupa, E.P., Kumar, U., Rao, P., Honnegowda, T. (2015) 'Role of Angiogenesis and Angiogenic Factors in Acute and Chronic Wound Healing', *Plastic and Aesthetic Research*. 2: 243. doi: 10.4103/2347-9264.165438.
- Kumar, R. and Hastir, A. (2017) 'Prospective Clinical Study: Mass Closure Versus Layer Closure of Abdominal Wall', *International Journal of Surgery and Medicine*, 3, p. 1. doi: 10.5455/ijsm.mass-closure-versus-layer-closure-abdominal-wall.
- Lambertz, A. *et al.* (2015) 'Polyvinylidene fluoride as a suture material: Evaluation of comet tail-like infiltrate and foreign body granuloma', *European Surgical Research*, 55(1-2), pp. 1-11. doi: 10.1159/000371797.
- Lau, F. H. and Pomahac, B. (2014) 'Wound healing in acutely injured fascia', *Wound Rep Reg*, 22, pp. 14-17. doi: 10.1111/wrr.12165.
- Lawrence, W.T., Norton, J.A., Sporn, M.B., Gorschboth, C., Grotendorst, G.R. (1986) 'The reversal of an Adriamycin induced healing impairment with chemoattractants and growth factors', *Ann Surg*. 203(2). pp. 142-7. doi: 10.1097/00000658-198602000-00006. [PubMed: 3511865].
- Lin, Z.-Q. (2003) 'Essential involvement of IL-6 in the skin wound-healing process as evidenced by delayed wound healing in IL-6-deficient mice', *Journal of Leukocyte Biology*, 73(6), pp. 713-721. doi: 10.1189/jlb.0802397.
- Lin, Z.-Q. (2003) 'Essential involvement of IL-6 in the skin wound-healing process as evidenced by delayed wound healing in IL-6-deficient mice', *Journal of Leukocyte Biology*, 73(6), pp. 713-721. doi: 10.1189/jlb.0802397.
- Liptan, G. L. (2010) 'Fascia: A missing link in our understanding of the pathology of fibromyalgia', *Journal of Bodywork and Movement Therapies*. Elsevier Ltd, 14(1), pp. 3-12. doi: 10.1016/j.jbmt.2009.08.003.

- Liu, J., Zhao, J., Li, S., Zhu, L., Yu, R., Qu, Y., *et al.* (2015) 'The expression level of TGF-beta1, TGF-beta3 and VEGF in transplanted oral mucosal and cutaneous wounds', *Clin Microbiol.* 4(2):198. pp. 1-7. doi:10.4172/2327-5073.1000198.
- Mary, C., Marois, Y. and King, M. W. (1998) 'Comparison of the in vivo behavior of polyvinylidene fluoride and polypropylene sutures used in vascular surgery', *ASAIO Journal*, pp. 199–206. Available at: <http://europaepmc.org/abstract/med/9617952>.
- Massague', J. (1990) 'The transforming growth factor-beta family'. *Annu Rev Cell Biol.* 6. pp. 597–641.
- Meijer, E. J. *et al.* (2013) 'The principles of abdominal wound closure', *Acta Chirurgica Belgica*, 113(4), pp. 239–244. doi: [http://dx.doi.org/10.1016/0890-4332\(93\)90047-Y](http://dx.doi.org/10.1016/0890-4332(93)90047-Y).
- Meijer, E.J., Timmermans, L., Jeekel, J., Lange, J.F., Muysoms, F.E. (2013) 'The principles of abdominal wound closure', *Acta Chir Belg*, 113. pp. 239-244.
- Mercandetti, A. M., Editor, C. and Molnar, J. A. (2015) 'Wound Healing and Repair: Overview, Types of Wound Healing, Categories of Wound Healing', pp. 1–7. Available at: <http://emedicine.medscape.com/article/1298129-overview>.
- Millan, F.A., Denhez, F., Kondaiah, P., Akhurst, R.J. (1991) 'Embryonic gene expression patterns of TGF beta 1, beta 2 and beta 3 suggest different developmental functions in vivo', *Development*. 111. pp. 131–143.
- Millbourn, D. (2009) 'Effect of Stitch Length on Wound Complications After Closure of Midline Incisions A Randomized Controlled Trial', *Archives of Surgery*, 144(11), p. 1056. doi: 10.1001/archsurg.2009.189.
- Millbourn, D., Cengiz, Y., Israelsson, L. A.(2009) 'Effect of stitch length on wound complications after closure of midline incisions : a randomized controlled trial', *Arch Surg*, 144. pp. 1056-1059.
- Mizell, J. S. (2015) 'Complications of abdominal surgical incisions', *UpToDate*, pp. 1–27.

- Murphy-Ullrich, J.E., Poczatek, M. (2000) 'Activation of latent TGF-beta by thrombospondin-1: mechanisms and physiology', *Cytokine Growth Factor Rev.* 11. pp. 59-69.
- Muysoms, F. E. *et al.* (2015) 'European Hernia Society guidelines on the closure of abdominal wall incisions', *Hernia : the journal of hernias and abdominal wall surgery*, 19(1), pp. 1–24. doi: 10.1007/s10029-014-1342-5.
- Naka, T., Nishimoto, N. and Kishimoto, T. (2002) 'The paradigm of IL-6: from basic science to medicine.', *Arthritis research*, 4 Suppl 3, pp. S233–S242. doi: 10.1186/ar565.
- Nall, A.V., Brownlee, R.E., Colvin, C.P., Schultz, G., Fein, D., Cassisi, N.J., *et al.* (1996) 'Transforming growth factor beta 1 improves wound healing and random flap survival in normal and irradiated rats', *Arch Otolaryngol Head Neck Surg.* 122(2). pp. 171-7. doi: 10.1001/archotol.1996.01890140057011. [PubMed: 8630211].
- Nout, E. *et al.* (2007) 'Creep Behavior of Commonly Used Suture Materials in Abdominal Wall Surgery', *Journal of Surgical Research*, 138(1), pp. 51–55. doi: 10.1016/j.jss.2006.06.001.
- O'Kane, S., Ferguson, M.W. (1997) 'Transforming growth factor beta s and wound healing', *Int J Biochem Cell Bio.* 29(1). pp. 63-78. doi: 10.1016/S1357-2725(96)00120-3. [PubMed: 9076942].
- O'Kane, S., Ferguson, M.W. (1997) 'Transforming growth factor beta s and wound healing', *Int J Biochem Cell Biol.* 29. pp, 63-78.
- Orsted, H.L., Keast, D., Forest-Lalade, L., Francoise, M., (2004) 'The basic principles of wound care'. *Wound Care Canada.* 9 (2). pp. 4-12.
- Osther, P. J. *et al.* (1995) 'Randomized comparison of polyglycolic acid and polyglyconate sutures for abdominal fascial closure after laparotomy in patients with suspected impaired wound healing', *British Journal of Surgery*, 82(8), pp. 1080–1082. doi: 10.1002/bjs.1800820824.
- Pelton, R.W., Dickinson, M.E., Moses, H.L., Hogan, B.L. (1990) 'In situ hybridization analysis of TGF beta 3 RNA expression during mouse

- development: comparative studies with TGF beta 1 and beta 2', *Development*. 110. pp. 609–620.
- Pelton, R.W., Saxena, B., Jones, M., Moses, H.L., Gold, L.I. (1991) 'Immunohistochemical localization of TGF beta 1, TGF beta 2, and TGF beta 3 in the mouse embryo: expression patterns suggest multiple roles during embryonic development', *J Cell Biol*. 115. pp. 1091–1105.
- Pereira, J. A., Pera, M., Grande, L. (2012) 'Incidence of incisional hernia after open and laparoscopic colorectal cancer resection, Elevada incidencia de hernia incisional tras reseccion abierta y laparoscopica por cancer colorrectal', *Cir Esp*.
- Popa, F. and Georgescu, A. V (2017) 'Abdominal Wall Reconstruction after Flap Surgery and the Effect on the Immune System', *Hindawi*, 2017, pp. 1–10.
- Radu, P. *et al.* (2013) 'Molecular factors of failure in incisional hernia surgery.', Rahbari, N. N. *et al.* (2009) 'Current practice of abdominal wall closure in elective surgery? Is there any consensus?', *BMC Surgery*, 9(1), pp. 1–8. doi: 10.1186/1471-2482-9-8.
- Ramshorst Gabrielle H van, Eker Hasan H, Hop Wim C.J., Jeekel Johannes, L. J. F. (2012) 'Impact of incisional hernia on health-related quality of life and body image : a prospective cohort study', *AJS*. Elsevier Inc., 204(2), pp. 144–150. doi: 10.1016/j.amjsurg.2012.01.012.
- Robson, M. C. (2000). Wound healing trajectories as predictors of effectiveness of therapeutic agents, *Archives of surgery*. doi: 10.1001/archsurg.135.7.773.
- Roses, R. E. and Morris, J. B. (2013) 'Incisions, Closures, And Management of The Abdominal Wound', in Zinner, M. J. and Ashley, S. W. (eds) *Maingot's Abdominal Operations*. 12th editi. New York: Mc Graw Hill Companies, pp. 99–122.
- Ruhrberg, C. (2001) 'Endogeneous inhibitors of angiogenesis', *J Cell Sci*. 114(Pt 18). pp. 3215-6. [PubMed: 11591810].
- Satteson, E. S. (2017) *Materials for Wound Closure: Wound Healing and Closure, Suture Characteristics, Suture Materials*, Medscape. Available at: <https://emedicine.medscape.com/article/1127693->

[overview%0Ahttps://emedicine.medscape.com/article/1127693-overview#a3](https://emedicine.medscape.com/article/1127693-overview#a3).

- Savari, R., Shafiei, M., Galehdari, H., Kesmati, M. (2019) 'Expression of VEGF and TGF- β genes in skin wound healing process induced using phenytoin in male rats', *Jundishapur J Health Sci.* 11(1):e86041. pp. 1-5. doi: 10.5812/jjhs.86041.
- Stewart, R. J. *et al.* (1981) 'the Wound Fibroblast and Macrophage .1. Wound Cell-Population Changes Observed in Tissue-Culture', *British Journal of Surgery*, 68(2), pp. 125–128.
- Thiruvoth, F. *et al.* (2015) 'Current concepts in the physiology of adult wound healing', *Plastic and Aesthetic Research*, 2(5), p. 250. doi: 10.4103/2347-9264.158851.
- Urban, E. *et al.* (1994) 'Why make monofilament sutures out of polyvinylidene fluoride?', *ASAIO journal (American Society for Artificial Internal Organs : 1992)*, pp. 145–156. Available at: http://journals.lww.com/asaiojournal/Abstract/1994/04000/Why_Make_Monofilament_Sutures_Out_of.6.aspx%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/8003751.
- van Ramshorst, G. H., Eker H. H., Hop W. C., Jeekel J., Lange J. F. (2012) 'Impact of incisional hernia on health-related quality of life and body image : a prospective cohort study', *Am J Surg.* 204. pp. 144-150.
- Veljkovic, R. *et al.* (2010) 'Prospective Clinical Trial of Factors Predicting the Early Development of Incisional Hernia after Midline Laparotomy', *Journal of the American College of Surgeons.* Elsevier Inc., 210(2), pp. 210–219. doi: 10.1016/j.jamcollsurg.2009.10.013.
- Weiland, D. E., Bay, R. C., Del Sordi, S. (1998) 'Choosing the best abdominal closure by meta-analysis', *Am J Surg*, 176. pp. 666-670.
- Wharton, K., Derynck, R. (2009) 'TGFbeta family signaling: novel insights in development and disease', *Development.* 136. pp. 3691–3697.
- Wu, M.Y., Hill, C.S. (2009) 'Tgf-beta superfamily signaling in embryonic development and homeostasis', *Dev Cell.* 16. pp. 329–343.

Xing, L. *et al.* (2013) 'Early laparotomy wound failure as the mechanism for incisional hernia formation', *Journal of Surgical Research*. Elsevier Ltd, 182(1), pp. 1–8. doi: 10.1016/j.jss.2012.09.009.