

- Ahmed, N., Argirov, O.K., Minhas, H.S., Cordeiro, C.A., Thornalley, P.J. 2002. Assay of advanced glycation endproducts (AGEs): surveying AGEs by chromatographic assay with derivatization by 6-aminoquinolyl-N-hydroxysuccinimide-carbamate and application to Nepsilon-carboxymethyl-lysine- and Nepsilon-(1-carboxyethyl)lysine-modified albumin. *Biochem. J.* 364:1–14.
- Akirav, E.M., Preston-Hurlburt, P., Garyu, J., Henegariu, O., Clynes, R., Schmidt, A. M., *et al.* 2012. RAGE expression in human T cells: a link between environmental factors and adaptive immune responses. *PloS One* 7 e34698.
- American Diabetes Association. 2019. Standards of Medical Care in Diabetes--2019. *Diabetes Care*, 37(Supplement_1), S14–S80. <https://doi.org/10.2337/dc14-S014>
- American Diabetes Association. 2020. 2. Classification and diagnosis of diabetes: Standards of Medical Care in Diabetes—2020. *Diabetes Care* ;43(Suppl. 1):S14–S31. <https://doi.org/10.2337/dc20-S002>
- Anandharaj, M., Sivansankari, B. and Rani, R.P. 2014. Effects of probiotics, probiotics, dan synbiotics on hypercholesterolemia: a review. *Chinese J. of Biology.* 2014, 1-7. <https://doi.org/10.1155/2014/572754>.
- Araki, N., Higashi, T., Mori, T., Shibayama, R., Kawabe, Y., Kodama, T., *et al.* 1995. Macrophage scavenger receptor mediates the endocytic uptake and degradation of advanced glycation end products of the Maillard reaction. *Eur. J. Biochem.* 230:408–415.
- Asmarani F, Wirjatmadi B, Adriani M, Gizi D, Masyarakat K. 2015. Pengaruh Pemberian Suplementasi Tepung Tempe Terhadap Kadar Gula Darah Tikus Wistar Diabetes Mellitus. *Ilm Kedokt.* 4(2):24–35.
- Aoki, H., Furuya, Y., Endo, Y., & Fujimoto, K. (2003). Effect of γ -aminobutyric acid-enriched tempeh-like fermented soybean (gaba-tempeh) on the blood pressure of spontaneously hypertensive rats. *Biosci, Biotech and Biochem*, 67(8), 1806–1808. <https://doi.org/10.1271/bbb.67.1806>.
- Aoki, H., Uda, I., Tagami, K., Furuya, Y., Endo, Y., & Fujimoto, K. 2003. The Production of a New Tempeh-like Fermented Soybean Containing a High Level of γ -Aminobutyric Acid by Anaerobic Incubation with *Rhizopus*. *Biosci, Biotech and Biochem*, 67(5), 1018–1023. <https://doi.org/10.1271/bbb.67.1018>.
- Aslam, A., Singh, J., and Rajbhandari, S. 2014. Pathogenesis of painful diabetic neuropathy. *Pain Res. Treat.* 2014:412041. doi: 10.1155/2014/412041.



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WYAN, DANI M, J.dr. Arta, Farmawati, Ph.D., Dr. Pramudji Hastuti, Apt., MS., Dr. Prasetyastuti, Apt., M.kes
Astawan, M., Marudhyah, Y. S., & Wihaya, C. H. 2018. Potential of Bioactive Components in

Tempe for the Treatment of Obesity. *Jurnal Gizi Dan Pangan*, 13(2), 79–86.

<https://doi.org/10.25182/jgp.2018.13.2.79-86>.

- Bierhaus, A., Humpert, P.M., Morcos, M., Wendt, T., Chavakis, T., Arnold, B., *et al.* 2005. Understanding RAGE, the receptor for advanced glycation end products. *J. Mol. Med.* 83:876–886.
- Bhattamisra, S.K., Mohapatra, L., Panda, B.P., Parida, S., 2013. Effect of Isoflavone rich Soya seed extract on glucose utilization and endurance capacity in diabetic rat. *Diabetol. Croat.* 42: 42–52.
- Birgani, G.A., Aharnargarpor, A., Khorsandi, L., Moghaddam, F.H., 2018. Anti-diabetic effect of betulinic acid on streptozotocin-nicotinamide induced diabetic male mouse model. *Braz. J. Pharm.* 54(2):e17171.
- Bintanah S, Kusuma HS. 2010. Pengaruh Pemberian Bekatul Dan Tepung Tempe Terhadap Profil Gula Darah Pada Tikus Yang Diberi Alloxan (The Influence of Rice bran and Flour Tempeh on Blood Sugar Profile in Rats Fed Alloxan). *Jurnal Gizi dan Pangan*. 1(2).
- Bonizzi G, Bebien M, Otero DC, Johnson-Vroom KE, Cao Y, Vu D, *et al.* 2004. Activation of IKK α target genes depends on recognition of specific κ B binding sites by RelB:p52 dimers". *The EMBO Journal*. 23 (21): 4202–10. doi:10.1038/sj.emboj.7600391.
- Brownlee, M. 2005. The pathobiology of diabetic complications: A unifying mechanism. *Diabetes*, 54(6), 1615–1625. <https://doi.org/10.2337/diabetes.54.6.1615>
- Cai D, Yuan M, Frantz DF, Melendez PA, Hansen L, Lee J, *et al.* 2005. Local and systemic insulin resistance resulting from hepatic activation of IKK β and NF- κ B. *Nat Med*. 11, 183–190.
- Cardozo AK, Heimberg H, Heremans Y, Leeman R, Kutlu B, Kruhoffer M, *et al.* 2001. A comprehensive analysis of cytokine-induced and nuclear factor- κ B dependent genes in primary rat pancreatic betacells. *J Biol Chem*. 276(48) 879–886.
- Carroll K.K. 1991. Review of clinical studies on cholesterol-lowering response to soy protein. *J. Am. Diet. Assoc.* 91:820–827
- Chavakis, T., Bierhaus, A., Al-Fakhri, N., Schneider, D., Witte, S., Linn, T., *et al.* 2003. The pattern recognition receptor (RAGE) is a counterreceptor for leukocyte integrins: a novel pathway for inflammatory cell recruitment. *J. Exp. Med.* 198:1507–1515.
- Chen, L., Chen, R., Wang, H., & Liang, F. 2015. Mechanisms Linking Inflammation to Insulin Resistance. *Int. J. of Endocrinology*, 2015. <https://doi.org/10.1155/2015/508409>



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WAYAN DANI M., d. Arta Farmawati, Ph.D.; Dr. Pramudji Hastuti, Apt, MS.; Dr. Prasetyastuti, Apt, M.kes
Chen, Y., Akirav, E.M., Chen, W., Henegarlu, O., Moser, B., Desai, D., et al. 2008. RAGE

ligation affects T cell activation and controls T cell differentiation. *J. Immunol.* 181:4272–4278.

- Chung, E.Y., Kim, B.H., Hong, J.T., Lee, C.K., Ahn, B., Nam, S.Y., et al. 2011. Resveratrol down-regulates interferon-gamma-inducible inflammatory genes in macrophages: molecular mechanism via decreased STAT-1 activation. *J. Nutr. Biochem.* 22:902–909.
- Dhakal, R., Bajpai, V. K., & Baek, K. H. 2012. Production of GABA (γ -aminobutyric acid) by microorganisms: A review. *Brazilian J. of Microbiology*, 43(4), 1230–1241. <https://doi.org/10.1590/S1517-83822012000400001>.
- Dao, M.C., Everard, A., Aron-Wisnewsky, J., Sokolovska, N., Prifti, E., Verger, E.O., et al. 2016. Akkermansia muciniphila improved metabolic health during a dietary intervention in obesity: relationship with gut microbiome richness and ecology. *Gut Microbes.* 65(3), 426-436. <https://doi.org/10.1136/gutjnl-2014-308778>.
- Dupre-Crochet, S., Erard, M., Nubetae, O. 2013. ROS production in phagocytes: why, when, and where?. *J. Leukoc. Biol.* 94:657–670.
- Dumitriu, I.E., Baruah, P., Valentinis, B., Voll, R.E., Herrmann, M., Nawroth, P.P., et al. 2005. Release of high mobility group box 1 by dendritic cells controls T cell activation via the receptor for advanced glycation end products. *J. Immunol.* 174:7506–7515
- Dyer, D.G., Dunn, J.A., Thorpe, S.R., Lyons, T.J., McCance, D.R., Baynes, J.W. 1992. Accumulation of Maillard reaction products in skin collagen in diabetes and aging. *Acad. Sci.* 663:421–422.
- Duru, K.C., Kovaleva, E.G., Danilova, I.G., van der Bijl, P., Belousova, A. V., 2018. The potential beneficial role of isoflavones in type 2 diabetes mellitus. *Nutr. Res.* 59: 1–15. doi:10.1016/j.nutres.2018.06.005
- Evans JL., Goldfine ID., Maddux BA., Grodsky GM. 2003. Are Oxidative Stress-Activated Signaling Pathways Mediators of Insulin Resistance and β -Cell Dysfunction? *Diabetes.* 52: 1-8.
- Everard, A., Belzer, C., Geurts, L., Ouwerkerk, J.P., Druart, C., Bindels, L.B., et al. 2012. Cross-talk between Akkermansia muciniphila and intestinal epithelium controls diet-induced obesity. *Proceedings of the National Academy of Sciences*, 110(22), 9066-9071. <https://doi.org/10.1073/pnas.1219451110>.
- Farquhar C.M., Marjoribanks J., Lethaby A., Lamberts Q., Suckling J.A., 2005:2015. Cochrane HT Study Group Long term hormone therapy for perimenopausal and postmenopausal women. *Cochrane Database Syst. Rev.* CD004143. doi: 10.1002/14651858



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS GADJAH MADA WYAN DANI M J, ds. Arta Farmawati, Ph.D.; Dr. Pramudji Hastuti, Apt. MS.; Dr. Prasetyastuti, Apt. M.kes
Firdaus, Marhyati, S.A., Roosita, K. 2018. Model Tikus Diabetes Yang Diinduksi

Streptozotocin- Sukrosa Untuk Pendekatan Penelitian Diabetes Streptozotocin , Sucrose-Induce Diabetic Male Rats Model for Research. *J. MKMI* 12: 29–34.

- Frommhold, D., Kamphues, A., Hepper, I., Pruenster, M., Lukic, I.K., Socher, I., *et al.* 2010. RAGE and ICAM-1 cooperate in mediating leukocyte recruitment during acute inflammation in vivo. *Blood*. 116:841–849.
- Grimm, S., Horlacher, M., Catalgol, B., Hoehn, A., Reinheckel, T., Grune, T. 2012. Cathepsins D and L reduce the toxicity of advanced glycation end products. *Free Radic. Biol. Med.* 52:1011–1023
- Grimm, S. , Ernst, L., Grotzinger, N., Hohn, A., Breusing, N., Reinheckel, T., Grune, T. 2010. Cathepsin D is one of the major enzymes involved in intracellular degradation of AGE-modified proteins. *Free Radic. Res.* 44:1013–1026.
- Grimm, S., Ott, C., Horlacher, M., Weber, D., Hohn, A., Grune, T. 2012. Advanced-glycation-end-product-induced formation of immunoproteasomes: involvement of RAGE and Jak2/STAT1. *Biochem. J.* 448:127–139.
- Han, X., Tao, Y.L., Deng, Y.P., Yu, J.W., Cai, J., Ren, G.F., *et al.* 2017. Metformin ameliorates insulinitis in STZ-induced diabetic mice. *PeerJ*:5.
- Haron, H., & Raob, N. 2014. Changes in Macronutrient, Total Phenolic and Anti-Nutrient Contents during Preparation of Tempeh. *J. of Nutri. & Food Sci.* 04(03). <https://doi.org/10.4172/2155-9600.1000265>
- Hofmann, S.M., Dong, H.J., Li, Z., Cai, W., Altomonte, J., Thung, S.N., Zeng, F., Fisher, E. A., Vlassara, H. 2002. Improved insulin sensitivity is associated with restricted intake of dietary glycoxidation products in the db/db mouse. *Diabetes* 51:2082–2089.
- Hoie L.H., Guldstrand M., Sjöholm A., Graubaum H.J., Gruenwald J., Zunft H.J.F., Lueder W. 2007. Cholesterol-lowering effects of a new isolated soy protein with high levels of nondenaturated protein in hypercholesterolemic patients. *Adv. Ther.* 24:439–447. doi: 10.1007/BF02849913.
- Hotamisligil GS, Arner P, Caro JF, Atkinson RF, Spiegelman BM. 1995. Increased adipose tissue expression of tumor necrosis factor- α in human obesity and insulin resistance. *J Clin Invest.* 95, 2409–2415.
- Huang, C.Y., Lai, K.Y., Hung, L.F., Wu, W.L., Liu, F.C. , Ho, L.J. 2011. Advanced glycation end products cause collagen II reduction by activating Janus kinase/signal transducer and activator of transcription 3 pathway in porcine chondrocytes. *Rheumatology* 50:1379–1389.



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WAYAN DANI M., dr. Arta Farmawati, Ph.D., Dr. Pramudji Hastuti, Apt, MS, Dr. Prasetyastuti, Apt, M.kes
Huang, J.S., Guh, J.Y., Chen, H.C., Hung, W.C., Lai, Y.H., Chuang, L.Y. 2001. Role of

receptor for advanced glycation end-product (RAGE) and the JAK/STAT- signaling pathway in AGE-induced collagen production in NRK-49F cells. *J. Cell. Biochem.* 81:102–113.

- Huttunen, H.J., Fages, C., Rauvala, H. 1999. Receptor for advanced glycation end products (RAGE)-mediated neurite outgrowth and activation of NF-kappaB require the cytoplasmic domain of the receptor but different downstream signaling pathways. *J. Biol. Chem.* 274:19919–19924.
- Hwang J.T, Park I.J, Shin J.I, Lee Y.K, Lee S.K, Baik H.W, *et al.* 2005. Genestein, EGCG and capcaisin inhibit adipocyte dif-ferentiation process via activating AMP- activated protein kinase. *Biochem Biophys Res Commun.* 338(2):694-699
- Hwang, J. H., Wu, S. J., Wu, P. L., Shih, Y. Y., and Chan, Y. C., 2018. Neuroprotective effect of tempeh against lipopolysaccharide-induced damage in BV-2 microglial cells. *Nutr. Neurosci.* doi: 10.1080/1028415X.2018.1456040.
- Ilatovskaya, D. V., Levchenko, V., Lowing, A., Shuyskiy, L. S., Palygin, O., Staruschenko, A. 2015. Podocyte injury in diabetic nephropathy: implications of angiotensin II – dependent activation of TRPC channels. *Sci. Rep.* 5:17637. doi: 10.1038/srep17637
- Ishihara, H., Sasaoka, T., Hori, H., Wada, T., Hirai, H., Haruta, T., *et al.* 1999. Molecular cloning of SH2- containing inositol phosphatase 2 (SHIP2) and its role in the regulation of insulin signaling. *Biochem. Biophys. Res. Commun.* 260, 265–272.
- John, W.G., Lamb, E.J. 1993. The Maillard or browning reaction in diabetes. *Eye* 7 (Pt 2):230–237.
- Jono, T., Miyazaki, A., Nagai, R., Sawamura, T., Kitamura, T., Horiuchi, S. 2002. Lectin- like oxidized low density lipoprotein receptor-1 (LOX-1) serves as an endothelial receptor for advanced glycation end products (AGE). *FEBS Lett.* 511:170–174.
- Jung, T., Catalgol, B., Grune, T. 2009. The proteasomal system. *Mol. Asp. Med.* 30:191–296.
- Kamioka, M., Ishibashi, T., Ohkawara, H., Nagai, R., Sugimoto, K., Uekita, H., *et al.* 2011. Involvement of membrane type 1-matrix metalloproteinase (MT1-MMP) in RAGE activation signaling path- ways. *J. Cell. Physiol.* 226:1554–1563.
- Karin, M., and Ben-Neriah, Y. 2000. Phosphorylation meets ubiquitination: the control of NF-kB activity. *Annu. Rev. Immunol.* 18, 621–663. doi: 10.1146/annurev.immunol.18.1.62.
- Kasper, M., Funk, R.H. 2001. Age-related changes in cells and tissues due to advanced glycation end products (AGEs). *Gerontol. Geriatr.* 32:233–243.
- Kierdorf, K., Fritz, G. 2013. RAGE regulation and signaling in inflammation and beyond. *J. Leukoc. Biol.* 94: 55–68.



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WAYAN DANI M., dr. Arta Farmawati, Ph.D.; Dr. Pramudji Hastuti, Apt., MS.; Dr. Prasetyastuti, Apt., M.kes
Kim, E.K., Kwon, K.B., Song, M.Y., Seo, S.W., Park, S.J., Ka, S.O., et al. 2007. Genistein
Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

protects pancreatic beta cells against cytokine-mediated toxicity. *Mol. Cell. Endocrinol.* 278, 18–28.

- Kitada, M., Zhang, Z., Mima, A., and King, GL. 2010. Molecular mechanisms of diabetic vascular complications. *J. Diabetes Invest.* 1, 77–89. doi: 10.1111/j.2040-1124.2010.00018.x
- Khoshnoodi, J., Pedchenko, V., Hudson, B.G. 2008. Mammalian collagen IV. *Microsc. Res. Tech.* 71:357–370.
- Koh, S. P., Jamaluddin, A., Alitheen, N. B., & Long, K. 2012. Nutritional values of tempe inoculated with different strains of *Rhizopus*: its γ -aminobutyric acid content and antioxidant property. *J. of Tropical Agric. and Food Sci.* 40(2), 181–192.
- Křížová, L., Dadáková, K., Kašparovská, J., & Kašparovský, T. 2019. Isoflavones. *Molecules (Basel, Switzerland)*, 24(6), 1076. doi:10.3390/molecules24061076.
- Lapolla, A., Fedele, D., Martano, L., Arico, N.C., Garbeglio, M., Traldi, P., et al. 2001. Advanced glycation end products: a highly complex set of biologically relevant compounds detected by mass spectrometry. *J. Mass Spectrom.*: JMS 36:370–378.
- Lee, J., Yee, S.T., Kim, J.J., Choi, M.S., Kwon, E.Y., Seo, K.I., Lee, M.K., 2010. Ursolic acid ameliorates thymic atrophy and hyperglycemia in streptozotocin-nicotinamide-induced diabetic mice. *Chem. Biol. Interact.* (3): 635-642.
- Lee DS, Lee SH. 2001. Genestein, a soy isoflavone, is a potent α -glucosidase inhibitor. *FEBS Letters.* 501(1):84-86.
- Lethaby A.E., Brown J., Marjoribanks J., Kronenberg F., Roberts H., Eden J. 2007. Phytoestrogens for vasomotor menopausal symptoms. *Cochrane Database Syst. Rev.* CD001395. doi: 10.1002/14651858
- Li, Y.M., Mitsuhashi, T., Wojciechowicz, D., Shimizu, N., Li, J., Stitt, A., et al. 1996. Molecular identity and cellular distribution of advanced glycation endproduct receptors: relationship of p60 to OST-48 and p90 to 80K-H membrane proteins. *Proc. Natl. Acad. Sci. USA* 93: 11047–11052.
- Liu, D., Zhen, W., Yang, Z., Carter, J.D., Si, H., Reynolds, K.A., 2006. Genistein acutely stimulates insulin secretion in pancreatic beta-cells through a cAMP- dependent protein kinase pathway. *Diabetes.* 55, 1043–1050.
- Liu, Y., Liang, C., Liu, X., Liao, B., Pan, X., Ren, Y., et al. 2010. AGEs increased migration and inflammatory responses of adventitial fibroblasts via RAGE, MAPK and NF-kappaB pathways. *Atherosclerosis.* 208:34–42.



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WAYAN DANI M., dr. Arta Farmawati, Ph.D., Dr. Pramudji Hastuti, Apt., MS., Dr. Prasetyastuti, Apt., M.kes
Liu, Y., Qu, Y., Wang, R., Ma, Y., Xia, C., Gao, C., et al. 2012. The alternative crosstalk
Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

between RAGE and nitrate thiolredoxin inactivation during diabetic myocardial ischemia-reperfusion injury. *Am. J. Physiol. Endocrinol. Metab.* 303:E841–852.

- Lopez-parra, V., Mallavia, B., Egido, J., dan Gomez-guerrero, C. 2012. Immunoinflammation in diabetic nephropathy: molecular mechanisms and therapeutic options. *Diabetic Nephropathy*, ed. J. S. D. Chan (Rijeka: InTech)
- Lu, M., Wang, R., Song, X., Chibbar, R., Wang, X., Wu, L., & Meng, Q. H. 2008. Dietary soy isoflavones increase insulin secretion and prevent the development of diabetic cataracts in streptozotocin-induced diabetic rats. *Nutrition Research*, 28, 464–471. <https://doi.org/10.1016/j.nutres.2008.03.009>.
- Maiti, D., Majumdar, M., 2012. Impact of bioprocessing on phenolic content and antioxidant activity of two edible seeds to improve hypoglycemic functionality. *Int. J. Shoulder Surg.* 6: 31–36. doi:10.4103/2229-5119.96958
- Madiraju AK., Erion DM., Rahimi Y., Zhang XM., Braddock DT., Albright RA., et al. 2014. Metformin suppresses gluconeogenesis by inhibiting mitochondrial glycerophosphate dehydrogenase. *Nature*. 510 (7506): 542–6. doi:10.1038/nature13270.
- Meng, XM., Tang, PMK., Li, J., dan Lan, HY. 2015. TGF- β /Smad signaling in renal fibrosis. *Front. Physiol.* 6:82. doi: 10.3389/fphys.2015.00082.
- Messina M., Kucuk O., Lampe J.W. 2006. An overview of the health effects of isoflavones with an emphasis on prostate cancer risk and prostate-specific antigen levels. *J. AOAC Int.* 89:1121–1134.
- Messina M., Hilakivi-Clarke L. 2009. Early intake appears to be the key to the proposed protective effects of soy intake against breast cancer. *Nutr. Cancer*. 61:792–798. doi: 10.1080/01635580903285015.
- Miyazaki, A., Nakayama, H., Horiuchi, S. 2002. Scavenger receptors that recognize advanced glycation end products. *Trends Cardiovasc. Med.* 12: 258–262
- Moore, M. 2018. Food as medicine: Diet, diabetes management, and the patient in twentieth century Britain. *J. of the History of Medicine and Allied Sci.* 73(2):150-167.
- Moser, B., Desai, D.D., Downie, M.P., Chen, Y., Yan, S.F., Herold, K., et al. 2007. Receptor for advanced glycation end products expression on T cells contributes to antigen-specific cellular expansion in vivo. *J. Immunol.* 179: 8051–8058.
- Nah, S.-S., Choi, I.-Y., Yoo, B., Kim, Y.G., Moon, H.-B., Lee, C.-K. 2007. Advanced glycation end products increases matrix metalloproteinase-1, -3, and -13, and TNF- α in human osteoarthritic chondrocytes. *FEBS Lett.* 581:1928–1932.



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WYAN DANIM J., dr. Arta Farmawati, Ph.D.; Dr. Pramudji Hastuti, Apt, MS.; Dr. Prasetyastuti, Apt, M.kes
Nakajima, N., Nozaki, N., Ishihara, K., Ishikawa, A., & Tsuji, H. 2005. Analysis of isoflavone
Universitas Gadjah Mada, 2020 | Diunduh dari <http://efd.repository.ugm.ac.id/>

content in tempeh, a fermented soybean, and preparation of a new isoflavone-enriched tempeh. *J. Biosci Bioeng.* 100(6), 685–687. <https://doi.org/10.1263/jbb.100.685>

- Nass, N., Bartling, B., Navarrete Santos, A., Scheubel, R.J., Borgermann, J., Silber, R. E., Simm, A. 2007. Advanced glycation end products, diabetes and ageing. *Gerontol. Geriatr.* 40:349–356.
- Neeper, M., Schmidt, A.M., Brett, J., Yan, S.D., Wang, F., Pan, Y.C., *et al.* 1992. Cloning and expression of a cell surface receptor for advanced glycosylation end products of proteins. *J. Biol. Chem.* 267:14998–15004.
- Nicholl, I.D., Bucala, R. 1998. Advanced glycation endproducts and cigarette smoking. *Cell. Mol. Biol.* 44:1025–1033.
- O'Brien, J., Morrissey, P.A. 1989. Nutritional and toxicological aspects of the Maillard browning reaction in foods. *Crit. Rev. Food Sci. Nutr.* 28:211–248.
- Ohashi, K., Takahashi, H.K., Mori, S., Liu, K., Wake, H., Sadamori, H., *et al.* 2010. Advanced glycation end products enhance monocyte activation during human mixed lymphocyte reaction. *Clin. Immunol.* 134:345–353.
- Ohgami, N., Nagai, R., Ikemoto, M., Arai, H., Kuniyasu, A., Horiuchi, S., Nakayama, H. 2001. CD36, a member of class B scavenger receptor family, is a receptor for advanced glycation end products. *Acad. Sci.* 947:350–355.
- Ohgami, N., Nagai, R., Ikemoto, M., Arai, H., Miyazaki, A., Hakamata, H., *et al.* 2002. CD36, serves as a receptor for advanced glycation endproducts (AGE). *J. Diabetes Complicat.* 16:56–59.
- Ohgami, N., Nagai, R., Miyazaki, A., Ikemoto, M., Arai, H., Horiuchi, S., Nakayama, H. 2001. Scavenger receptor class B type I-mediated reverse cholesterol transport is inhibited by advanced glycation end products. *J. Biol. Chem.* 276:13348–13355.
- Palanisamy, N., Kannappan, S., Anuradha, C.V., 2011. Genistein modulates NF- κ B-associated renal inflammation, fibrosis and podocyte abnormalities in fructose-fed rats. *Eur. J. Pharmacol.* 667, 355–364.
- Patel, S., & Santani, D. 2009. Role of NF- κ B in the pathogenesis of diabetes and its associated complications. *Pharmacol. Reports*, 61(4), 595–603. [https://doi.org/10.1016/S1734-1140\(09\)70111-2](https://doi.org/10.1016/S1734-1140(09)70111-2)
- Peng, Y., Kim, J.M., Park, H.S., Yang, A., Islam, C., Lakatta, E.G., *et al.* 2016. AGE-RAGE signal generates a specific NF- κ B RelA “barcode” that directs collagen I expression. *Sci. Rep.* 6(January), 1–10. <https://doi.org/10.1038/srep18822>



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WYAN DANI M. J., dr. Arta Farmawati, Ph.D., Dr. Pramudji Hastuti, Apt, MS, Dr. Prasetyastuti, Apt., M.kes
Pittenger, G., Vinik, A., dan Pittenger, G.L. 2003. Nerve growth factor and diabetic neuropathy.
Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Exp. Diabesity Res. 4, 271–285. doi: 10.1080/15438600390249718.

- Platta, H.W., Stenmark, H. 2011. Endocytosis and signaling. *Curr. Opin. Cell Biol.* 23:393–403.
- Polianskyte-prause, Z., Tolvanen, T. A., Lindfors, S., Dumont, V., Van, M., Wang, H., *et al.* 2018. Metformin increases glucose uptake and acts renoprotectively by reducing SHIP2 activity. *The FASEB j.* 1–12. doi 10.1096/fj.201800529RR.
- Pollreis, A., Hudson, B.I., Chang, J.S., Qu, W., Cheng, B., Papapanou, P.N., *et al.* 2010. Receptor for advanced glycation endproducts mediates pro-atherogenic responses to periodontal infection in vascular endothelial cells. *Atherosclerosis* 212:451–456.
- Prud'Homme GJ, Glinka Y, Udovyk O, Hasilo C, Paraskevas S, Wang Q. 2014. GABA protects pancreatic beta cells against apoptosis by increasing SIRT1 expression and activity. *Biochem Biophys Res Commun.* 452(3):649–54.
- Puteri, N. E., Astawan, M., Palupi, N. S., Wresdiyati, T., & Takagi, Y. 2018. Characterization of biochemical and functional properties of water-soluble tempe flour. *Food Sci. and Tech.*, 38, 147–153. <https://doi.org/10.1590/fst.13017>
- Rena, G., Hardie, D. G., & Pearson, E. R. 2017. The mechanisms of action of metformin. *Diabetologia*, Vol. 60, pp. 1577–1585. <https://doi.org/10.1007/s00125-017-4342-z>
- Reutens, AT., and Atkins, RC. 2011. Epidemiology of diabetic nephropathy. *Contrib. Nephrol.* 170, 1–7. doi: 10.1159/000324934.
- Sami, W., Ansari, T., Butt, N. S., & Hamid, M. 2017. Effect of diet on type 2 diabetes mellitus: A review. *Int. J. of health scie.*, 11(2), 65–71.
- Sano, H., Higashi, T., Matsumoto, K., Melkko, J., Jinnouchi, Y., Ikeda, K., *et al.* 1998. Insulin enhances macrophage scavenger receptor-mediated endocytic uptake of advanced glycation end products. *J. Biol. Chem.* 273 :8630–8637.
- Saono, S., Hull, R.R., Dhamcharee, B., 1986. A Concise Handbook of Indi-genous Fermented Foods in the Asia Countries. Indonesian Institute of Sciences, Jakarta, Indonesia.
- Saputra, N.T., Suartha, I.N., Dharmayudha, A.A.G.O., 2018. Agen Diabetagonik Streptozotocin untuk Membuat Tikus Putih Jantan Diabetes Mellitus. *Bul. Vet. Udayana* 10: 116. doi:10.24843/bulvet.2018.v10.i02.p02
- Shoelson SE, Lee J, Goldfine AB. 2006. Inflammation and insulin resistance. *J Clin Invest.* 116: 1793-1801.



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WYAN DANIM, dr. Arta Farmawati, Ph.D., Dr. Pramudji Hastuti, Apt, MSc, Dr. Prasetyastuti, Apt, M.kes
Shu X.O., Zheng Y., Cai H., Gu K., Chen Z., Zheng W., Lu W. 2009. Soy food intake and

breast cancer survival. *JAMA*. 302:2437–2443. doi: 10.1001/jama.2009.1783.

- Signorini, C., Carpen, A., Coletto, L., Borgonovo, G., Galanti, E., Capraro, J., *et al.* 2018. Enhanced vitamin B12 production in an innovative lupin tempeh is due to synergic effects of *Rhizopus* and *Propionibacterium* in cofermentation. *Int. J. of Food Sci. and Nutri.* 69(4), 451–457. <https://doi.org/10.1080/09637486.2017.1386627>
- Sorkin, A., Zastrow, M.V. 2009. Endocytosis and signalling: intertwining molecular networks, *Nat. Rev. Mol. Cell Biol.* 10:609–622.
- Stephanie, Kartawidjajaputra, F., Silo, W., Yogiara, Y., & Suwanto, A. 2019. Tempeh consumption enhanced beneficial bacteria in the human gut. *Food Research*, 3(1), 57–63. [https://doi.org/10.26656/fr.2017.3\(1\).230](https://doi.org/10.26656/fr.2017.3(1).230).
- Stokes SE, Winn LM. 2014. NF-kappa B signaling is increased in HD3 cells following exposure to 1, 4-ben- zoquinone: role of reactive oxygen species and p38- MAPK. *Toxicol Sci SCI.* 137: 303-310
- Stryer L. 1995. *Biochemistry* (Fourth ed.). New York: W.H. Freeman and Company. pp. 773–74. [ISBN 0-7167-2009-4](https://doi.org/10.1016/0-7167-2009-4).
- Sun, C., Li, X., Liu, L., Canet, MJ., Guan, Y., Fan, Y., *et al.* 2016. Effect of fasting time on measuring mouse blood glucose level. *Int. J. Clin. Exp. Med.* 9(2), 4186–4189.
- Suryavanshi, SV., & Kulkarni, YA. 2017. NF-κβ: A potential target in the management of vascular complications of diabetes. *Front. Pharmacol.*, 8(NOV), 1–12. <https://doi.org/10.3389/fphar.2017.00798>
- Tamura, Y., Adachi, H., Osuga, J.-I., Ohashi, K., Yahagi, N., Sekiya, M., *et al.* 2003. FEEL-1 and FEEL-2 are endocytic receptors for advanced glycation end products. *J. Biol. Chem.* 278:12613–12617.
- Tan, S. Y., Ling, J., Wong, M., Sim, Y. J., Wong, S. S., Abdelgadir, S., *et al.* 2018. Type 1 and 2 diabetes mellitus: A review on current treatment approach and gene therapy as potential intervention. *Diabetes & Metab. Syndr.: Clin. Res. & Rev.* <https://doi.org/10.1016/j.dsx.2018.10.008>
- Teede H.J., Dalais F.S., Kotsopoulos D., Liang Y.L., Davis S., McGrath B.P. 2001. Dietary soy has both beneficial and potentially adverse cardiovascular effects: A placebo-controlled study in men and postmenopausal women. *J. Clin. Endocrinol. Metab.* 86:3053–3060. doi: 10.1210/jc.86.7.3053.
- Thornalley, P.J., Langborg, A., Minhas, H.S. 1999. Formation of glyoxal, methylglyoxal and 3-deoxyglucosone in the glycation of proteins by glucose. *Biochem. J.* 344 (Pt 1):109–116.



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WAYAN DANIM J. de Arta Firmawati, Ph.D.; Dr. Pramudji Hastuti, Apt, MS.; Dr. Prasejyastuti, Apt, M.kes
Tian, J., Avalos, A.M., Mao, S.-Y., Chen, B., Senthil, K., 2007. Toll-like receptor 9-dependent

activation by DNA-containing immune complexes is mediated by HMGB1 and RAGE. *Nat. Immunol.* 8: 487–496.

- Tian, J., Dang, H. N., Yong, J., Chui, W. S., Dizon, M. P. G., Yaw, C. K. Y., *et al.* 2011. Oral treatment with γ -aminobutyric acid improves glucose tolerance and insulin sensitivity by inhibiting inflammation in high fat diet-fed mice. *PLoS ONE*, 6(9), 1–7. <https://doi.org/10.1371/journal.pone.0025338>.
- Utari DM, Rimbawan R, Riyadi H, Muhilal M, Purwastyastuti P. 2011. Potensi Asam Amino pada Tempe untuk Memperbaiki Profil Lipid dan Diabetes Mellitus. *Kesmas Natl Public Heal* J.5(4):166.
- Valencia, J.V., Weldon, S.C., Quinn, D., Kiers, G.H., DeGroot, J., TeKoppele, J.M., Hughes, T. E. 2004. Advanced glycation end product ligands for the receptor for advanced glycation end products: biochemical characterization and formation kinetics. *Anal. Biochem.* 324:68–78
- Vlassara, H., Li, Y.M., Imani, F., Wojciechowicz, D., Yang, Z., Liu, F.T., Cerami, A. 1995. Identification of galectin-3 as a high-affinity binding protein for advanced glycation end products (AGE): a new member of the AGE-receptor complex. *Mol. Med.* (Cambridge, MA) 1:634–646.
- Vlassara, H. 2001. The AGE-receptor in the pathogenesis of diabetic complications. *Diabetes/Metab. Res. Rev.* 17:436–443.
- Wang, Y., Wang, H., Piper, M.G., McMaken, S., Mo, X., Opalek, J., Schmidt, A.M., Marsh, C. B. 2010. sRAGE induces human monocyte survival and differentiation. *J. Immunol.* 185:1822–1835.
- Wang G. 2014. Raison d'être of insulin resistance: the adjustable threshold hypothesis. *J. R. Soc., Interface.* 11 (101): 20140892. doi:10.1098/rsif.2014.0892.
- Wang, Z., Zhang, J., Chen, L., Li, J., Zhang, H., Guo, X., 2019. Glycine suppresses AGE/RAGE signaling pathway and subsequent oxidative stress by restoring Glo1 function in the aorta of diabetic rats and in HUVECs. *Oxid. Med. Cell. Longev.* 2019. doi:10.1155/2019/4628962
- Wei, A.C., Ren, X., Jiang, Y., Jin, H., Liu, N., Li, J. 2013. Advanced glycation end products accelerate rat vascular calcification through RAGE/oxidative stress. *BMC Cardiovasc. Disord.* 13:13.
- WHO Global Report. 2016. Global Report on Diabetes. *Isbn*, 978, 6–86. Retrieved from <http://www.who.int/about/licensing/>



PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED GLYCATION END PRODUCTS DAN NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS MENCIT MODEL DIABETES MELLITUS TIPE 2

UNIVERSITAS
GADJAH MADA

WYAN DANIM, J. dr. Arta Farmawati, Ph.D.; Dr. Pramudji Hastuti, Apt, MS; Dr. Prasetyastuti, Apt, M.kes
Wong, A., Dukic-Stefanovic, S., Gasic-Milenkovic, J., Schinzel, R., Wiesinger, H., Riederer, P., & Münch, G. 2001. Anti-inflammatory antioxidants attenuate the expression of inducible nitric oxide synthase mediated by advanced glycation endproducts in murine microglia. *Eur. J. of Neurosci.* 14(12), 1961–1967. <https://doi.org/10.1046/j.0953-816X.2001.01820.x>

Xie, F., Zhu, J., Hou, B., Wang, Y., Meng, F., Ren, Z., & Ren, S. 2017. Inhibition of NF- κ B activation improves insulin resistance of L6 cells. *Endocr. J.*, 64(7), 685–693. <https://doi.org/10.1507/endocrj.EJ17-0012>.

Xing, L., Gao Hong, L., Dai, GY., Sun, HM., Xu, HQ. 2016. Food advanced glycation end products aggravate the diabetic vascular complications via modulating the AGEs/RAGE pathway. *Chin. J. Nat. Med.* 14, 844–855. doi: 10.1016/S1875-5364(16)30101-7.

Yagame, M., Kim, Y., Zhu, D., Suzuki, D., Eguchi, K., Nomoto, Y., *et al.* 1995. Differential distribution of type IV collagen chains in patients with diabetic nephropathy in non-insulin-dependent diabetes mellitus. *Nephron*, 70:42–48.

Yagihashi, S., Mizukami, H., dan Sugimoto, K. 2011. Mechanism of diabetic neuropathy: where are we now and where to go? *J. Diabetes Invest.* 2, 18–32. doi: 10.1111/j.2040-1124.2010.00070.x

Ye Y.-B., Tang X.-Y., Verbruggen M.A., Su Y.-X. 2006. Soy isoflavones attenuate bone loss in early postmenopausal Chinese women: A single-blind randomized, placebo-controlled trial. *Eur. J. Nutr.* 45:327–334. doi: 10.1007/s00394-006-0602-2

Yonekura, H., Yamamoto, Y., Sakurai, S., Petrova, R.G., Abedin, M.J., Li, H., *et al.* 2003. Novel splice variants of the receptor for advanced glycation end-products expressed in human vascular endothelial cells and pericytes, and their putative roles in diabetes-induced vascular injury. *Biochem. J.* 370: 1097–1109.

Zastrow, M.V., Sorkin, A. 2007. Signaling on the endocytic pathway. *Curr. Opin. Cell Biol.* 19:436–445.

Zhang, F., Kent, K.C., Yamanouchi, D., Zhang, Y., Kato, K., Tsai, S., Nowygrad, R., Schmidt, A. M., Liu, B. 2009. Anti-receptor for advanced glycation end products therapies as novel treatment for abdominal aortic aneurysm. *Ann. Surg.* 250:416–423.

Zhang, L., Bukulin, M., Kojro, E., Roth, A., Metz, V.V., Fahrenholz, F., *et al.* 2008. Receptor for advanced glycation end products is subjected to protein ectodomain shedding by metalloproteinases. *J. Biol. Chem.* 283:35507–35516.

Zhang, M., Kho, A.L., Anilkumar, N., Chibber, R., Pagano, P.J., Shah, A.M., Cave, A.C. 2006. Glycated proteins stimulate reactive oxygen species production in cardiac myocytes: involvement of Nox2 (gp91phox)-containing NADPH oxidase. *Circulation.* 113:1235–1243.



**PENGARUH SUPLEMEN BERBASIS TEMPE TERHADAP EKSPRESI RECEPTOR ADVANCED
GLYCATION END PRODUCTS DAN
NUCLEAR FACTOR KAPPA- LIGHT-CHAIN-ENHANCER OF ACTIVATED B CELLS PADA PANKREAS
MENCIT MODEL DIABETES
MELLITUS TIPE 2**

UNIVERSITAS
GADJAH MADA

WAYAN DANI M., dr. Arta Farmawati, Ph.D., Dr. Pramudji Hastuti, Apt. MS., Dr. Prasetiyastuti, Apt. M.kes
Zhao, Y., Krishnamurthy, B., Mohan, ZU., Kay, I.W., dan Thomas, HE. 2011. NF- κ B in type I

diabetes. *Inflamm. Allergy Drug Targets.* 10, 208–217. doi:
10.2174/187152811795564046.