

## DAFTAR PUSTAKA

- Adams, R.M., McCarl, B.A. & Mearns, L.O. 2003. The effects of spatial scale of climate scenarios on economic assessments: An example from U.S. agriculture. *Climatic Change*, 60(1–2): 131–148.
- Anandababu, D., Purushothaman, B.M. & Suresh Babu, S. 2018. Estimation of Land Surface Temperature using LANDSAT 8 Data. *International Journal of Advance Research*, 4(2): 177–186. Tersedia di [www.IJARIT.com](http://www.IJARIT.com).
- Bakar, S.B.A., Pradhan, B., Lay, U.S. & Abdullahi, S. 2016. Spatial assessment of land surface temperature and land use/land cover in Langkawi Island. *IOP Conference Series: Earth and Environmental Science*.
- Benali, A., Carvalho, A.C., Nunes, J.P., Carvalhais, N. & Santos, A. 2012. Estimating air surface temperature in Portugal using MODIS LST data. *Remote Sensing of Environment*, 124: 108–121. Tersedia di <http://dx.doi.org/10.1016/j.rse.2012.04.024>.
- Brecht 2018. *Remote Sensing Indices*. <https://medium.com/>. Tersedia di <https://medium.com/regen-network/remote-sensing-indices-389153e3d947>.
- Córdova, M., Célleri, R., Shellito, C.J., Orellana-Alvear, J., Abril, A. & Carrillo-Rojas, G. 2016. Near-surface air temperature lapse rate over complex terrain in the Southern Ecuadorian Andes: implications for temperature mapping. *Arctic, Antarctic, and Alpine Research*, 48(4): 678–684. Tersedia di <http://www.bioone.org/doi/10.1657/AAAR0015-077>.
- Dikau, R., Brabb, E.E. & Mark, R.M. 1991. *Landform Classification of New Mexico by Computer*. U.S. Department of The Interior U.S. Geological Survey, .
- Dodson, R. & Marks, D. 1997. Daily air temperature interpolated at high spatial resolution over a large mountainous region. *Climate Research*, 8: 1–20.
- Effendy, S. 2009. Dampak pengurangan RTH perkotaan terhadap peningkatan suhu udara dengan metode penginderaan jauh. *Jurnal Agromet*, 23(November): 169–181.
- Gallo, K.P., McNab, A.L., Karl, T.R., Brown, J.F., Hood, J.J. & Tarpley, J.D. 1993.

- The Use of NOAA AVHRR Data for Assessment of the Urban Heat Island Effect. *Journal of Applied Meteorology*, 32(5): 899–908. Tersedia di [https://doi.org/10.1175/1520-0450\(1993\)032%3C0899:TUONAD%3E2.0.CO](https://doi.org/10.1175/1520-0450(1993)032%3C0899:TUONAD%3E2.0.CO).
- Guillevic, P., Göttsche, F., Nickeson, J., Hulley, G., Ghent, D., Yu, Y., Trigo, I., Hook, S., Sobrino, J.A., Remedios, J., Román, M. & Camacho, F. 2018. Land surface temperature product validation best practice protocol version 1.1. *Best Practice for Satellite-Derived Land Product Validation (p. 58): Land Product Validation Subgroup (WGCV/CEOS)*, doi. Tersedia di [http://ceos.org/document\\_management/Working\\_Groups/WGCV/CEOS\\_LST\\_PROTOCOL\\_Oct2017\\_v1.0.0.pdf](http://ceos.org/document_management/Working_Groups/WGCV/CEOS_LST_PROTOCOL_Oct2017_v1.0.0.pdf).
- Hammond, E.H. 1954. Small-Scale Continental Landform Maps. *Annals of the Association of American Geographers*, 44(1): 33–42. Tersedia di <https://www.tandfonline.com/doi/abs/10.1080/00045605409352120>.
- Hengl, T. 2006. Finding the right pixel size. *Computers and Geosciences*, 32(9): 1283–1298.
- Jacobson, M.Z. 2005. *Fundamentals of atmospheric modeling*. Second Ed ed. Cambridge, UK: Cambridge University Press. Tersedia di [www.cambridge.org/9780521839709](http://www.cambridge.org/9780521839709).
- Jatmiko, R.H. 2016. *Penggunaan citra saluran inframerah termal untuk studi perubahan liputan lahan dan suhu sebagai indikator perubahan iklim perkotaan di Yogyakarta*. Gadjah Mada University.
- Karagülle, D. & Frye, C. n.d. *Improved Hammond 's Landform Classification and Method for Global 250-m Elevation Data*.
- Katpatal, Y.B., Kute, A. & Satapathy, D.R. 2008. Surface and air temperature studies in relation to land use/ land cover of Nagpur urban area using Landsat 5 TM data. *Journal of Urban Planning and Development*, 134(3): 110–118.
- Kattel, D.B., Yao, T., Yang, K., Tian, L., Yang, G. & Joswiak, D. 2013. Temperature lapse rate in complex mountain terrain on the southern slope of the central Himalayas. *Theoretical and Applied Climatology*, 113(3–4): 671–682.

- Kawashima, S., Ishida, T., Minomura, M. & Miwa, T. 2000. Relations between surface temperature and air temperature on a local scale during winter nights. *Journal of Applied Meteorology*, 39(9): 1570–1579.
- Khandelwal, S., Goyal, R., Kaul, N. & Mathew, A. 2018. Assessment of land surface temperature variation due to change in elevation of area surrounding Jaipur, India. *Egyptian Journal of Remote Sensing and Space Science*, 21(1): 87–94. Tersedia di <https://doi.org/10.1016/j.ejrs.2017.01.005>.
- Kshetri, T. 2018. *NDVI, NDBI & NDWI Calculation Using Landsat 7,8*. <https://www.linkedin.com/>. Tersedia di <https://www.linkedin.com/pulse/ndvi-ndbi-ndwi-calculation-using-landsat-7-8-tek-bahadur-kshetri>.
- Lai, Y.J., Li, C.F., Lin, P.H., Wey, T.H. & Chang, C.S. 2012. Comparison of MODIS land surface temperature and ground-based observed air temperature in complex topography. *International Journal of Remote Sensing*, 33(24): 7685–7702.
- Li, Y., Zeng, Z., Zhao, L. & Piao, S. 2015. Spatial patterns of climatological temperature lapse rate in mainland China: a multi-time scale investigation. *Geophysical Research: Atmospheres*, 120: 2661–2675.
- Liu, S., Su, H., Zhang, R., Tian, J. & Wang, W. 2016. Estimating the surface air temperature by remote sensing in Northwest China using an improved advection-energy balance for air temperature model. *Advances in Meteorology*, 2016: 11.
- Minder, J.R., Mote, P.W. & Lundquist, J.D. 2010. Surface temperature lapse rates over complex terrain: lessons from the Cascade Mountains. *Journal of Geophysical Research Atmospheres*, 115(14): 1–13.
- Morgan, J.M. & Lesh, A. 2005. Developing Landform Maps Using ESRI'S ModelBuilder. *ESRI User Conference Proceedings*, 1–11. Tersedia di <http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Developing+landform+maps+using+esri?s+ModelBuilder#0>.
- Mutiibwa, D., Strachan, S. & Albright, T. 2015. Land surface temperature and surface air temperature in complex terrain. *IEEE Journal of Selected Topics in*

*Applied Earth Observations and Remote Sensing*, 8(10): 4762–4774.

- Neteler, M. 2010. Estimating daily land surface temperatures in mountainous environments by reconstructed MODIS LST data. *Remote Sensing*, 2(1): 333–351.
- Nichol, J.E. & Wong, M.S. 2008. Spatial variability of air temperature and appropriate resolution for satellite - derived air temperature estimation. *International Journal of Remote Sensing*, 29(24): 7213–7223.
- Nieto, H., Sandholt, I., Aguado, I., Chuvieco, E. & Stisen, S. 2011. Air temperature estimation with MSG-SEVIRI data: Calibration and validation of the TVX algorithm for the Iberian Peninsula. *Remote Sensing of Environment*, 115(1): 107–116. Tersedia di <http://dx.doi.org/10.1016/j.rse.2010.08.010>.
- Prata, A.J. 1994. Land surface temperatures derived from the advanced very high resolution radiometer and the along-track scanning radiometer: 2. Experimental results and validation of AVHRR algorithms. *Journal of Geophysical Research: Atmospheres*, 99(D6): 13025–13058. Tersedia di <https://doi.org/10.1029/94JD00409>.
- Rajeshwari, A. & Mani, N.D. 2014. Estimation of Land Surface Temperature of Dindigul District Using Landsat 8 Data. *International Journal of Research in Engineering and Technology*, 03(05): 122–126.
- Rozenstein, O., Qin, Z., Derimian, Y. & Karnieli, A. 2014. Derivation of land surface temperature for landsat-8 TIRS using a split window algorithm. *Sensors (Switzerland)*, 14(4): 5768–5780.
- Samodra, G., Chen, G., Sartohadi, J. & Kasama, K. 2017. Comparing data-driven landslide susceptibility models based on participatory landslide inventory mapping in Purwosari area, Yogyakarta, Java. *Environmental Earth Sciences*, 76(4): 1–19.
- Skykas, D. 2020. *Spectral Indices with Multispectral Satellite Data*. Tersedia di <https://www.geo.university/>.

<https://www.geo.university/pages/spectral-indices-with-multispectral-satellite-data>.

Sobrino, J.A., Jiménez-Muñoz, J.C., Sòria, G., Romaguera, M., Guanter, L., Moreno, J., Plaza, A. & Martínez, P. 2008. Land surface emissivity retrieval from different VNIR and TIR sensors. *IEEE Transactions on Geoscience and Remote Sensing*, 46(2): 316–327.

Steinacker, R., Ratheiser, M., Bica, B., Chimani, B., Dorninger, M., Gepp, W., Lotteraner, C., Schneider, S. & Tschannett, S. 2006. A mesoscale data analysis and downscaling method over complex terrain. *Monthly Weather Review*, 134(10): 2758–2771.

Stisen, S., Sandholt, I., Nørgaard, A., Fensholt, R. & Eklundh, L. 2007. Estimation of diurnal air temperature using MSG SEVIRI data in West Africa. *Remote Sensing of Environment*, 110(2): 262–274.

Stull, B.R. 1988. *An introduction to boundary layer meteorology*. First Edit ed. AA Dordrecht, The Netherlands: Kluwer Academic Publishers. Tersedia di <http://link.springer.com/10.1007/978-94-009-3027-8>.

Sun, Y., Wang, J., Zhang, R., Gillies, R.R., Xue, Y. & Bo, Y. 2005. Air temperature retrieval from remote sensing data based on thermodynamics. *Theoretical and Applied Climatology*, 80: 37–48.

Sutikno 1996. Geomorphology of Yogyakarta area and its surrounding proposed as a geomorphological field laboratory. *The Indonesian Journal of Geography*, 28(71): 1–10.

Tsou, J., Zhuang, J., Li, Y. & Zhang, Y. 2017. Urban Heat Island Assessment Using the Landsat 8 Data: A Case Study in Shenzhen and Hong Kong. *Urban Science*, 1(1): 10.

USGS, D. of the I. 2016. *Landsat 8 (L8) data users handbook version 2.0*. Sioux Falls, South Dakota: United State Geological Survey.

Wang, Y., Wang, L., Li, X. & Chen, D. 2018. Temporal and spatial changes in estimated near-surface air temperature lapse rates on Tibetan Plateau.

*International Journal of Climatology*, 38(7): 2907–2921.

- Weiss, S.B. 2005. *Topoclimate and microclimate in the Monarch Butterfly biosphere reserve*.
- Weng, Q., Fu, P. & Gao, F. 2014. Generating daily land surface temperature at Landsat resolution by fusing Landsat and MODIS data. *Remote Sensing of Environment*, 145: 55–67. Tersedia di <http://dx.doi.org/10.1016/j.rse.2014.02.003>.
- Xu, H. 2006. Modification of normalised difference water index (NDWI) to enhance open water features in remotely sensed imagery. *International Journal of Remote Sensing*, 27(14): 3025–3033.
- Xu, Y., Knudby, A. & Ho, H.C. 2014. Estimating daily maximum air temperature from MODIS in British Columbia, Canada. *International Journal of Remote Sensing*, 35(24): 8108–8121.
- Xu, Y., Qin, Z. & Shen, Y. 2012. Study on the estimation of near-surface air temperature from MODIS data by statistical methods. *International Journal of Remote Sensing*, 33(24): 7629–7643.
- Yang, Y., Cai, W. & Yang, J. 2017. Evaluation of MODIS land surface temperature data to estimate near-surface air temperature in Northeast China. *Remote Sensing*, 9(5): 410. Tersedia di <http://www.mdpi.com/2072-4292/9/5/410>.
- Yang, Z., Hanna, E. & Callaghan, T. V. 2011. Modeling surface-air-temperature variation over complex terrain around Abisko, Swedish Lapland: Uncertainties of measurements and models at different scales. *Geografiska Annaler, Series A: Physical Geography*, 93(2): 89–112.
- Yu, X., Guo, X. & Wu, Z. 2014. Land surface temperature retrieval from Landsat 8 TIRS-comparison between radiative transfer equation-based method, split window algorithm and single channel method. *Remote Sensing*, 6(10): 9829–9852.
- Zakšek, K. & Schroedter-Homscheidt, M. 2009. Parameterization of air temperature in high temporal and spatial resolution from a combination of the SEVIRI and

MODIS instruments. *ISPRS Journal of Photogrammetry and Remote Sensing*, 64(4): 414–421. Tersedia di <http://dx.doi.org/10.1016/j.isprsjprs.2009.02.006>.

Zhu, W., Lu, A. & Jia, S. 2013. Estimation of daily maximum and minimum air temperature using MODIS land surface temperature products. *Remote Sensing of Environment*, 130: 62–73. Tersedia di <http://dx.doi.org/10.1016/j.rse.2012.10.034>.