

DAFTAR PUSTAKA

- Al-Ishaq, R. K. *et al.* (2019) 'Flavonoids and their anti-diabetic effects: Cellular mechanisms and effects to improve blood sugar levels', *Biomolecules*, 9(9). doi: 10.3390/biom9090430.
- Bai, L. *et al.* (2019) 'Antidiabetic Potential of Flavonoids from Traditional Chinese Medicine: A Review', *American Journal of Chinese Medicine*, 47(5), pp. 933–957. doi: 10.1142/S0192415X19500496.
- Basciano, H., Federico, L. and Adeli, K. (2005) 'Fructose, insulin resistance, and metabolic dyslipidemia', *Nutrition and Metabolism*, 2, pp. 1–14. doi: 10.1186/1743-7075-2-5.
- Bouayed, J. and Bohn, T. (2010) 'Exogenous antioxidants - Double-edged swords in cellular redox state: Health beneficial effects at physiologic doses versus deleterious effects at high doses', *Oxidative Medicine and Cellular Longevity*, 3(4), pp. 228–237. doi: 10.4161/oxim.3.4.12858.
- Browning, J. D. *et al.* (2004) 'Molecular mediators of hepatic steatosis and liver injury Find the latest version: Molecular mediators of hepatic steatosis and liver injury', *The Journal of Clinical Investigation*, 114(2), pp. 147–152. doi: 10.1172/JCI200422422.The.
- Bumke-Vogt, C. *et al.* (2014) 'The flavones apigenin and luteolin induce FOXO1 translocation but inhibit gluconeogenic and lipogenic gene expression in human cells', *PLoS ONE*, 9(8). doi: 10.1371/journal.pone.0104321.
- Carpentier, Y. and Sobotka, L. (2008) 'Basics in clinical nutrition: Lipid metabolism', *e-SPEN*, 3(5), pp. 188–191. doi: 10.1016/j.eclnm.2008.06.005.
- Chayati, I. *et al.* (2019) 'Anthocyanin Extract of Purple Corn Improves Hyperglycemia and Insulin Resistance of Rats Fed High Fat and Fructose Diet via GLP1 and GLP1R Mechanism', *Journal of Food and Nutrition Research*, 7(4), pp. 303–310. doi: 10.12691/jfnr-7-4-7.
- Chung, A. P. Y. S. *et al.* (2018) 'Geraniin Protects High-Fat Diet-Induced Oxidative Stress in Sprague Dawley Rats', 5(March), pp. 1–14. doi: 10.3389/fnut.2018.00017.
- Chyau, C. C. *et al.* (2020) 'Antrodan alleviates high-fat and high-fructose diet-induced fatty liver disease in C57BL/6 mice model via AMPK/Sirt1/SREBP-1c/PPAR γ pathway', *International Journal of Molecular Sciences*, 21(1). doi: 10.3390/ijms21010360.

- Clifford, T. *et al.* (2015) ‘The potential benefits of red beetroot supplementation in health and disease’, *Nutrients*, 7(4), pp. 2801–2822. doi: 10.3390/nu7042801.
- Danuyanti, I. G. A. N., Farmawati, A. and Sunarti (2018) ‘High Fiber Diet Suppression of TLR4 and NF κ β Gene Expression Correlated with TNF- α and IL-6 Levels in Hypertriglyceridemia Rats’, *Jfmr*, 6(12), pp. 725–730. doi: 10.12691/jfmr-6-12-2.
- Danuyanti, I. G. A. N., Farmawati, A. and Sunarti (2019) ‘Beneficial effects of high-fiber diet on the expression and level of intercellular adhesion molecule-1 of hypercholesterolemia rats’, *Mediterranean Journal of Nutrition and Metabolism*, 12(3), pp. 305–313. doi: 10.3233/MNM-190288.
- Deshmukh, G. P., Sindhav, R. and Jose, N. (2018) ‘Application of Beetroot as Natural Coloring Pigment and Functional Ingredient in Dairy and Food Products’, 7(12), pp. 2010–2016.
- El-Hawary, S. S. *et al.* (2017) ‘Investigation of some chemical constituents, cytotoxicity and antioxidant activities of beta vulgaris var. altissima cultivated in Egypt’, *Rasayan Journal of Chemistry*, 10(4), pp. 1391–1401. doi: 10.7324/rjc.2017.1041936.
- Elliott, S. S. *et al.* (2002) ‘Fructose, weight gain, and the insulin resistance syndrome’, *American Journal of Clinical Nutrition*, 76(5), pp. 911–922. doi: 10.1093/ajcn/76.5.911.
- Embries, S. A. S., Ongowski, G. E. D. and Ehrla, K. A. M. (2006) ‘Physiological Effects of Extraction Juices from Apple , Grape , and Red Beet Pomaces in Rats’.
- Evans, J. L. (2007) ‘Antioxidants: Do they have a role in the treatment of insulin resistance?’, *Indian Journal of Medical Research*, 125(3), pp. 355–372.
- Gao, H. *et al.* (2012) ‘Treatment with ginger ameliorates fructose-induced fatty liver and hypertriglyceridemia in rats: Modulation of the hepatic carbohydrate response element-binding protein-mediated pathway’, *Evidence-based Complementary and Alternative Medicine*, 2012. doi: 10.1155/2012/570948.
- Geidl-Flueck, B. and Gerber, P. A. (2017) ‘Insights into the hexose liver metabolism—glucose versus fructose’, *Nutrients*, 9(9). doi: 10.3390/nu9091026.
- Hannou, S. A. *et al.* (2018) ‘Fructose metabolism and metabolic disease’, *Journal of Clinical Investigation*, 128(2), pp. 545–555. doi: 10.1172/JCI96702.
- Harikumar, K. *et al.* (2013) ‘A Review on Hyperlipidemic’, *Nternational Journal of Novel Trends in Pharmaceutical Sciences*, 3(4), pp. 69–80. Available at:

http://www.ijntps.org/File_Folder/0043.pdf.

- Hayasaka, N. *et al.* (2018) ‘Absorption and Metabolism of Luteolin in Rats and Humans in Relation to in Vitro Anti-inflammatory Effects’, *Journal of Agricultural and Food Chemistry*, 66(43), pp. 11320–11329. doi: 10.1021/acs.jafc.8b03273.
- Herningtyas, E. H. and Ng, T. S. (2019) ‘Prevalence and distribution of metabolic syndrome and its components among provinces and ethnic groups in Indonesia’, *BMC Public Health*. *BMC Public Health*, 19(1), pp. 1–12. doi: 10.1186/s12889-019-6711-7.
- Hou, R., Panda, C. and Voruganti, V. S. (2019) ‘Heterogeneity in metabolic responses to dietary fructose’, *Frontiers in Genetics*, 10(OCT), pp. 1–10. doi: 10.3389/fgene.2019.00945.
- Janevski, M. *et al.* (2012) ‘Fructose containing sugars modulate mRNA of lipogenic genes ACC and FAS and protein levels of transcription factors ChREBP and SREBP1c with no effect on body weight or liver fat’, *Food and Function*, 3(2), pp. 141–149. doi: 10.1039/c1fo10111k.
- Jung, U. J., Cho, Y. and Choi, M. (2016) ‘Apigenin Ameliorates Dyslipidemia, Hepatic Steatosis and Insulin Resistance by Modulating Metabolic and Transcriptional Profiles in the Liver of High-Fat Diet-Induced Obese Mice’. doi: 10.3390/nu8050305.
- Kammoun, H. L. *et al.* (2009) ‘GRP78 expression inhibits insulin and ER stress–induced SREBP-1c activation and reduces hepatic steatosis in mice’, *Molecular Neurobiology*, 119(5), pp. 1201–1215. doi: 10.1172/JCI37007DS1.
- Kavya, B. (2017) ‘Lipids and its Metabolism’, *Journal of Cardiology & Cardiovascular Therapy*, 4(2). doi: 10.19080/jocct.2017.04.555635.
- Khitan, Z. and Kim, D. H. (2013) ‘Fructose: A key factor in the development of metabolic syndrome and hypertension’, *Journal of Nutrition and Metabolism*, 2013. doi: 10.1155/2013/682673.
- Kim, J., Wei, Y. and R. Sowers, J. (2010) ‘Role of Mitochondrial Dysfunction in Insulin Resistance’, *PMC*, 102(4), pp. 401–414. doi: 10.1161/CIRCRESAHA.107.165472.Role.
- Kolderup, A. and Svihus, B. (2015) ‘Fructose Metabolism and Relation to Atherosclerosis, Type 2 Diabetes, and Obesity’, *Journal of Nutrition and Metabolism*, 2015. doi: 10.1155/2015/823081.
- Koo, H. Y. *et al.* (2009) ‘Replacing dietary glucose with fructose increases ChREBP activity and SREBP-1 protein in rat liver nucleus’, *Biochemical and*

- Biophysical Research Communications*. Elsevier Inc., 390(2), pp. 285–289.
doi: 10.1016/j.bbrc.2009.09.109.
- Köseler, E. *et al.* (2018) ‘The effects of glucose and fructose on body weight and some biochemical parameters in rats’, *Progress in Nutrition*, 20(1), pp. 46–51.
doi: 10.23751/pn.v20i1.5956.
- Kwon, E. Y. *et al.* (2015) ‘Luteolin attenuates hepatic steatosis and insulin resistance through the interplay between the liver and adipose tissue in mice with diet-induced obesity’, *Diabetes*, 64(5), pp. 1658–1669. doi: 10.2337/db14-0631.
- Li, S. *et al.* (2018) ‘The Potential and Action Mechanism of Polyphenols in the Treatment of Liver Diseases’, *Oxidative Medicine and Cellular Longevity*, 2018. doi: 10.1155/2018/8394818.
- Liebert, M. A. (2005) ‘The Molecular Basis for Oxidative Stress-Induced Insulin Resistance’, *ANTIOXIDANTS & REDOX SIGNALING*, 7(7 & 8 July 2005), pp. 1040–1052. doi: 10.1089/ars.2005.7.1040.
- Lin, C. F. *et al.* (2018) ‘Epidemiology of Dyslipidemia in the Asia Pacific Region’, *International Journal of Gerontology*, 12(1), pp. 2–6. doi: 10.1016/j.ijge.2018.02.010.
- Linden, A. G. *et al.* (2018) ‘Interplay between ChREBP and SREBP-1c coordinates postprandial glycolysis and lipogenesis in livers of mice’, *Journal of Lipid Research*, 59(3), pp. 475–487. doi: 10.1194/jlr.M081836.
- Liu, C. *et al.* (2017) ‘Flavonoid-rich extract of paulownia fortunei flowers attenuates diet-induced hyperlipidemia, hepatic steatosis and insulin resistance in obesity mice by ampk pathway’, *Nutrients*, 9(9), pp. 1–15. doi: 10.3390/nu9090959.
- Livak, K. J. and Schmittgen, T. D. (2001) ‘Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method’, *Methods*, 25(4), pp. 402–408. doi: 10.1006/meth.2001.1262.
- Lozano, I. *et al.* (2016) ‘High-fructose and high-fat diet-induced disorders in rats: Impact on diabetes risk, hepatic and vascular complications’, *Nutrition and Metabolism*. Nutrition & Metabolism, 13(1). doi: 10.1186/s12986-016-0074-1.
- Ma, L., Robinson, L. N. and Towle, H. C. (2006) ‘ChREBP-Mlx is the principal mediator of glucose-induced gene expression in the liver’, *Journal of Biological Chemistry*, 281(39), pp. 28721–28730. doi: 10.1074/jbc.M601576200.
- Maithilikarpagaselvi, N. *et al.* (2016) ‘Curcumin inhibits hyperlipidemia and hepatic fat accumulation in high-fructose-fed male Wistar rats’, *Pharmaceutical*

- Biology*, 54(12), pp. 2857–2863. doi: 10.1080/13880209.2016.1187179.
- Mancini, A. *et al.* (2015) ‘Oxidative stress and metabolic syndrome: Effects of a natural antioxidants enriched diet on insulin resistance’, *Clinical Nutrition ESPEN*. Elsevier Ltd, 10(2), pp. e52–e60. doi: 10.1016/j.clnesp.2014.11.002.
- Mhatre V. Ho, Ji-Ann Lee, and K. C. M. (2012) ‘Fatty acid synthase and liver triglyceride metabolism: housekeeper or messenger’, *Bone*, 23(1), pp. 1–7. doi: 10.1038/jid.2014.371.
- Montgomery, M. K. and Turner, N. (2015) ‘Mitochondrial dysfunction and insulin resistance: an update’, *Endocrine Connection*, 4(1), pp. 1–15. doi: 10.1530/EC-14-0092.
- Moreno-Fernández, S. *et al.* (2018) ‘High fat/high glucose diet induces metabolic syndrome in an experimental rat model’, *Nutrients*, 10(10), pp. 1–15. doi: 10.3390/nu10101502.
- Nguyen, P. *et al.* (2008) ‘Liver lipid metabolism’, 92, pp. 272–283. doi: 10.1111/j.1439-0396.2007.00752.x.
- Noeman, S. A., Hamooda, H. E. and Baalash, A. A. (2011) ‘Biochemical study of oxidative stress markers in the liver, kidney and heart of high fat diet induced obesity in rats’, *Diabetology and Metabolic Syndrome*. BioMed Central Ltd, 3(1), p. 17. doi: 10.1186/1758-5996-3-17.
- Nomura, K. and Yamanouchi, T. (2012) ‘The role of fructose-enriched diets in mechanisms of nonalcoholic fatty liver disease’, *Journal of Nutritional Biochemistry*. Elsevier Inc., 23(3), pp. 203–208. doi: 10.1016/j.jnutbio.2011.09.006.
- Nouh, F., Omar, M. and Younis, M. (2019) ‘Risk Factors and Management of Hyperlipidemia (Review)’, 2(1), pp. 1–10. doi: 10.9734/AJCR/2019/45449.
- Ozkan, H. and Yakan, A. (2019) ‘Annals of Hepatology Dietary high calories from sunflower oil , sucrose and fructose sources alters lipogenic genes expression levels in liver and skeletal muscle in rats’, 18, pp. 715–724.
- Panchal, S. *et al.* (2011) ‘Rutin Attenuates Metabolic Changes , Nonalcoholic Steatohepatitis , and ...’, *Journal of Nutrition*, 20(20), pp. 1062–1069. doi: 10.3945/jn.111.137877.hyperglycemia.
- Pappachan, J. M. *et al.* (2017) ‘Non-alcoholic Fatty Liver Disease: A Clinical Update’, *Journal of Clinical and Translational Hepatology*, XX(XX), pp. XX–XX. doi: 10.14218/jcth.2017.00013.
- Rahal, A. *et al.* (2014) ‘Oxidative Stress , Prooxidants , and Antioxidants : The Interplay’, *Biomed Research International*. Hindawi Publishing Corporation,

2014. doi: 10.1155/2014/761264.

- Refaat, W. A. and El-Nassag, D. E. (2019) 'Antioxidant Activity of Beetroot Powder on Alloxan Induced Diabetic Rats', *Journal of Health Science*, 2019(22), pp. 111–126. doi: 10.21608/jedu.2019.73657.
- Regnault, T. R. *et al.* (2013) 'Fructose, pregnancy and later life impacts', *Clinical and Experimental Pharmacology and Physiology*, 40(11), pp. 824–837. doi: 10.1111/1440-1681.12162.
- Salehi, B. *et al.* (2019) 'The therapeutic potential of Apigenin', *International Journal of Molecular Sciences*, 20(6). doi: 10.3390/ijms20061305.
- De Silva, G. S. *et al.* (2019) 'Circulating serum fatty acid synthase is elevated in patients with diabetes and carotid artery stenosis and is LDL-associated', *Atherosclerosis*. Elsevier, 287(May), pp. 38–45. doi: 10.1016/j.atherosclerosis.2019.05.016.
- Song, Z., Xiaoli, A. M. and Yang, F. (2018) 'Regulation and metabolic significance of De Novo lipogenesis in adipose tissues', *Nutrients*, 10(10), pp. 1–22. doi: 10.3390/nu10101383.
- Steele, C. C., Pirkle, J. R. A. and Kirkpatrick, K. (2017) 'Diet-induced impulsivity: Effects of a high-fat and a high-sugar diet on impulsive choice in rats', *PLoS ONE*, 12(6), pp. 1–16. doi: 10.1371/journal.pone.0180510.
- Strable, M. and Ntambi, J. (2010) 'Genetic control of de novo lipogenesis', *Biochem Mol Biol*, 45(3), pp. 199–214. doi: 10.3109/10409231003667500.Genetic.
- Tangvarasittichai, S. (2015) 'Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus', *World Journal of Diabetes*, 6(3), p. 456. doi: 10.4239/wjd.v6.i3.456.
- Tranchida, F. *et al.* (2012) 'Long-term high fructose and saturated fat diet affects plasma fatty acid profile in rats', *Journal of Zhejiang University: Science B*, 13(4), pp. 307–317. doi: 10.1631/jzus.B1100090.
- Tranchida, F. *et al.* (2018) 'Long-term high fructose and saturated fat diet affects plasma fatty acid profile in rats To cite this version : HAL Id : hal-01734922 Long-term high fructose and saturated fat diet affects plasma fatty acid profile in rats'.
- Wong, R. H. F. *et al.* (2009) 'A Role of DNA-PK for the Metabolic Gene Regulation in Response to Insulin', *Cell*, 136(6), pp. 1056–1072. doi: 10.1016/j.cell.2008.12.040.
- Wootton-Beard, P. C. and Ryan, L. (2011) 'A beetroot juice shot is a significant and

convenient source of bioaccessible antioxidants’, *Journal of Functional Foods*. Elsevier Ltd, 3(4), pp. 329–334. doi: 10.1016/j.jff.2011.05.007.

Wroblewska, M., Juskiwicz, J. and Wiczowski, W. (2011) ‘Physiological properties of beetroot crisps applied in standard and dyslipidaemic diets of rats’, *Lipids in Health and Disease*, 10, pp. 1–8. doi: 10.1186/1476-511X-10-178.

Xu, X. *et al.* (2013) ‘Transcriptional control of hepatic lipid metabolism by SREBP and ChREBP’, *Seminars in Liver Disease*, 33(4), pp. 301–311. doi: 10.1055/s-0033-1358523.

Zhang, D. M., Jiao, R. Q. and Kong, L. D. (2017) ‘High dietary fructose: Direct or indirect dangerous factors disturbing tissue and organ functions’, *Nutrients*, 9(4). doi: 10.3390/nu9040335.

Zwarts, I. *et al.* (2019) ‘Identification of the fructose transporter GLUT5 (SLC2A5) as a novel target of nuclear receptor LXR’, *Scientific Reports*, 9(1). doi: 10.1038/s41598-019-45803-x.