



DAFTAR PUSTAKA

- Abdel-Kader, K., Unruh, M. L., & Weisbord, S. D. (2009). *Symptom burden, depression, and quality of life in chronic and end-stage kidney disease.* *Clinical Journal of the American Society of Nephrology*, 4(6), 1057–1064. <https://doi.org/10.2215/CJN.00430109>
- Aisara, S., Azmi, S., & Mefri, Y. (2018). Gambaran Klinis Penderita Penyakit Ginjal Kronik yang Menjalani Hemodialisis di RSUP Dr. M. Djamil Padang. *Jurnal Kesehatan Andallas*, 7(1), 42–50.
- Alhaj, E., Alhaj, N., Rahman, I., Niazi, T. O., Berkowitz, R., & Klapholz, M. (2013). Uremic cardiomyopathy: An underdiagnosed disease. *Congestive Heart Failure*, 19(4). <https://doi.org/10.1111/chf.12030>
- Almutary, H., Bonner, A., & Douglas, C. (2016). Original Research: Which Patients With Chronic Kidney Disease Have the Greatest *Symptom burden*? a Comparative Study of Advanced Ckd Stage and Dialysis Modality. *Journal of Renal Care*, 73–82.
- Alshammari, B., Noble, H., McAneney, H., & O'Halloran, P. (2019). An exploration of *symptom burden* and its management, in Saudi Arabian patients receiving haemodialysis, and their caregivers: A mixed methods study protocol. *BMC Nephrology*, 20(1), 1–9. <https://doi.org/10.1186/s12882-019-1424-9>
- Anita, D. C., & Novitasari, D. (2017). Kepatuhan Pembatasan Asupan Cairan Terhadap Lama Menjalani Hemodialisa. *Jurnal Kesehatan Masyarakat*, 1(1), 104–112.
- Atmojo, S. darmo. (1999). Prinsip dan Indikasi Hemodialisis. *Pertemuan Tahunan Ilmu Penyakit Dalam*, 167–175.
- Badariah, Kusuma, F. H. D., & Dewi, N. (2017). Karakteristik pasien penyakit ginjal kronik yang menjalani hemodialisis di rsud kabupaten kota baru. *Nursing News*, 2(Nomor 2), 281–285.
- Bayhakki, B., & Hasneli, Y. (2018). Hubungan Lama Menjalani Hemodialisis dengan Inter-Dialytic Weight Gain (IDWG) pada Pasien Hemodialisis. *Jurnal Keperawatan Padjadjaran*, 5(3), 242–248. <https://doi.org/10.24198/jkp.v5i3.646>
- Bokyoung, J. K., Park, K. K., Park, J. C. J. S. S., & Kim, C. K. Y. (2013). *Health-related quality of life with KDQOL-36 and its association with self-efficacy and treatment satisfaction in Korean dialysis patients.* 753–758. <https://doi.org/10.1007/s11136-012-0203-x>
- Brown, S. A., Tyrer, F. C., Clarke, A. L., Lloyd-Davies, L. H., Stein, A. G., Tarrant, C., Burton, J. O., & Smith, A. C. (2017). *Symptom burden in*



patients with chronic kidney disease not requiring renal replacement therapy.
Clinical Kidney Journal, 10(6), 788–796. <https://doi.org/10.1093/ckj/sfx057>

Caninsti, R. (2013). Kecemasan dan Depresi pada Pasien Gagal Ginjal Kronis yang Menjalani Terapi Hemodialisis. *Jurnal Psikologi Ulayat*, 1(2). <https://doi.org/10.24854/jpu22013-22>

Chang, V. T., Hwang, S. S., Thaler, H. T., Kasimis, B. S., & Portenoy, R. K. (2004). Memorial Symptom Assessment Scale. *Expert Review of Pharmacoeconomics & Outcomes Research*, 4(2), 171–178. <https://doi.org/10.1586/14737167.4.2.171>

Cleeland, C. S., Zhao, F., Chang, V. T., Sloan, J. A., O'Mara, A. M., Gilman, P. B., Weiss, M., Mendoza, T. R., Lee, J.-W., & Fisch, M. J. (2014). The *symptom burden* of cancer: evidence for a core set of cancer-related and treatment-related symptoms from the Eastern Cooperative Oncology Group's Symptom Outcomes and Practice Patterns Study. *NIH Public Access*, 23(1), 1–7. <https://doi.org/10.1038/jid.2014.371>

Dąbrowska-Bender, M., Dykowska, G., Żuk, W., Milewska, M., & Staniszewska, A. (2018). The impact on quality of life of dialysis patients with renal insufficiency. *Patient Preference and Adherence*, 12, 577–583. <https://doi.org/10.2147/PPA.S156356>

Davison, S. N., & Jhangri, G. S. (2010). Impact of Pain and *Symptom burden* on the Health-Related Quality of Life of Hemodialysis Patients. *Journal of Pain and Symptom Management*, 39(3), 477–485. <https://doi.org/10.1016/j.jpainsympman.2009.08.008>

Davison, S. N., Jhangri, G. S., & Johnson, J. A. (2006). Longitudinal validation of a modified Edmonton symptom assessment system (ESAS) in haemodialysis patients. *Nephrology Dialysis Transplantation*, 21(11), 3189–3195. <https://doi.org/10.1093/ndt/gfl380>

De Pasquale, C., Pistorio, M. L., Corona, D., Mistretta, A., Zerbo, D., Sinagra, N., Giaquinta, A., Tallarita, T., Esker, B., Mociskyte, D., Leonardi, A., Gula, R., Veroux, P., & Veroux, M. (2012). Correlational study between psychic symptoms and quality of life among hemodialysis patients older than 55 years of age. *Transplantation Proceedings*, 44(7), 1876–1878. <https://doi.org/10.1016/j.transproceed.2012.06.015>

Eckerblad, J., Theander, K., Ekdahl, A., Unosson, M., Wirehn, A., & Milberg, A. (2015). *Symptom burden in community-dwelling older people with multimorbidity : a cross-sectional study*. 1–9.

Farida, A. (2010). Pengalaman Klien Hemodialisis terhadap Kualitas Hidup dalam Konteks Asuhan Keperawatan di RSUP Fatmawati Jakarta. In *Tesis Universitas Indonesia*.

Fresenius Medical Care. (2014). *Annual report 2014*.



- Fromer, M., & Hogan, M. (2007). ESAs. *Oncology Times*, 29(10), 23–25.
<https://doi.org/10.1097/01.cot.0000282463.09167.44>
- Gapstur, R. L. (2007). *Symptom burden : A Concept Analysis and Implications for Oncology Nurses*. 34(3), 673–681. <https://doi.org/10.1188/07.ONF.673-680>
- Gerasimoula, K., Lefkothea, L., Maria, L., Victoria, A., Paraskevi, T., & Maria, P. (2015). Quality of Life in Hemodialysis Patients. *Mater Sociomed*, 305–309.
- Green, J. A., Mor, M. K., Shields, A. M., Sevick, M. A., Palevsky, P. M., Fine, M. J., Arnold, R. M., & Weisbord, S. D. (2011). Prevalence and Demographic and Clinical Associations Hemodialysis. *Clinical Journal of the American Society of Nephrology*, 6(June), 1354–1360.
<https://doi.org/10.2215/CJN.09761110>
- Hajghanbari, B., Holsti, L., Road, J. D., & Reid, W. D. (2012). Pain in people with chronic obstructive pulmonary disease (COPD). *Respiratory Medicine*, 106(7), 998–1005. <https://doi.org/10.1016/j.rmed.2012.03.004>
- Haryani, S., & Misniarti. (2016). Masalah kebutuhan seksual pada pasien gagal ginjal kronik yang menjalani terapi haemodialisa. *Jurnal Media Kesehatan*, 9(1), 001-113.
- Heidarzadeh, M., Atashpeikar, S., & Jalilazar, T. (2010). Relationship between quality of life and self-care ability in patients receiving hemodialysis. *Iranian Journal of Nursing and Midwifery Research*, 15(2), 71–76.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3093176/>
- Herman, I., Yonata, A., Tjiptaningrum, A., Berawi, K. N., Kedokteran, F., Lampung, U., Ilmu, B., Dalam, P., Kedokteran, F., Lampung, U., Klinik, B. P., Kedokteran, F., Lampung, U., Fisiologi, B., Kedokteran, F., Lampung, U., & Moeloek, A. (2015). *Hubungan Lama Hemodialisis dengan Fungsi Kognitif Pasien Penyakit Ginjal Kronik yang Menjalani Hemodialisis di RSUD Abdul Moeloek Bandar Lampung The Relationship between Duration of Hemodialysis and Cognitive Function Chronic Kidney Disease Patient in Hem*. 7, 47–53.
- Hidayat, R., Azmi, S., & Pertiwi, D. (2012). *Artikel Penelitian Hubungan Kejadian Anemia dengan Penyakit Ginjal Kronik pada Pasien yang Dirawat di Bagian Ilmu Penyakit Dalam RSUP dr M Djamil Padang Tahun 2010*. 5(3), 546–550.
- Hintistan, S., & Deniz, A. (2018). Evaluation of Symptoms in Patients Undergoing Hemodialysis. *Bezmialem Science*, 6(2), 112–118.
<https://doi.org/10.14235/bs.2018.1530>
- Ikee, R., Yano, K., & Tsuru, T. (2019). Constipation in chronic kidney disease: It is time to reconsider. *Renal Replacement Therapy*, 5(1), 1–10.
<https://doi.org/10.1186/s41100-019-0246-3>
- Imelda, F., Susalit, E., Marbun, M. B. M., & Rumende, C. M. (2017). Gambaran



Klinis dan Kualitas Hidup Pasien Penyakit Ginjal Tahap Akhir yang Menjalani Hemodialisis Dua Kali Dibandingkan Tiga Kali Seminggu. *Jurnal Penyakit Dalam Indonesia*, 4(3), 128. <https://doi.org/10.7454/jpdi.v4i3.139>

International Society of Nephrology. (2012). *KDIGO 2012: Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease* (G. Eknayan & N. Lameire (eds.); Vol. 3, Issue 1).
<https://doi.org/10.3182/20140824-6-za-1003.01333>

Isroin, L. (2014). Manajemen Cairan pada Pasien Hemodialisis Untuk Meningkatkan Kualitas Hidup di RSUD Dr. Harjono Ponorogo. *IJNP (Indonesian Journal of Nursing Practices)*, 1(2), 146–156.
<http://journal.umy.ac.id/index.php/ijnp/article/view/655/807>

Joshi, V. D., Mooppil, N., & Lim, J. F. (2010). Validation of the Kidney Disease Quality of Life-Short Form: a cross-sectional study of a dialysis-targeted health measure in Singapore. *BMC Nephrology*, 11(1), 36.
<https://doi.org/10.1186/1471-2369-11-36>

Kamil, A. R., & Setiyono, E. (2018). *Symptom burden* dan Kualitas Tidur pada Pasien Hemodialysis. *Indonesian Journal of Nursing Sciences and Practice*, 27–37.

Kemenkes, R. (2017). *InfoDATIN: Situasi Penyakit Ginjal Kronis di Indonesia*. www.depkes.go.id › download › infodatin › infodatin ginjal 2017

Kemenkes, R. (2018). Hasil Utama Rikesdas. *Journal of Physics A: Mathematical and Theoretical*, 44(8), 085201. <https://doi.org/10.1088/1751-8113/44/8/085201>

Kirkova, J., Davis, M. P., Walsh, D., Tiernan, E., Leary, N. O., Legrand, S. B., Lagman, R. L., & Russell, K. M. (2019). *Cancer Symptom Assessment Instruments : A Systematic Review*. 24(9).
<https://doi.org/10.1200/JCO.2005.02.8332>

Krol, G. D. (2011). Chronic Kidney Disease Staging and Progression. In *Chronic Kidney Disease (CKD)* (6.0, p. 4). Henry Ford Health System.

Kubrusly, M., Oliveira, C. M. C. de, Santos, D. C. de O., Mota, R. S., & Pereira, M. L. (2012). Análise comparativa entre a albumina pré- e pós-dialise como indicadores do risco nutricional e de morbimortalidade em hemodiálise. *Jornal Brasileiro de Nefrologia*, 34(1), 27–35.
<https://doi.org/10.1590/S0101-28002012000100005>

Lakshmi, B., Kumar, A. C. V., Reddy, H., Gopal, J., Chaitanya, V., Chandra, V., Sandeep, P., Nagaraju, R., Ram, R., & Kumar, V. (2017). Employment status of patients receiving maintenance dialysis – peritoneal and hemodialysis: A cross-sectional study. *Indian Journal of Nephrology*, 27(5), 384.
https://doi.org/10.4103/ijn.IJN_151_16

Li, H., Xie, L., Yang, J., & Pang, X. (2018). International Journal of Nursing



Sciences *Symptom burden* amongst patients suffering from end-stage renal disease and receiving dialysis : A literature review. *International Journal of Nursing Sciences*, 5(4), 427–431. <https://doi.org/10.1016/j.ijnss.2018.09.010>

Mahdavi-mazdeh, M., Nozari, B., Hatmi, Z. N., Zamyadi, M., & Mahdavi, A. (2009). The Impact of Gender and Marital Status on Therapeutic Outcomes of Maintenance Hemodialysis Patients. *International Journal Nephrology and Urology*, 1(2), 124–128.

Manns, B. J., Johnson, J. A., Taub, K., Mortis, G., Ghalli, W. A., & Donaldson, C. (2002). Dialysis Adequacy and Health Related Quality of Life in. *ASAIO Journal, March*, 565–569.
<https://doi.org/10.1097/01.MAT.0000026361.66193.17>

Mendonça, A. E. O. de, Dantas, J. da G., Andrade, D. A. de, Segato, C. T., & Torres, G. de V. (2015). Socio-Demographic and Clinical Profile of Older Adults Receiving Hemodialysis. *Cogitare Enferm*, 20(1), 60–66.

Milik, A., & Hrynkiewicz, E. (2014). On translation of LD, IL and SFC given according to IEC-61131 for hardware synthesis of reconfigurable logic controller. *IFAC Proceedings Volumes (IFAC-PapersOnline)*, 19(1), 4477–4483.

Moghadasian, S., Hasan, M., Hagh, S., & Hokmabadi, L. A. (2012). *Lifestyle of Hemodialysis Patients in Comparison with Outpatients*. 1(2), 101–107.
<https://doi.org/10.5681/jcs.2012.015>

Muntina, D. V. (2019). *Pengaruh Pendidikan Kesehatan Terstruktur Terhadap Kualitas Hidup pada Pasien Hemodialisis di Rumah Sakit Akademik UGM*. Universitas Gadjah Mada.

Murphy, E. L., Murtagh, F. E. M., Carey, I., & Sheerin, N. S. (2008). Understanding Symptoms in Patients with Advanced Chronic Kidney Disease Managed without Dialysis: Use of a Short Patient-Completed Assessment Tool. *Nephron Clinical Practice*, 111(1), c74–c80.
<https://doi.org/10.1159/000183177>

Mushi, L., Marschall, P., & Fleßa, S. (2015). The cost of dialysis in low and middle-income countries : a systematic review. *BMC Health Services Research*, 1–10. <https://doi.org/10.1186/s12913-015-1166-8>

Nugroho, P. (2017). Apakah Hemodialisis Tiga Kali Seminggu Lebih Baik? *Jurnal Penyakit Dalam Indonesia*, 4(3), 103.
<https://doi.org/10.7454/jpdi.v4i3.142>

Pernefri. (2014). *10 th Report Of Indonesian Renal Registry 2017 10 th Report Of Indonesian Renal Registry 2017*.

Rahman, A. R. A., Rudiansyah, M., & Triawanti. (2020). Hubungan Antara Adekuasi Hemodialisis Dan Kualitas Hidup Pasien Di Rsud Ulin Banjarmasin : Tinjauan Terhadap Pasien Penyakit Ginjal Kronik Yang



UNIVERSITAS
GADJAH MADA

Gambaran Symptom Burden pada Pasien yang Menjalani Terapi Hemodialisis di Rumah Sakit TNI Angkatan

Udara Dr. Suhardi Hardjolukito Yogyakarta

AMALIA LATIFA A, Melyza Perdana, S.Kep., Ns., MS; Dr. Heny Suseani P., S.Kp., M.Kes.

Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Menjalani Hemodialisis Rutin. *Berkala Kedokteran*, 9(2), 151–160.
<https://doi.org/10.20527/jbk.v9i2.945>

Rahman, M., & Chowdhury, A. S. (2018). *Measuring the Health-Related Life Quality of Patients with Chronic Kidney Disease Title.*

Ratnasari, S., & Suleeman, J. (2017). Perbedaan Regulasi Emosi Perempuan dan Laki-Laki di Perguruan Tinggi. *Jurnal Psikologi Sosial*, 15(1), 35–46.
<https://doi.org/10.7454/jps.2017.4>

Rosdiana, I., Yetty, K., & Sabri, L. (2014). *BERHUBUNGAN DENGAN KEJADIAN INSOMNIA PADA PASIEN GAGAL GINJAL KRONIK Pendahuluan Metode Hasil.* 17(2), 39–47.

Rostami, Z., Einollahi, B., Lessan-pezeshki, M., Na-, A. S., Kebar, S. M., Shahbazian, H., Makhlough, A., Makhdoomi, K., Salesi, M., & Jalalzadeh, M. (2013). *Health-Related Quality of Life in Hemodialysis Patients : An Iranian Multi-Center Study.* 5(4), 901–912.

Ryan, T., Ingleton, C., Gardiner, C., Parker, C., Gott, M., & Noble, B. (2013). *Symptom burden*, palliative care need and predictors of physical and psychological discomfort in two UK hospitals. *BMC Palliative Care*, 12(1), 11. <https://doi.org/10.1186/1472-684X-12-11>

Senanayake, S., Gunawardena, N., Palihawadana, P., Bandara, P., & Haniffa, R. (2017). *Symptom burden in chronic kidney disease ; a population based cross sectional study.* 1–8. <https://doi.org/10.1186/s12882-017-0638-y>

Shdaifat, E. A., & Manaf, M. R. (2013). Haemodialysis in Jordan : Socio-Demographic Data , Clinical Analysis and Projecting Burden. *Middle-East Journal of Scientific Research*, 14(January), 624–632.
<https://doi.org/10.5829/idosi.mejsr.2013.14.5.73162>

Smeltzer, S. C., & Bare, B. G. (2013). *Buku Ajar Keperawatan Medikal Bedah* (2nd ed.). EGC.

Stokes, J. B. (2010). *CONSEQUENCES OF FREQUENT HEMODIALYSIS : COMPARISON TO CONVENTIONAL HEMODIALYSIS AND TRANSPLANTATION Years General Transplant.* 122, 124–136.

Sumardjito, S. (1999). Permasalahan Perkotaan Dan Kecenderungan Perilaku Individualis Penduduknya. *Jurnal Cakrawala Pendidikan*, 2(2).

Susianti, N. A., Astuti, Asmedi, A., Abdul, G., & Djarnwoto, B. (2015). Profil Fungsi Kognitif Pasien gagal Ginjal Kronik yang Menjalani Hemodialisis. *Jurnal Neurona*, 32(4). <http://www.neurona.web.id/paper-detail.do?id=945>

Susilawati, E., Latief, K., & Khomarudin, K. (2018). Efikasi Diri Dan Dukungan Sosial Pasien Hemodialisa Dalam Meningkatkan Kepatuhan Pembatasan Cairan. *Faletehan Health Journal*, 5(1), 39–48.
<https://doi.org/10.33746/fhj.v5i1.7>



Tandi, dkk, 2014 dalam Hasneli, Y., & Bayhakki. (2017). Hubungan Lama Menjalani Hemodialisis dengan Inter-Dialytic Weight Gain (IDWG) pada Pasien Hemodialisis Long-Term Relationship In Hemodialysis With Inter-Dialytic Weight Gain (IDWG) pada Pasien Hemodialisis. *Jurnal Keperawatan Universitas Padjajaran*, 5(3), 242–248.
jkp.fkep.unpad.ac.id/index.php/jkp/article/view/646

Tania, F., & Diskusi, H. (2016). *Biaya dan Outcome Hemodialisis di Rumah Sakit Kelas B dan C Pendahuluan. 1.*

Titisari, A. (2017). *Hubungan antara dukungan sosial keluarga dengan resiliensi pada penyakit ginjal kronis yang menjalani terapi hemodialisis* [Universitas Muhammadiyah Surakarta]. <http://eprints.ums.ac.id/51233/11/a.pdf>

Tranmer, J. E., & Heyland, D. (2003). *Measuring the Symptom Experience of Seriously Ill Cancer and Noncancer Hospitalized Patients Near the End of Life with the Memorial Symptom Assessment Scale*. 25(5), 420–429.
[https://doi.org/10.1016/S0885-3924\(03\)00074-5](https://doi.org/10.1016/S0885-3924(03)00074-5)

Vivekanand, P., Dm, J., Garcia-garcia, P. G., Iseki, P. K., Li, Z., Naicker, P. S., Plattner, B., Saran, P. R., Wang, P. A. Y., & Yang, P. C. (2013). Chronic kidney disease : Global dimension and perspectives Global Kidney Disease 3 Chronic kidney disease : global dimension and perspectives. *The Lancet*, 382(9888), 260–272. [https://doi.org/10.1016/S0140-6736\(13\)60687-X](https://doi.org/10.1016/S0140-6736(13)60687-X)

Wang, R., Tang, C., Chen, X., Zhu, C., Feng, W., Li, P., & Lu, C. (2016). Poor sleep and reduced quality of life were associated with symptom distress in patients receiving maintenance hemodialysis. *Health and Quality of Life Outcomes*, 14(1), 1–8. <https://doi.org/10.1186/s12955-016-0531-6>

Weisbord, S. D., Fried, L. F., Arnold, R. M., Rotondi, A. J., Fine, M. J., Levenson, D. J., & Switzer, G. E. (2004). Development of a symptom assessment instrument for chronic hemodialysis patients: The dialysis symptom index. *Journal of Pain and Symptom Management*, 27(3), 226–240. <https://doi.org/10.1016/j.jpainsymman.2003.07.004>

Wright, S. (2009). *Chapter 20 : Hemodialysis in Elderly Patients*. 1–4.

Yarlas, A. S., White, M. K., & Saris-baglama, M. Y. R. N. (2011). *Measuring the health status burden in hemodialysis patients using the SF-36 Ò health survey*. 383–389. <https://doi.org/10.1007/s11136-010-9764-8>

Yu, I. C., Huang, J. Y., & Tsai, Y. F. (2012). Symptom cluster among hemodialysis patients in Taiwan. *Applied Nursing Research*, 25(3), 190–196. <https://doi.org/10.1016/j.apnr.2010.11.002>