

- Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>
1. Muysoms FE, Antoniou SA, Bury K, et al. European Hernia Society guidelines on the closure of abdominal wall insisions. *Hernia*. 2015;19(1):1-24. doi:10.1007/s10029-014-1342-5
 2. Lau FH, Pomahac B. Wound healing in acutely injured fasia. *Wound Repair Regen*. 2014;22:14-17. doi:10.1111/wrr.12165
 3. Höer JJ, Junge K, Schachtrupp A, Klinge U, Schumpelick V. Influence of laparotomy closure technique on collagen synthesis in the insisional region. *Hernia*. 2002;6(3):93-98. doi:10.1007/s10029-002-0070-4
 4. Jairam AP, Timmermans L, Eker HH, et al. Prevention of insisional hernia with prophylactic onlay and sublay mesh reinforcement versus primary suture only in midline laparotomies (PRIMA): 2-year follow-up of a multicentre, double-blind, randomised controlled trial. *Lancet*. 2017;390(10094):567-576. doi:10.1016/S0140-6736(17)31332-6
 5. Fortelny RH, Baumann P, Thasler WE, et al. Effect of suture technique on the occurrence of insisional hernia after elective midline abdominal wall closure: Study protocol for a randomized controlled trial. *Trials*. 2015;16(1):1-8. doi:10.1186/s13063-015-0572-x
 6. Millbourn D. Effect of Stitch Length on Wound Complications After Closure of Midline Insisions_{title>A Randomized Controlled Trial}<alt-title>Stitch Length and Midline Insision Complications</alt-title>. *Arch Surg*. 2009;144(11):1056. doi:10.1001/archsurg.2009.189
 7. Radu P, Brătucu M, Garofil D, et al. Molecular factors of failure in insisional hernia surgery. *Chirurgia (Bucur)*. 2013;108(2):193-198. <http://www.ncbi.nlm.nih.gov/pubmed/23618569>.
 8. Kibe Y, Takenaka H, Kishimoto S. Spatial and temporal expression of basic fibroblast growth factor protein during wound healing of rat skin. 2000:720-727.
 9. Kreszinger M, Delimar D, Kos J, et al. Wound strength after midline laparotomy: a comparison of four closure techniques in rats. *Vet Arh*. 2007;77(5):397-408.
 10. Dart a J, Dart CM. 6.636 - Suture Material: Conventional and Stimuli Responsive. *Compr Biomater*. 2011;7:573-587. doi:10.1016/B978-0-08-055294-1.00245-2
 11. Mayberry JC, Burgess E a, Goldman RK, Pearson TE, Brand D, Mullins RJ.

- closure for trauma: the plain truth. *J Trauma*. 2004;57(1):157-162-163. doi:10.1097/01.TA.0000102411.69521.80
12. Nout E, Lange JF, Salu NE, et al. Creep Behavior of Commonly Used Suture Materials in Abdominal Wall Surgery. *J Surg Res*. 2007;138(1):51-55. doi:10.1016/j.jss.2006.06.001
13. Kirilova M, Stoytchev S, Pashkouleva D, Kavardzhikov V. Experimental study of the mechanical properties of human abdominal fascia. *Med Eng Phys*. 2011;33(1):1-6. doi:10.1016/j.medengphy.2010.07.017
14. Dahl M, Hansen P, Stål P, Edmundsson D, Magnusson SP. Stiffness and thickness of fascia do not explain chronic exertional compartment syndrome. *Clin Orthop Relat Res*. 2011;469(12):3495-3500. doi:10.1007/s11999-011-2073-x
15. Stewart RJ, Duley JA, Rosman I, Fraser R, Allardyce RA. the Wound Fibroblast and Macrophage .1. Wound Cell-Population Changes Observed in Tissue-Culture. *Br J Surg*. 1981;68(2):125-128.
16. Dubay DA, Wang X, Kuhn MA, Robson MC, Franz MG. The prevention of insisional hernia formation using a delayed-release polymer of basic fibroblast growth factor. *Ann Surg*. 2004;240(1):179-186. doi:10.1097/01.sla.0000131576.12153.ab
17. Franz MG, Smith PD, Wachtel TL, et al. fascia insisions heal faster than skin: A new model of abdominal wall repair. *Surgery*. 2001;129(2):203-208. doi:10.1067/msy.2001.110220
18. Skorupski P, Rechberger T, Postawski K, Woessner JF, Jakowicki JA. Is diminished pubocervical fascia collagen content a risk factor for failure of surgical management of genuine stress urinary incontinence in women? *Eur J Obstet Gynecol Reprod Biol*. 2002;102(2):195-198. doi:10.1016/S0301-2115(01)00587-5
19. Klinge U, Si ZY, Zheng H, Schumpelick V, Bhardwaj RS, Klosterhalfen B. Collagen I/III and matrix metalloproteinases (MMP) 1 and 13 in the fascia of patients with insisional hernias. *J Investig Surg*. 2001;14(1):47-54. doi:10.1080/089419301750072202
20. Koźma EM, Olczyk K, Głowacki A, Bobiński R. An accumulation of proteoglycans in scarred fascia. *Mol Cell Biochem*. 2000;203(1-2):103-112. <http://www.ncbi.nlm.nih.gov/pubmed/10724338>.

21. Hodgson NCF, Malthaner RA, Østbye T. The search for an ideal method of abdominal fascia closure: A meta- analysis. *Ann Surg.* 2000;231(3):436-442. doi:10.1097/00000658-200003000-00018
22. Knolmayer J, Cornell KM, Bowyer MW, Mccullough JS, Koenig W, Afb T. for fascia Healing. 1996;9610(96).
23. Gelse K, Pöschl E, Aigner T. Collagens - Structure, function, and biosynthesis. *Adv Drug Deliv Rev.* 2003;55(12):1531-1546. doi:10.1016/j.addr.2003.08.002
24. Hulmes DJS. Collagen diversity, synthesis and assembly. *Collagen Struct Mech.* 2008;15-47. doi:10.1007/978-0-387-73906-9_2
25. Eryilmaz E, Teizer W, Hwang W. In Vitro Analysis of the Co-Assembly of Type-I and Type-III Collagen. *Cell Mol Bioeng.* 2017;10(1):41-53. doi:10.1007/s12195-016-0466-3
26. Stadelmann WK, Digenis AG, Toin GR. Physiology and healing dynamics of chronic cutaneous wounds. *Am J Surg.* 1998;176(2A suppl):26S-38S.