

DAFTAR PUSTAKA

- Abdalaziz, F.A., Ismail, El-Sherbini, S.A., Behairy, A., 2018. The use of bedside echocardiography for measuring cardiac index and systemic vascular resistance in pediatric patients with septic shock. *Rev. Bras. Ter. Intensiva* 30: 460–470. doi:10.5935/0103-507X.20180067
- Agustini, N.M.A., Wati, D.K., Suparyatha, I., 2018. The relationship between bacterial types and antibiotic resistance with the clinical outcomes of sepsis patients in Pediatric Intensive Care Unit at Sanglah Hospital Denpasar, Bali-Indonesia. *Indones. J. Biomed. Sci.* 12: 13–18. doi:10.15562/ijbs.v12i1.144
- Amaral, G., Bushee, J., Keynes, M., 2013. Uscom for predict fluid responsiveness. *J. Petrol.* 369: 1689–1699. doi:10.1017/CBO9781107415324.004
- Arikawa, S., Rollins, N., Newell, M., Becquet, R., 2016. Mortality risk and associated factors in HIV-exposed, uninfected children. *Trop. Med. Int. Heal.* 21: 720–734. doi:10.1111/tmi.12695
- Assadi, F., 2017. Passive leg raising: Simple and reliable technique to prevent fluid overload in critically ill patients. *Int. J. Prev. Med.* 8: 1–9. doi:10.4103/ijpvm.IJPVM_11_17
- Basu, S., Frank, L.H., Fenton, K.E., Sable, C.A., 2012. Two-dimensional speckle tracking imaging detects impaired myocardial performance in children with septic shock, not recognized by conventional echocardiography. *Pediatr. Crit. Care Med.* 13: 259–264. doi:10.1097/PCC.0b013e3182288445
- Beltramo, F., Mentee, J., Razavi, A., Khemani, R.G., Ross, P.A., 2016. Validation of an Ultrasound Cardiac Output Monitor as a Bedside Tool for Pediatric Patients. *Pediatr. Cardiol.* 37: 177–183. doi:10.1007/s00246-015-1261-y
- Biban, P., Gaffuri, M., Spaggiari, S., Zaglia, F., Serra, A., Santuz, P., 2012. Early recognition and management of septic shock in children. *Pediatr. Rep.* 4. doi:10.4081/pr.2012.e13
- Bonanno, F.G., 2012. Shock-A reappraisal: The holistic approach. *J. Emergencies, Trauma Shock* 5: 167–177. doi:10.4103/0974-2700.96487
- Bridges, E., 2013. Using functional hemodynamic indicators to guide fluid therapy. *Am. J. Nurs.* 113: 42–50. doi:10.1097/01.NAJ.0000429754.15255.eb
- Brierley, J., Carcillo, J.A., Choong, K., 2015. Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: *Am. Coll. Crit. Care Med.* 40: 1291–1296. doi:10.1097/CCM.0b013e31819323c6.Clinical
- Brierley, J., Peters, M.J., 2008. Distinct hemodynamic patterns of septic shock at presentation to pediatric intensive care. *Pediatrics* 122: 752–759. doi:10.1542/peds.2007-1979
- Carcillo, J.A., Han, K., Lin, J., Orr, R., 2007. Goal-Directed Management of Pediatric Shock in the Emergency Department. *Clin. Pediatr. Emerg. Med.* 8: 165–175. doi:10.1016/j.cpem.2007.07.002
- Cattermole, Leung, P.Y.M., Ho, G.Y.L., Lau, P.W.S., Smith, B.E., 2017. The normal ranges of cardiovascular parameters measured using the ultrasonic cardiac output monitor. *Physiol. Rep.* 5: e13195. doi:10.14814/phy2.13195
- Chaiyakulsil, C., Chantira, M., Katanyuwong, P., 2018. Comparison of three non-

- invasive hemodynamic monitoring methods in critically ill children. *PLoS One* 13: 1–12. doi:10.1371/journal.pone.0199203
- Cheng, Y.W., Xu, F., Li, J., 2018. Identification of volume parameters monitored with a noninvasive ultrasonic cardiac output monitor for predicting fluid responsiveness in children after congenital heart disease surgery. *Med. (United States)* 97. doi:10.1097/MD.00000000000012289
- Chong, S.W., Peyton, P.J., 2012. A meta-analysis of the accuracy and precision of the ultrasonic cardiac output monitor (USCOM). *Anaesthesia* 67: 1266–1271. doi:10.1111/j.1365-2044.2012.07311.x
- Chris, N., 2019. Fluid Responsiveness. *Crit. Care* 35: 1–9.
- Cohen, J., 2002. The immunopathogenesis of sepsis. *Nature* 420: 885–891. doi:10.1038/nature01326
- Cole, E.T., Harvey, G., Urbanski, S., Foster, G., Thabane, L., 2014. Rapid paediatric fluid resuscitation: A randomised controlled trial comparing the efficiency of two provider-endorsed manual paediatric fluid resuscitation techniques in a simulated setting. *BMJ Open* 4. doi:10.1136/bmjopen-2014-005028
- Convertino, V.A., Wirt, M.D., Glenn, J.F., Lein, B.C., 2016. The compensatory reserve for early and accurate prediction of hemodynamic compromise: A review of the underlying physiology. *Shock* 45: 580–590. doi:10.1097/SHK.0000000000000559
- Danusantoso, M.M., 2016. Pengukuran Indeks Syok untuk Deteksi Dini Syok Hipovolemik pada Anak dengan Takikardia: telaah terhadap perubahan indeks isi sekuncup. *Sari Pediatr.* 15: 319. doi:10.14238/sp15.5.2014.319-24
- Davis, A.L., Carcillo, J.A., Nguyen, T.C., 2017. The American College of Critical Care Medicine Clinical Practice Parameters for Hemodynamic Support of Pediatric and Neonatal Septic Shock: Executive Summary. *Pediatr. Crit. Care Med.* 18: 884–890. doi:10.1097/PCC.0000000000001259
- Dellinger, R.P., Levy, M., 2013. Surviving sepsis campaign: International guidelines for management of severe sepsis and septic shock: 2012. *Crit. Care Med.* 41: 580–637. doi:10.1097/CCM.0b013e31827e83af
- Dey, I., Sprivilis, P., 2005. Emergency physicians can reliably assess emergency department patient cardiac output using the USCOM continuous wave Doppler cardiac output monitor. *Emerg Med Australas* 17: 193–199. doi:10.1111/j.1742-6723.2005.00722.x
- Dhanani, S., Barrowman, N.J., Ward, R.E., Murto, K.T., 2011. Intra- and inter-observer reliability using a noninvasive ultrasound cardiac output monitor in healthy anesthetized children. *Paediatr. Anaesth.* 21: 858–864. doi:10.1111/j.1460-9592.2010.03480.x
- Elgendy, A., Seppelt, I.M., Lane, A.S., 2017. Comparison of continuous-wave doppler ultrasound monitor and echocardiography to assess cardiac output in intensive care patients. *Crit. Care Resusc.* 19: 222–229.
- Fallis, A., 2013. SEPTIC SYOK. *J. Chem. Inf. Model.* 53: 1689–1699. doi:10.1017/CBO9781107415324.004
- Fathi, E.M., Narchi, H., Chedid, F., 2018. Noninvasive hemodynamic monitoring of septic shock in children. *World J. Methodol.* 8: 1–8. doi:10.5662/wjm.v8.i1.1

- Fisher, J.D., Nelson, D.G., Beyersdorf, H., Satkowiak, L.J., 2010. Clinical spectrum of shock in the pediatric emergency department. *Pediatr. Emerg. Care* 26: 622–625. doi:10.1097/PEC.0b013e3181ef04b9
- Gidwani, U.K., Mohanty, B., Chatterjee, K., 2013. The Pulmonary Artery Catheter. A Critical Reappraisal. *Cardiol. Clin.* 31: 545–565. doi:10.1016/j.ccl.2013.07.008
- Guarracino, F., Baldassarri, R., Pinsky, M.R., 2013. Ventriculo-arterial decoupling in acutely altered hemodynamic states. *Crit. Care* 17. doi:10.1186/cc12522
- Gwet, K.L., 2014. Handbook of inter-rater reliability: the definitive guide to measuring the extent of agreement among raters ; [a handbook for researchers, practitioners, teachers & students], 4. ed. ed. Advanced Analytics, LLC, Gaithersburg, MD.
- Hartawan, I.N.B., Pudjiadi, A.H., Latief, A., Dewi, R., Yuniar, I., 2016. Validitas Stroke Volume Variation dengan Ultrasonic Cardiac Output Monitor (USCOM) untuk Menilai Fluid Responsiveness. *Sari Pediatr.* 17: 367. doi:10.14238/sp17.5.2016.367-372
- Hodgson, L.E., Forni, L.G., Venn, R., Samuels, T.L., Wakeling, H.G., 2016. A comparison of the non-invasive ultrasonic cardiac output monitor (USCOM) with the oesophageal Doppler monitor during major abdominal surgery. *J. Intensive Care Soc.* 17: 103–110. doi:10.1177/1751143715610785
- Hodgson, L.E., Venn, R., Forni, L.G., Samuels, T.L., Wakeling, H.G., 2015. Measuring the cardiac output in acute emergency admissions: use of the non-invasive ultrasonic cardiac output monitor (USCOM) with determination of the learning curve and inter-rater reliability. *J. Intensive Care Soc.* 17: 122–128. doi:10.1177/1751143715619186
- Holley, A., Lukin, W., Paratz, J., Hawkins, T., Boots, R., Lipman, J., 2012. Review article: Part two: Goal-directed resuscitation - Which goals? Perfusion targets. *EMA - Emerg. Med. Australas.* 24: 127–135. doi:10.1111/j.1742-6723.2011.01515.x
- Horster, S., Stemmler, H.J., Strecker, N., Brettner, F., Geiger, S., 2012. Cardiac output measurements in septic patients: Comparing the accuracy of USCOM to PiCCO. *Crit. Care Res. Pract.* 2012. doi:10.1155/2012/270631
- Inwald, D.P., Canter, R., Woolfall, K., Mouncey, P., Harrison, D.A., 2019. Restricted fluid bolus volume in early septic shock: Results of the Fluids in Shock pilot trial. *Arch. Dis. Child.* 104: 426–431. doi:10.1136/archdischild-2018-314924
- Joffe, R., Duff, J., Garcia Guerra, G., Pugh, J., Joffe, A.R., 2016. The accuracy of blood pressure measured by arterial line and non-invasive cuff in critically ill children. *Crit. Care* 20: 177. doi:10.1186/s13054-016-1354-x
- Jozwiak, M., Monnet, X., Teboul, J.-L., 2018. Prediction of fluid responsiveness in ventilated patients. *Ann. Transl. Med.* 6: 352–352. doi:10.21037/atm.2018.05.03
- Kawasaki, T., 2017. Update on pediatric sepsis: A review. *J. Intensive Care* 5: 1–12. doi:10.1186/s40560-017-0240-1
- Kelm, D.J., Perrin, J.T., Cartin-Ceba, R., Kennedy, C.C., 2015. Fluid overload in patients with severe sepsis and septic shock treated with early goal-directed

- therapy is associated with increased acute need for fluid-related medical interventions and hospital death. *Shock* 43: 68–73. doi:10.1097/SHK.0000000000000268
- Knirsch, W., Kretschmar, O., Weiss, M., 2008. Cardiac output measurement in children: Comparison of the Ultrasound Cardiac Output Monitor with thermodilution cardiac output measurement. *Intensive Care Med.* 34: 1060–1064. doi:10.1007/s00134-008-1030-y
- Kruskal, W.H., Tanur, J.M., Williams, E.J., Stanley, J.C., 2009. International Encyclopedia of Statistics.
- Kurniyanta, P., Utariani, A., Hanindito, E., Ryalino, C., 2019. The ultrasonic cardiac output monitor (USCOM) as a tool in evaluating fluid responsiveness in pediatric patients underwent emergency surgery. *Bali J. Anaesth.* 31: 856–864. doi:10.1093/bja/aer090
- Latief, A., Hadinegoro, S.R.S., Pudjiadi, A.H., Malisie, R.F., Alam, A., 2016. Konsensus diagnosis dan tata laksana sepsis pada anak. *Badan Penerbit IDAI* 1–47.
- Lee, J.H., Kim, E.H., Jang, Y.E., Kim, H.S., Kim, J.T., 2019. Fluid responsiveness in the pediatric population. *Korean J. Anesthesiol.* 72: 429–440. doi:10.4097/kja.19305
- Lemson, J., Nusmeier, A., Van Der Hoeven, J.G., 2011. Advanced hemodynamic monitoring in critically ill children. *Pediatrics* 128: 560–571. doi:10.1542/peds.2010-2920
- Li, C., Lin, F.Q., Fu, S.K., Chen, G.Q., Li, Q., 2013. Stroke volume variation for prediction of fluid responsiveness in patients undergoing gastrointestinal surgery. *Int. J. Med. Sci.* 10: 148–155. doi:10.7150/ijms.5293
- Lichtenthal, P.R., Phillips, R.A., Sloniger, J.A., Copeland, J.G., 2006. USCOM- Non Invasive Doppler – Are Cardiac Output Measurements Accurate in Both Infants and Adults ? *Anesthesiology* 2006.
- Lukito, V., Djer, M.M., Pudjiadi, A.H., Munasir, Z., 2012. The role of passive leg raising to predict fluid responsiveness in pediatric intensive care unit patients. *Pediatr. Crit. Care Med.* 13: 155–160. doi:10.1097/PCC.0b013e3182388ab3
- Mackenzie, D.C., Noble, V.E., 2014. Assessing volume status and fluid responsiveness in the emergency department. *Clin. Exp. Emerg. Med.* 1: 67–77. doi:10.15441/ceem.14.040
- Madigan, V.M., Smith, B.E., 2013. Non-invasive method for rapid bedside estimation of inotropy: Theory and preliminary clinical validation. *Br. J. Anaesth.* 111: 580–588. doi:10.1093/bja/aet118
- Maitland, K., Kiguli, S., Olupot-olupot, P., Reyburn, H., Lang, T., 2011. Mortality after Fluid Bolus in African Children with Severe Infection. *N. Engl. J. Med.* 1959–1968. doi:10.1056/NEJMoa1404595
- Malbrain, M.L.N.G., Huygh, J., Peeters, Y., Bernards, J., 2016. Hemodynamic monitoring in the critically ill: An overview of current cardiac output monitoring methods. *F1000Research* 5: 1–9. doi:10.12688/f1000research.8991.1
- Marik, P.E., Monnet, X., Teboul, J.L., 2011. Hemodynamic parameters to guide fluid therapy. *Ann. Intensive Care* 1: 2–9. doi:10.1186/2110-5820-1-1

- Martin, K., Weiss, S.L., 2015. Initial resuscitation and management of pediatric septic shock. *Minerva Pediatr.* 67: 141–158.
- Mendelson, J., 2018. Management of Pediatric Shock. *pubmed* 36: 427–440.
- Monnet, X., Marik, P.E., Teboul, J.L., 2016. Prediction of fluid responsiveness: an update. *Ann. Intensive Care* 6: 1–11. doi:10.1186/s13613-016-0216-7
- Morelli, A., Ertmer, C., Westphal, M., Raffone, C., Venditti, M., 2013. Effect of heart rate control with esmolol on hemodynamic and clinical outcomes in patients with septic shock: A randomized clinical trial. *JAMA - J. Am. Med. Assoc.* 310: 1683–1691. doi:10.1001/jama.2013.278477
- Mukhtar, A., Awad, M., Elayashy, M., Hussein, A., 2019. Validity of mini-fluid challenge for predicting fluid responsiveness following liver transplantation. *BMC Anesthesiol.* 19: 4–9. doi:10.1186/s12871-019-0728-4
- Muller, L., Toumi, M., Bousquet, P.J., Riu-Poulenc, B., Louart, G., 2011. An increase in aortic blood flow after an infusion of 100 ml colloid over 1 minute can predict fluid responsiveness: The mini-fluid challenge study. *Anesthesiology* 115: 541–547. doi:10.1097/ALN.0b013e318229a500
- Nguyen, H.B., Banta, D.P., Stewart, G., Kim, T., Bansal, R., 2010. Cardiac index measurements by transcutaneous doppler ultrasound and transthoracic echocardiography in adult and pediatric emergency patients. *J. Clin. Monit. Comput.* 24: 237–247. doi:10.1007/s10877-010-9240-6
- Parker, M.M., 2014. Hemodynamic Management in Pediatric Septic Shock: Objective Tools May Complement Clinical Assessment*. *www.pccmjournal.org* 15: 90–91. doi:10.1097/01.pcc.0000436126.70886.e8
- Pasaribu, H., 2019. Mikrosirkulasi Pada Resusitasi Cairan. *J. Ilmu Kedokt.* 12: 63. doi:10.26891/jik.v12i2.2018.63-71
- Peake, S.L., Delaney, A., Bailey, M., Bellomo, R., Cameron, P.A., 2014. Goal-directed resuscitation for patients with early septic shock. *N. Engl. J. Med.* 371: 1496–1506. doi:10.1056/NEJMoa1404380
- Pigozzi, L., 2017. Evaluation of fluid responsiveness in circulatory shock. *Trauma Emerg. Care* 2: 1–2. doi:10.15761/tec.1000143
- Ross, P.A., Klein, M.J., Nguyen, T., 2019. Body Habitus and Risk of Mortality in Pediatric Sepsis and Septic Shock: A Retrospective Cohort Study. *J. Pediatr.* 210: 178–183. doi:10.1097/00005392-199503001-00016
- Rusmawatiningtyas, D., Nurnaningsih, N., 2017. Mortality rates in pediatric septic shock. *Paediatr. Indones.* 56: 304. doi:10.14238/pi56.5.2016.304-10
- Rusmawatiningtyas, D., Wiguna, P.A., Kumara, I.F., Arguni, E., Pudjiadi, A., 2019. Initial Hemodynamic Profiles of Children with Dengue Shock Syndrome in Referral Settings. *Am. J. Pediatr.* 5: 260–266. doi:10.11648/j.ajp.20190504.26
- Self, W.H., Semler, M.W., Steingrub, J.S., Douglas, I.S., Shapiro, N.I., 2018. Liberal Versus Restrictive Intravenous Fluid Therapy for Early Septic Shock: Rationale for a Randomized Trial. *Ann. Emerg. Med.* 72: 457–466. doi:10.1016/j.annemergmed.2018.03.039
- Smith, B.E., 2013. The USCOM and Haemodynamics A Guide for Junior Medical and Nursing Staff 15–20.
- Spenceley, N., MacLaren, G., Kissoon, N., MacRae, D.J., 2011. Monitoring in pediatric cardiac critical care: A worldwide perspective. *Pediatr. Crit. Care*

- Med.* 12. doi:10.1097/PCC.0b013e3182211d66
- Stapleton, L., 2015. Minimally invasive hemodynamic monitoring. *Hemodynamic Monit. Made Incred. Vis. Third Ed.* 1–6. doi:10.1007/978-3-642-00418-6_1895
- Stewart, G.M., Nguyen, H.B., Kim, T.Y., 2008. Inter-rater reliability for noninvasive measurement of cardiac function in children. *Pediatr. Emerg. Care* 24: 433–437. doi:<http://dx.doi.org/10.1097/PEC.0b013e31817de143>
- Toscani, L., Aya, H.D., Antonakaki, D., Arulkumaran, N., 2017. What is the impact of the fluid challenge technique on diagnosis of fluid responsiveness? A systematic review and meta-analysis. *Crit. Care* 21. doi:10.1186/s13054-017-1796-9
- Turnham, H., Brierley, J., 2015. New Methods of Monitoring Shock in Children. *Curr. Treat. Options Pediatr.* 1: 15–24. doi:10.1007/s40746-014-0009-x
- Ueyama, H., Kiyonaka, S., 2017. Predicting the Need for Fluid Therapy-Does Fluid Responsiveness Work? *J. Intensive Care* 5: 1–6. doi:10.1186/s40560-017-0210-7
- Vincent, J., Opal, S.M., Marshall, J.C., Tracey, K.J., 2013. Viewpoint Sepsis definitions : time for change 381: 774–775.
- Weiss, S.L., Peters, M.J., Inwald, D.P., Nadel, S., Brierley, J., Carcillo, J., 2020. Surviving sepsis campaign international guidelines for the management of septic shock and sepsis-associated organ dysfunction in children, *Intensive Care Medicine*. doi:10.1007/s00134-019-05878-6
- Wiguna, P.A., Rusmawatiningtyas, D., Indah Kartika Murni, 2019. Kesepakatan Antar Penilai Ultrasound Cardiac Output Monitoring Di Instalasi Rawat Intensif Anak. *Sari Pediatri*.
- Wulandari, A., Martuti, S., Kaswadi, P., 2018. Perkembangan diagnosis sepsis pada anak. *Sari Pediatr.* 19: 237. doi:10.14238/sp19.4.2017.237-44
- Yong Y. Han, M., Joseph A. Carcillo, M., Michelle A. 2015. Early Reversal of Pediatric-Neonatal Septic Shock by Community Physicians Is Associated With Improved Outcome University of Pittsburgh School of Medicine. *pediatrics.aappublications.org* 112: 458–461.
- Yuliarto, S., Kadafi, K., Anitasari, D., 2018. Restrictive versus Liberal Fluid Resuscitation in Children with Dengue Shock Syndrome: the differences in Clinical Outcomes and Hemodynamic Parameters. *Int. J. Pediatr.* 7: 9215–9224.
- Zorko, D.J., Choong, K., Gilleland, J., 2013. Urgent Ultrasound Guided Hemodynamic Assessments by a Pediatric Medical Emergency Team: A Pilot Study. *PLoS One* 8: 6–11. doi:10.1371/journal.pone.0066951