

## DAFTAR PUSTAKA

Anderson RCE, Kan P, Klimo P, Brockmeyer DL, Walker ML, Kestle JRW.  
Complication of intracranial monitoring in children with head trauma. *J Neurosurg*  
2004; 101 (2 Suppl): 53-8.

American College of Surgeons, 1997, *Advance Trauma Life Support*. United States  
of America: First Impression, 1997.

Arief Sjamsul, *Radikal Bebas*, Bagian Ilmu Kesehatan Anak Fakultas Kedokteran Universitas  
Airlangga Rumah Sakit dr Soetomo Surabaya, 2003: 1-9.

Arifianto M, Ma'ruf A, Ibrahim A. Efficacy comparison of mannitol and  
hypertonic saline for traumatic brain injury (TBI) treatment. *Bali Med J*. 2016;5:170–  
6. [Google Scholar]

Aloisi, F., Care, A., Borsellino, G., Gallo, P., Rosa, S., Bassani, A., *et al.* (1992).  
Production of hemolympopoietic cytokines (IL-6, IL-8, colony-stimulating factors) by  
normal human astrocytes in response to IL 1beta and tumor necrosis factor-  
alpha. *J. Immunol.* 149, 2358–2366.

Angeloni, C., Prata C., 2 Segal, F.V.D., Piperno R. and Hrelia. S. 2015, Traumatic Brain Injury  
and NADPH Oxidase: A Deep Relationship, *Oxidative Medicine and Cellular  
Longevity*, 2015: 1-10. Benveniste, E.N. (1998). Cytokine actions in the central nervous  
system. *Cytokine Growth Factor Rev.* 9, 259–275.

Bartnik, B.L., Lee, S.M., Hovda, D.A., and Sutton, R.L. (2007). The fate of glucose during the  
period of decreased metabolism after fluid percussion injury: a <sup>13</sup>C NMR study. *J.  
Neurotrauma* 24, 1079–1092. doi: 10.1089/neu.2006.0210

Bartnik, B.L., Sutton, R. L., Fukushima, M., Harris, N.G., Hovda, D.A., and Lee, S.M. (2005).  
Upregulation of pentose phosphate pathway and preservation of tricarboxylic acid cycle

Bisri T, Utomo BA, Fuadi I. Exogenous lactate infusion improved neurocognitive function of patients with mild traumatic brain injury. *Asian J Neurosurg.* 2016;11:151–9. [PMC free article] [PubMed] [Google Scholar]

Boone MD, Oren-Grinberg A, Robinson TM, Chen CC, Kasper EM. Mannitol or hypertonic saline in the setting of traumatic brain injury: What have we learned? *Surg Neurol Int.* 2015;6:177. [PMC free article] [PubMed] [Google Scholar]

Budi Harto Batubara, Nazaruddin Umar, Chairul M. Mursin, Perbandingan Osmolaritas Plasma Setelah Pemberian Manitol 20% 3 mL/kgBB dengan Natrium Laktat Hipertonik 3 mL/kgBB pada Pasien Cedera Otak Traumatik Ringan-Sedang, *Jurnal Anestesi Perioperatif [JAP]*. 2016;4(3): 154–61]

Brain Trauma Foundation. Guidelines for the Management of Severe Traumatic Brain Injury. BTF. 2007;24:S1-S106.

Brisby, H., Olmarker, K., Rosengren, L., Cederlund, C. G., and Rydevik, B. (1999). Markers of nerve tissue injury in the cerebrospinal fluid in patients with lumbar disc herniation and sciatica. *Spine* 24, 742–746.

Brough, D., Tyrrell, P. J., and Allan, S. M. (2011). Regulation of interleukin-1 in acute brain injury. *Trends Pharmacol. Sci.* 32, 617–622.

Brustovetsky, N., Brustovetsky, T., Jemmerson, R. and Dubinsky, J. M. 2002, Calcium induced cytochrome c release from CNS mitochondria is associated with the permeability transition and rupture of the outer membrane, *Journal of neurochemistry*, 80 (2): 207–18. Cernak, I. 2005, Animal Models of Head Trauma, *The Journal of the American Society for Experimental NeuroTherapeutics*, 2: 410–422.

- Carney N, Totten AM, O'Reilly C, Ullman JS, Hawryluk GW, Bell MJ, et al. Guidelines for the management of severe traumatic brain injury, fourth edition. *Neurosurgery*. 2017;80:6–15. [PubMed] [Google Scholar]
- Carpenter, K.L., Jalloh, I., Gallagher, C.N., Grice, P., Howe, D.J., Mason, A., et al. (2014). (13)C-labelled microdialysis studies of cerebral metabolism in TBI patients. *Eur.J.Pharm.Sci.* 57,87–97. doi:10.1016/j.ejps.2013.12.012
- Casey PA, Mckenna MC, Fiskum G, Saraswati M, Robertson Cl. Early and sustained alterations in cerebral metabolism after traumatic brain injury in immature rats. *J Neurotrauma*. 2008;25:603-14.
- Castro MA, Beltran FA, Brauchi S, Concha II. A metabolic switch in brain; glucose and lactate metabolism modulation by ascorbic acid. *J Neurochem*. 2009;110:423-40
- Catala-Temprano A, Teruel GC, Lasaosa FJC, Odena MP, Julian AN, Rico AP. Intracranial pressure and cerebral perfusion as risk factors in children with traumatic brain injuries. *J Neurosurg*. 2007;106(6 Suppl):463-6.
- Cernak, I. 2005, Animal Models of Head Trauma, *The Journal of the American Society for Experimental NeuroTherapeutics*, 2: 410–422.
- Chiry O, Fishbein WN, Merezhinskaya N, Clarke S, Galuske R, Magistretti PJ, et al. Distribution of the monocarboxylate transporter MCT2 in human cerebral cortex: An immunohistochemical study. *Brain Res*. 2008;1226:61-9.
- Chiry O, Pellerin L, Monnet-Tschudi F, Fishbein WN, Merezhinskaya N, Magistretti PJ, et al. Expression of the mono carboxylate transporter MCT1 in the adult human brain cortex. *Brain Res*. 2006;1070:65-70.
- Chung, I. Y., and Benveniste, E. N. (1990). Tumor necrosis factor-alpha production by astrocytes. Induction by lipopolysaccharide, IFN-gamma, and IL 1beta. *J. Immunol.* 144, 2999–3007. Coburn, K. (1992). Traumatic brain injury: the silent epidemic. *AACN Clin. Issues Crit.CareNurs*. 3, 9–18.

- Clayton TJ, Nelson RJ, Manara AR. Reduction in mortality from severe head injury following introduction of a protocol for intensive care management. *British Journal of Anaesthesia*; 2004. 93. h. 761–7.
- Coburn, K. (1992). Traumatic brain injury: the silent epidemic. *AACN Clin. Issues Crit.CareNurs.* 3, 9–18.
- Dahlan, S.M. 2011, *Statistik untuk Kedokteran dan Kesehatan: Deskriptif Bivariat, dan Multivariat*, Edisi 5, Salemba Medika, Jakarta.
- David A. Bender, PhD & Peter A. Mayes, PhD, DSc, Rodwell. *Harper's Illustrated Biochemistry*, 30<sup>th</sup> edition, section IV, Metabolism of Carbohydrates. 2015. 139-210
- De Waal Malefyt, R., Abrams, J., Bennett, B., Figdor, C. G., and De Vries, J.E. (1991). Interleukin-10 (IL-10) inhibits cytokine synthesis by human monocytes and an auto regulatory role of IL-10 produced by monocytes. *J. Exp.Med.* 174, 1209–1220.
- Dusick, J.R., Glenn, T.C., Lee, W.N., Vespa, P.M., Kelly, D.F., Lee, S.M., *et al.* (2007). Increased pentosephosphate pathway flux after clinical traumatic brain injury: a [1,2-<sup>13</sup>C] glucose labeling study in humans. *J.Cereb.BloodFlow Metab.* 27, 1593–1602. doi:10.1038/sj.jcbfm.9600458
- D'Andrea, A., Aste-Amezaga, M., Valiante, N. M., Ma, X., Kubin, M., and Trinchieri, G. (1993). Interleukin-10 (IL-10) inhibits human lymphocyte interferon gamma production by suppressing natural killer cell stimulatory factor/IL-12 synthesis in accessory cells. *J. Exp. Med.* 178, 1041–1048.
- Dikmen *et al*, 2009 Dikmen Sureyya S., PhD; John D. Corrigan, PhD, ABPP; Harvey S. Levin, PhD; Joan Machamer, MA; William Stiers, PhD, ABPP; Marc G. Weisskopf, PhD, ScD. 2009. Cognitive Outcome Following Traumatic Brain Injury. *J Head Trauma Rehabil.* Vol. 24, No. 6, pp. 430–438

*Annu.Rev. Immunol.* 27, 519–550.

Fan, L., Young, P. R., Barone, F. C., Feuerstein, G. Z., Smith, D. H., and McIntosh, T. K. (1995). Experimental brain injury induces expression of interleukin 1 beta RNA in the rat brain. *Brain Res.Mol.BrainRes.* 30, 125–130.

Fiorentino, D. F., Zlotnik, A., Mosmann, T. R., Howard, M., and O'Garra, A. (1991). IL-10 inhibits cytokine production by activated macrophages. *J. Immunol.* 147, 3815–3822.  
Foda, M. A., and Marmarou, A. (1994). A new model of diffuse brain injury in rats. Part II: morphological characterization. *J.Neurosurg.* 80, 301–313.

Gallagher, C.N., Carpenter, K.L., Grice, P., Howe, D.J., Mason, A., Timofeev, I., *et al.* (2009). The human brain utilizes lactate via the tricarboxylic acid cycle: a <sup>13</sup>C-labelled microdialysis and high – resolution nuclear magnetic resonance study. *Brain* 132 (Pt10), 2839–2849. doi:10.1093/brain/awp202

Goldstein, M. (1990). Traumatic brain injury: a silent epidemic. *Ann.Neurol.* 27, 327.

Gruber, M.F., Williams, C.C., and Gerrard, T.L. (1994). Macrophage-colony stimulating factor expression by anti-CD45 stimulated human monocytes is transcriptionally upregulated by IL-1beta and inhibited by IL-4 and IL-10. *J. Immunol.* 152, 1354–1361.

Hansson, M.J., Mansson, R., Morota, S., Uchino, H., Kallur, T., Sumi, T., *et al.* 2008, Calcium-induced generation of reactive oxygen species in brain mitochondria is mediated by permeability transition, *Free Radic Biol Med*, 45(3): 284–94.

Hanna, Ahmad MR. Effect of equiosmolar solutions of hypertonic sodium lactate versus mannitol in craniectomy patients with moderate traumatic brain injury. *Fakultas Kedokteran Universitas Hasanuddin. Med J Indonesia.* 2014;23;30–5.

Hickey Robert, Adelson David, *et al.* (2007) Cyclooxygenase-2 Activity Following Traumatic Brain Injury in the Developing Rat. 3, 271-276.

- Hutchinson, P. J., O'Connell, M. T., Seal, A., Nortje, J., Timofeev, I., Al-Rawi, P. G., *et al.* (2009). A combined microdialysis and FDG-PET study of glucose metabolism in head injury. *Acta Neurochir. (Wien)* 151, 51–61; discussion 61. doi:10.1007/s00701-008-0169-1
- Ide, K., Schmalbruch, I. K., Quistorff, B., Horn, A., and Secher, N. H. (2000). Lactate, glucose and O<sub>2</sub> uptake in human brain during recovery from maximal exercise. *J. Physiol.* 522 (Pt1), 159–164. doi:10.1111/j.1469-7793.2000.t01-2-00159.xm
- IONI, Pusat Informasi Obat Nasional, Bab 9.2.1 <http://pionas.pom.go.id/ioni/bab-9-gizi-dan-darah/92-cairan-dan-elektrolit/921-gangguan-keseimbangan-cairan-dan-elektrolit-4>,  
totilac
- Iskandar, Japardi, 2004, Penatalaksanaan Cedera Kepala *secara Operatif*. Sumatra Utara: USU Press 2004, 1-27.
- Jalloh, I., Helmy, A., Shannon, R. J., Gallagher, C. N., Menon, D. K., Carpenter, K. L., *et al.* (2013). Lactate uptake by the injured human brain: evidence from an arteriovenous gradient and cerebral microdialysis study. *J. Neurotrauma* 30, 2031–2037. doi:10.1089/neu.2013.2947
- Kaufmann WE, Worley PF, Taylor CV, Bremer M, Isakson PC 1997 Cyclooxygenase-2 expression during rat neocortical development and in Rett syndrome. *Brain Dev* 19:25–34
- Kamm, K., Vanderkolk, W., Lawrence, C., Jonker, M., and Davis, A. T. (2006). The effect of traumatic brain injury upon the concentration and expression of interleukin-1beta and interleukin-10 in the rat. *J. Trauma* 60, 152–157.
- Kinoshita, K., Chatzipanteli, K., Vitarbo, E., Truettner, J. S., Alonso, O. F., and Dietrich, W. D. (2002). Interleukin-1 beta messenger ribonucleic acid and protein levels after fluidpercussion brain injury in rats: importance of injury severity and brain temperature. *Neurosurgery* 51, 195–203; discussion 203.

Knoblach, S. M., and Faden, A. I. (1998). Interleukin-10 improves outcome and alters proinflammatory cytokine expression after experimental traumatic brain injury. *Exp. Neurol.* 153, 143–151.

Krueger, J. M. (2008). The role of cytokines in sleep regulation. *Curr. Pharm. Des.* 14, 3408–3416

Lu, B., Rutledge, B. J., Gu, L., Fiorillo, J., Lukacs, N. W., Kunkel, S. L. (1998). Abnormalities in monocyte recruitment and cytokine expression in monocyte chemoattractant protein 1-deficient mice. *J. Exp. Med.* 187, 601–608.

Lee *et al.*, 2004 Lee, L.L., Galo, E., Lyeth, B.G., Muizelaar, J.P. and Berman, R.F. 2004, Neuroprotection in the rat lateral fluid percussion model of traumatic brain injury by SNX-185, an N-type voltage-gated calcium channel blocker, *Exp Neurol*, 190: 70–78.

Levasseur JE, Alessandri B, Reinert M, Clausen T, Zhou Z, Altemeni N, dkk. Lactate, not glucose, up-regulates mitochondrial oxygen consumption both in sham and lateral fluid percussed rat brain. *J Neurosurg.* 2006;59(5):1122-31.

Lifshitz, J., Friberg, H., Neumar, R.W., Raghupathi, R., Welsh, F.A., Janmey, P., *et al.* 2003, Structural and functional damage sustained by mitochondria after traumatic brain injury in the rat: evidence for differentially sensitive populations in the cortex and hippocampus, *J Cereb Blood Flow Metab*, 23 (2): 219–31.

Marijata, 2006. Pengantar Dasar Bedah Klinis. Unit Pelayanan Kampus (UPK) Fakultas Kedokteran Universitas Gajah Mada. Yogyakarta. 2006: 83-108.

Mauritz W, Wilbacher I, Majdan M, *et al.* *Epidemiology, Treatment and Outcome of Patients after Severe Traumatic Brain Injury in European Regions with Different Economic Status.* *The European Journal of Public Health.* 2008;18:575-580.

Meierhans R, Brandi G, Fasshauer M, Sommerfeld J, Schupbach R, Bechir M, Stover

- J (2012) Arterial lactate above 2 mM is associated with increased brain lactate and decreased brain glucose in patients with severe traumatic brain injury. *Minerva Anestesiol* 78:185–193
- Moppet, I.K. 2007, Traumatic Brain Injury: Assessment, Resuscitation, and Early Management, *Br J Anaesth*, 99 (1): 18-31.
- Mustafa, A.G. and Al-Shboul, O. 2013, The role of free radicals and reactive species following traumatic brain injury, *OA Biotechnology*, 2 (3): 23.
- O'Connor, J.J., and Coogan, A. N. (1999). Actions of the proinflammatory cytokine IL-1 beta on central synaptic transmission. *Exp. Physiol.* 84, 601–614.
- Pellerin L. Lactate as a pivotal element in neuron–glia metabolic cooperation. *Neurochem Intern.* 2003;43 331-8.
- Qureshi AI, Suarez JI. Hypertonic saline solutions for treatment of intracranial hypertension. *Curr Opin Anaesthesiol.* 2000;5:3301-3.
- Rice, A.C., Zsoldos, R., Chen, T., Wilson, M.S., Alessandri, B., Hamm, R.J., *et al.* (2002). Lactate administration attenuates cognitive deficits following traumatic brain injury. *Brain Res.* 928, 156–159. doi:10.1016/S0006-8993(01)03299-1
- Rockswold GL, Solid CA, Paredes-Andrade E, Rockswold SB, Jancik JT, Quickel RR. Hypertonic saline and its effect on intracranial pressure, cerebral perfusion pressure, and brain tissue oxygen. *Neurosurgery.* 2009;65(6):1035-42.
- Rothwell, N. (2003). Interleukin-1 and neuronal injury: mechanisms, modification, and therapeutic potential. *BrainBehav.Immun.* 17, 152–157.
- Sai Ambika Tadepalli, Zsolt Kristóf Bali, Nóra Bruszt, Lili Veronika Nagy, Krisztina Amrein, Bálint Fazekas, András Büki, Endre Czeiter, István Hernádi. Long-term cognitive impairment without diffuse axonal injury following

Salmond C.H, D.A.Chatfield, D.K.Menon, J.D.Pickard, B.J.Sahakian. 2005. Cognitive sequelae of head injury: involvement of basal forebrain and associated structures. *Brain* Vol. 128 No.1: 189–200

Schurr A. Lactate: The ultimate cerebral oxidative energy substrate? *J Cereb Blood Flow Metab.* 2006;26:142–52. [PubMed] [Google Scholar]

Stiefel MF, Udoetuk JD, Spiotta AM, Gracias VH, Goldberg A, Maloney-Wilensky, dkk. Conventional neurocritical care and cerebral oxygenation after traumatic brain injury. *J Neurosurg.* 2006;105(5):68-75.

Timofeev,I.,Carpenter,K.L.,Nortje,J.,Al-Rawi,P.G.,O’Connell,M.T.,Czosnyka, M., *et al.*(2011a). Cerebral extra cellular chemistry and outcome following traumatic brain injury: a microdialysis study of 223 patients. *Brain* 134(Pt2), 484–494.doi: 10.1093/brain/awq353

Timofeev,I.,Czosnyka,M.,Carpenter,K.L.,Nortje,J.,Kirkpatrick,P.J.,Al-Rawi, P.G., *et al.* (2011b). Interaction between brain chemistry and physiology after traumatic brain injury :impact of autoregulation and microdialysis catheter location. *J.Neurotrauma* 28,849–860.doi:10.1089/neu.2010.1656

Timofeev,I.,Nortje,J.,Al-Rawi,P.G.,Hutchinson,P.J.,andGupta,A.K. (2013). Extra cellular brain pH with or without hypoxiaisa marker of profound metabolic derangement and increased mortality after traumatic brain injury. *J.Cereb.BloodFlowMetab.* 33,422–427.doi:10.1038/jcbfm.2012.186

Tyson,R.L.,Gallagher,C.,andSutherland,G.R.(2003).13C-Labeled substrates and the cerebral metabolic compartmentalization of acetate and lactate. *Brain Res.* 992,43–52.doi:10.1016/j.brainres.2003.08.027

vanHall, G., Stromstad, M., Rasmussen, P., Jans, O., Zaar, M., Gam, C., *et al.* (2009). Blood lactate is an important energy source for the human brain. *J.Cereb.BloodFlowMetab.* 29,1121–1129. doi:10.1038/jcbfm.2009.3

Winarsih, H. 2007, *Antioksidan Alami dan Radikal Bebas*, Edisi 5, Kanisius, Yogyakarta, pp 122-204.

Woiciechowsky, C., Schoning, B., Cobanov, J., Lanksch, W. R., Volk, H. D., and Docke, W. D. (2002). Early IL-6 plasma concentrations correlate with severity of brain injury and pneumonia in brain-injured patients. *J. Trauma* 52, 339–345

Woodcock Thomas, Cristina Maria, 2013, *The role of markers of inflammation in traumatic brain injury*, Fontiersin.org, Australia

Xiong, Y., Gu, Q., Peterson, P.L., Muizelaar, J.P., and Lee, C.P. (1997). Mitochondrial dysfunction and calcium perturbation induced by traumatic brain injury. *J.Neurotrauma* 14,23–34. doi:10.1089/neu.1997.14.23

Yan, E.B., Hellewell, S.C., Bellander, B. M., Agyapomaa, D. A., and Morganti Kossmann, M. C. (2011). Post-traumatic hypoxia exacerbates neurological deficit, neuro inflammation and cerebral metabolism in rats with diffuse traumatic brain injury. *J. Neuroinflammation* 8, 147

Ziebell dan Morganti-Kossmann 2010 Ziebell, J.M. and Morganti-Kossmann, M.C. 2010, Involvement of Pro- and Anti-Inflammatory Cytokines and Chemokines in the Pathophysiology of Traumatic Brain Injury, *The Journal of the American Society for Experimental NeuroTherapeutics*, 7: 22–30.