

## REFERENCES

- Agarwala, B. K., and P. R. Choudhury. 2013. Host races of the cotton aphid, *Aphis gossypii*, in asexual populations from wild plants of taro and brinjal . Journal of Insect Science 13(34): 1–13.
- Bertleff, D., J. Diekmann, S. Brand, A. G. Ünlü, and R. Bucher. 2020. Predation and avoidance behavior in aphid-ladybird interactions of native and invasive ladybirds in Europe. Ecological Entomology: 1-7.
- Binz, H., R. Bucher, M.H. Entling, and F. Menzel. 2014. Knowing the risk: crickets distinguish between spider predators of different size and commonness. Ethology 120(1): 99-110.
- Borowiak-Sobkowiak, B., R. Durak, and B. Wilkaniec. 2017. Morphology, biology and behavioral aspects of aphid *craccivora* (Hemiptera: Aphididae) on robinia pseudoacacia. Acta Sci. Pol. Hortorum Cultus 16(1): 39–49.
- Brisson, J. A., and D. L. Stern. 2006. The pea aphid, *Acyrtosiphon pisum*: an emerging genomic model system for ecological, developmental and evolutionary studies. BioEssays 28(7): 745-755.
- Bucher, R., H. Binz, F. Menzel, and M.H. Entling. 2014. Effects of spider chemotactile cues on arthropod behavior. J. Insect Behav 27: 567-580.
- CABI. 2019. *Acyrtosiphon pisum* (pea aphid). <<https://www.cabi.org/isc/datasheet/3147#totaxonomicTree>>. Accessed on 7 June 2020.
- CABI. 2019. *Aphis craccivora* (groundnut aphid). <<https://www.cabi.org/isc/datasheet/6192#totaxonomicTree>>. Accessed on 07 June 2020.
- CABI. 2019. *Aphis gossypii* (cotton aphid). <<https://www.cabi.org/isc/datasheet/6204#totaxonomicTree>>. Accessed on 07 June 2020.
- Dixon, A.F.G. 2000. Insect Predator-Prey Dynamics Ladybird Beetles & Biological Control. Cambridge University Press, United Kingdom.
- Evans, E. 2000. Morphology of invasion: body size patterns associated with establishment of *Coccinella septempunctata* (Coleoptera: Coccinellidae) in western North America. European Journal of Entomology 97: 469–474.
- Evans, E. W. 1991. Intra versus interspecific interactions of ladybeetles (Coleoptera: Coccinellidae) attacking aphids. Oecologia 87(3): 401–408.
- Fan, L. P., F. Ouyang, J. W. Su, and F. Ge. 2018. Adaptation of Defensive Strategies

by the Pea Aphid Mediates Predation Risk from the Predatory Lady Beetle.  
*Journal of Chemical Ecology* 44(1): 40–50.

- Fernandes, F. S., F. S. Ramalho, A. R. B. Nascimento, J. B. Malaquias, C. A. D. da Silva, and J. C. Zanuncio. 2012. Within-plant distribution of cotton aphid (hemiptera: aphididae), in cotton with colored fibers and cotton-fennel intercropping system. *Annals of the Entomological Society of America* 105(4): 599–607.
- Fill, A., E.Y. Long, and D.L.Finke. 2012. Non-consumptive effects of a natural enemy on a non-prey herbivore population. *Ecological Entomology* 37(1): 43-50.
- Francke, D. L., J. P. Harmon, C. T. Harvey, and A. R. Ives. 2008. Pea aphid dropping behavior diminishes foraging efficiency of a predatory ladybeetle. *Entomologia Experimentalis et Applicata* 127(2):118 – 124.
- Goławska, S. 2010. Effect of various host-plants on the population growth and development of the pea aphid. *Journal of Plant Protection Research* 50(2): 224–228.
- Hatano, E., G. Kunert, J.P. Michaud, and W.W. Weisser. 2008. Chemical cues mediating aphid location by natural enemies. *Eur. J. Entomol.* 105: 797-806.
- Hoki, E., Losey, J., Ugine, T.A., 2014. Comparing the consumptive and non-consumptive effects of a native and introduced lady beetle on pea aphids (*Acyrtosiphon pisum*). *Biological Control* 70: 78–84.
- Jean, P., and S. Jean-Christophe. 2010. The pea aphid complex as a model of ecological speciation. *Ecological Entomology* 35(s1):119 – 130.
- Kamphuis, L. G., L. Gao, and K. B. Singh. 2012. Identification and characterization of resistance to cowpea aphid (*Aphis craccivora* Koch) in *Medicago truncatula*. *BMC Plant Biology* 12(1): 1.
- Kanturski, M., Świętek, P., Trela, J., Borowiak-Sobkowiak, B., Wiczorek, K., 2020. Micromorphology of the model species pea aphid *Acyrtosiphon pisum* (Hemiptera, Aphididae) with special emphasis on the sensilla structure. *European Zoology Journal* 87(1): 336–356.
- Karl, K. and M. Poteser. 2009. Relationship between body size and spatial vision in the praying mantis - an ontogenic study. *Journal of Orthoptera Research* 18(2): 153-158.
- Kitajima, E. W., B. K. De Alcântara, P. M. Madureira, P. Alfenas-Zerbini, J. A. M. Rezende, and F. M. Zerbini. 2008. A mosaic of beach bean (*Canavalia rosea*) caused by an isolate of Cowpea aphid-borne mosaic virus (CABMV) in Brazil. *Archives of Virology* 153(4): 743–747.

- Koch, A. 1999. Cowpea Aphid. *Kansas Crop Pests*. 49–51.
- Koch, R. L. 2003. The multicolored Asian lady beetle, *Harmonia axyridis*: a review of its biology, uses in biological control, and non-target impacts. *Journal of Insect Science* 3(32): 1–16.
- Liu, X. D., T. T. Xu, and H. X. Lei. 2017. Refuges and host shift pathways of host-specialized aphids *Aphis gossypii*. *Scientific Reports* 7(1): 1–9.
- Minoretti, N., and W. W. Weisser. 2000. The impact of individual ladybirds (*Coccinella septempunctata*, Coleoptera: Coccinellidae) on aphid colonies. *European Journal of Entomology*, 97(4): 475–479.
- Mondor, E.B., D.S. Baird, K.N. Slessor, and B.D Roitberg. 2000. Ontogeny of alarm pheromone secretion in pea aphid, *Acyrtosiphon pisum*. *Journal of Chemical Ecology* 26(12): 2875-2882.
- Mondor, E.B and B.D. Roitberg. 2002. Pea aphid, *Acyrtosiphon pisum*, cornicle ontogeny as an adaptation to differential predation risk. *Canadian Journal of Zoology* 80: 2131-2136.
- Nelson, E. H., and J. A. Rosenheim. 2006. Encounters between aphids and their predators: The relative frequencies of disturbance and consumption. *Entomologia Experimentalis et Applicata* 118(3): 211–219.
- Ninkovic, V., Y. Feng, U. Olsson, J. Pettersson. 2013. Ladybird footprints induce aphid avoidance behavior. *Biological Control* 65: 63-71.
- Ofuya, T.I. 1997. Control of the cowpea aphid, *Aphis craccivora* Koch (Homoptera: Aphididae), in cowpea, *Vigna unguiculate* (L.) Walp. *Integrated Pest Management* 2: 199-207.
- Ostertagová, E., O. Ostertag, and J. Kovác. 2014. Methodology and application of the Kruskal-Wallis test. *Applied Mechanics and Materials* 611: 115-120.
- Purandare, S. R., and B. Tenhumberg. 2012. Influence of aphid honeydew on the foraging behaviour of *Hippodamia convergens* larvae. *Ecological Entomology* 37(3): 184–192.
- Preisser, E.L., D.I. Bolnick., and M.F. Benard. 2005. Scared to death? The effects of intimidation and consumption in predator -prey interactions. *Ecology* 86(2): 501-509.
- Preisser, E. and J. Orrock. 2012. The allometry of fear: interspecific relationships between body size and response to predation risk. *Ecosphere* 3(9): 1-27.
- Reynolds, P.G. and K. Cuddington. 2012. Effects of plant gross morphology on

- predator searching behaviour. *Environmental Entomology* 41(3): 516-522.
- Rice, K.B., Eubanks, M.D., 2013. No Enemies Needed: Cotton Aphids (Hemiptera: Aphididae) Directly Benefit from Red Imported Fire Ant (Hymenoptera: Formicidae) Tending. *Florida Entomol.* 96, 929–932.
- Sarwar, M.K., Azam, I., Iram, N., Iqbal, W., Rashda, A., Anwer, F., Atta, K., Ali, R., 2014. Cotton aphid *Aphis gossypii* L. (Homoptera; Aphididae); a challenging pest; biology and control strategies: A review. *International Journal of Applied Biology and Pharmaceutical Technology* 5(1): 288-294.
- Sun, Y. X., and T. X. Liu. 2018. *Harmonia axyridis* does not have obvious fitness gain and preference to the red morph of *Acyrtosiphon pisum*: A case study on a laboratory strain. *Frontiers in Ecology and Evolution* 6(1): 1–7.
- Tamai, K., and Y. Choh. 2018. Antipredator response of pea aphids *Acyrtosiphon pisum* (Hemiptera: Aphididae): effects of predation risks from an alternative patch on a current patch. *Applied Entomology and Zoology* 53(2): 267–274.
- Tamai, K., and Y. Choh. 2019. Previous exposures to cues from conspecifics and ladybird beetles prime antipredator responses in pea aphids 54(3): 277-283.
- Tayeh, A., A. Estoup, G. Laugier, A. Loiseau, J. Turgeon, S. Toepfer, and B. Facon. 2012. Evolution in biocontrol strains: insight from the harlequin ladybird *Harmonia axyridis*. *Evolutionary Applications* 5(5):481-488.
- Uiterwaal, S.F and J.P. DeLong. 2018. Multiple factors, including arena size, shape the functional responses of ladybird beetles. *Journal of Applied Ecology* 55(5): 2429-2438.
- Ünlü, A.G., Terlau, J.F., Bucher, R., 2020. Predation and avoidance behavior of the pea aphid *Acyrtosiphon pisum* confronted with native and invasive lady beetles in Europe. *Biological Invasions* 22(5): 1647-1656.
- Vonsteen, I., W.W. Weisser, and G. Kunert. 2016. Is there any evidence that aphid alarm pheromones work as prey and host finding kairomones for natural natural enemies?. *Ecological Entomology* 41(1): 1-12.
- Younes, A. 2006. Searching and predatory behavior of the predator, *Sphodromantis viridis* Forskal. *Egypt Egyptian Journal of Pest Control* 16(1): 33-37.
- Youren, F. 2012. Responses of aphids to semiochemicals released from ladybirds. *Sveriges Iantbruksuniversitet Swedish University of Agricultural Science. Department of Ecology, Sweden.*
- Zhang, R., B. Wang, G. Grossi, P. Falabella, Y. Liu, S. Yan, J. Lu, J. Xi, and G. Wong. 2017. Molecular basis of alarm pheromone detection in aphids. *Current Biology* 27(1): 55-61.