

DAFTAR PUSTAKA

- Abraham, V. A., Al-Shuaibi, M., Faleiro, J. R., Abozuhairah, R. A., & Vidyasagar, P. S. P. V. 1998. An integrated approach for the management of red palm weevil *Rhynchophorus ferrugineus* Oliv.: A key pest of date palm in the Middle East. *Sultan Qaboos University Journal for Scientific Research, Agricultural Science*, 3(1), 77–83.
- Astuti, M., Hafiza, Yuningsih, E., Nasution, I. M., Mustikawati, D., & Wasingun, A. R. 2014. *Pedoman Budidaya Kelapa (Cocos nucifera) Yang Baik*. Kementerian Pertanian Direktorat Jendral Perkebunan.
- Ávalos, J. A., Balasch, S., & Soto, A. 2016. Flight behaviour and dispersal of *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae) adults using mark-release-recapture method. *Bulletin of Entomological Research*, 106(5), 606–614.
- Azmi, W. A., Chong, :, Lian, J., Hazlina, :, Zakeri, A., Norhayati, :, Bayani, W., Omar, W., Yong, :, Wai, K., Ainatun, :, Zulkefli, N., & Hussain, M. H. 2017. The Red Palm Weevil, *Rhynchophorus ferrugineus*: Current Issues and Challenges in Malaysia. *Oil Palm Bulletin*, 74(August), 17–24.
- Azmi, W. A., Daud, S. N., Hussain, M. H., Wai, Y. K., Chik, Z., & Sajap, A. S. 2014. Field trapping of adult red palm weevil *rhynchophorus ferrugineus* olivier (coleoptera: Curculionidae) with food baits and synthetic pheromone lure in a coconut plantation. *Philippine Agricultural Scientist*, 97(4), 409–415.
- BAPEDA. 2009. *Profil Daerah Provinsi D.I. Yogyakarta*. Badan Perencanaan Pembangunan Daerah Provinsi Daerah Istimewa Yogyakarta.

Bouchard, P., Bousquet, Y., Davies, A. E., Alonso-Zarazaga, M. A., Lawrence, J. F., Lyal, C. H.

C., Newton, A. F., Reid, C. A. M., Schmitt, M., Ślipiński, S. A., & Smith, A. B. T. 2011.

Family-group names in Coleoptera (Insecta). *ZooKeys*, 88(SPEC. ISSUE), 1–972.

Chown, S. L., & Nicolson, S. W. 2004. *Insect Physiological Ecology : Mechanism and Patterns*.

Oxford University Press Inc.

Ditjenbun. 2009. Pengendalian Hama Kumbang Sagu (*Rhynchophorus ferrugineus* Oliver) pada Tanaman Kelapa.

<Http://Disbun.Jatimprov.Go.Id/Pustaka/Phocadownload/Hama%20kumbang%20sagu.Pdf>.

Diakses Pada 20 Januari 2020.

El-Mergawy, R., & Al-Ajlan, A. M. 2011. Red palm weevil, *rhynchophorus ferrugineus* (olivier):

Economic importance, biology, biogeography and integrated pest management. *Journal of Agricultural Science and Technology A*, 1(January), 1–23.

El-Sabea, A. M. R., Faleiro, J. R., & Abo-El-saad, M. M. 2009. The threat of red palm weevil

Rhynchophorus ferrugineus to date plantations of the Gulf region in the Middle-East: An economic perspective. *Outlooks on Pest Management*, 20(3), 131–134.

Grill, C. P., & Moore, A. J. 1998. Effects of a larval antipredator response and larval diet on adult

phenotype in an aposematic ladybird beetle. *Oecologia*, 114(2), 274–282.

Gunawardena, N. 1994. Electroantennogram Response of The Coconut Pest, *Rhynchophorus*

ferrugineus (Coleoptera : Curculionidae) To Alcohols. *J.Nant. Sci. Coun.*, 22(1), 25–33.

- Gunawardena, N., & Bandarage, U. 1995. 4-Methyl-5-Nonanol (Ferrugineol) as an aggregation pheromone of the coconut pest, *Rhynchophorus ferrugineus* F. (Coleoptera: Curculionidae)^osynthesis and use in a preliminary field assay. *Journal of the National Science Foundation of Sri Lanka*, 23(2), 71.
- Gunawardena, N. E., & Gunatilake, R. 1993. Preliminary Studies On A Host Attractant Of The Coconut Pest *Rhynchophorus ferrugineus* (Coleoptera : Curculionidae). *J. Natn. Sci. Coun.*, 21(1), 93–101.
- Hallett, R. H., Crespi, B. J., & Borden, J. H. 2004. Synonymy of *Rhynchophorus ferrugineus* (Olivier), 1790 and *R. vulneratus* (Panzer), 1798 (Coleoptera, Curculionidae, Rhynchophorinae). *Journal of Natural History*, 38(22), 2863–2882.
- Hallett, R. H., Gries, G., Gries, R., Borden, J. H., Czyzewska, E., Oehlschlager, A. C., Pierce, H. D., Angerilli, N. P. D., & Rauf, A. 1993. Aggregation pheromones of two asian palm Weevils, *Rhynchophorus ferrugineus* and *R. vulneratus*. *Naturwissenschaften*, 80(7), 328–331.
- Hoddle, M. S., Hoddle, C. D., Alzubaidy, M., Kabashima, J., Nisson, J. N., Millar, J., & Dimson, M. 2017. The palm weevil *Rhynchophorus vulneratus* is eradicated from Laguna Beach. *California Agriculture*, 71(1), 23–29.
- Hussain, A., Elsharabasy, S., Megahed, M., & Abd elmagid, A. 2016. Population Abundance of the Red Palm Weevil *Rhynchophorus ferrugineus* (oliv.) Adults on date Palm Plantations in Baharia Oases Giza Governorate - Egypt. *Journal of Plant Protection and Pathology*, 7(10), 649–654.

Indriyanti, D. R., Anggraeni, S. D., & Slamet, M. 2017. Density and composition of *Oryctes rhinoceros* (Coleoptera: Scarabaeidae) stadia in field. *ARPN Journal of Engineering and Applied Sciences*, 12(22), 6364–6371.

Jogjapro. 2014. Topografi Provinsi Daerah Istimewa Yogyakarta.
<https://jogjapro.go.id/berita/detail/topografi>. Diakses pada 30 Januari 2020

Justin, C. G. L., Leelamathi, M., Thangaselvabai, T., & Johnson, S. N. 2008. Bioecology and management of the red palm weevil, *Rhynchophorus ferrugineus* Oliv. (Coleoptera: Curculionidae) on coconut - a review. *Agricultural Reviews*, 29(2), 117–124.

Kaakeh, W. 2005. Longevity, fecundity, and fertility of the red palm weevil, *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) on natural and artificial diets. *Emirates Journal of Food and Agriculture*, 17(1), 23–33.

Kaakeh, W., El-Ezaby, F., Aboul-Nour, M. M., & Khamis, A. 2001. Management of The Red Palm Weevil, *Rhynchophorus Ferrugineus* OLIV., by a Pheromone/Food-Based Trapping System. *Proceedings of The Second International Conference on Date Palms*, 325–343.

Kalshoven, L. G. . 1981 *Pests Of Crops* (Revised an). P.T. Ichtar Baru - Van Hoeve.

Löhr, B., Vásquez-Ordóñez, A. A., & Lopez-Lavalle, L. A. B. 2015. *Rhynchophorus palmarum* in Disguise: Undescribed Polymorphism in the “black” Palm Weevil. *PLoS ONE*, 10(12), 1–14.

Manzoor, M., Ahmad, J. N., Ahmad, S. J. N., Naqvi, S. A., Umar, U. U. D., Rasheed, R., & Haider, M. S. 2020. Population dynamics, abundance and infestation of the red palm weevil, *rhynchophorus ferrugineus* (Olivier) in different geographical regions of date palm in Pakistan. *Pakistan Journal of Agricultural Sciences*, 57(2), 381–391.

Mardiatmoko, G., & Ariyanti, M. 2018. *Produksi Tanaman Kelapa (Cocos nucifera L.)*. Badan Penerbit Fakultas Pertanian Universitas Pattimura (BPFP - UNPATTI).

Metwally, H. A. A., & Basheer, A. M. 2019. *The Behavior and Activity of the Red Palm Weevil Rhynchophorus ferrugineus Throughout the Year under Baharia Oasis Conditions , Egypt*. 797–807.

Michie, L. J., Mallard, F., Majerus, M. E. N., & Jiggins, F. M. 2010. Melanic through nature or nurture: Genetic polymorphism and phenotypic plasticity in *Harmonia axyridis*. *Journal of Evolutionary Biology*, 23(8), 1699–1707.

Mizzi, S., Dandria, D., Mifsud, D., & Longo, S. 2009. The Red Palm Weevil, *Rhynchophorus ferrugineus* (Olivier, 1790) in Malta. *Bulletin of the Entomological Society of Malta*, 2(September), 111–121.

Murphy, S., & Briscoe, B. 1999. The red palm weevil as an alien invasive: Biology and the prospects for biological control as a component of IPM. *Biocontrol News and Information*, 20(1), 35–46.

Nuriyanti, D. D., Widhiono, I., & Suyanto, A. 2017. Faktor-Faktor Ekologis yang Berpengaruh terhadap Struktur Populasi Kumbang Badak (*Oryctes rhinoceros* L.). *Biosfera*, 33(1), 13.

Petren, K. 2001. Habitat and Niche, Concept of. *Encyclopedia of Biodiversity: Second Edition*, 3, 39–49. <https://doi.org/10.1016/B978-0-12-384719-5.00069-1>

Pracaya. 2007. *Hama dan Penyakit Tanaman Edisi Revisi*. Penebar Swadaya.

Price, P. W., Denno, R. F., Eubanks, M. D., Finke, D. I., & Kaplan, I. 2011. *Insect Ecology : Behaviour, Populations and Communities*. Cambridge University Press.

- Rugman-Jones, P. F., Hoddle, C. D., Hoddle, M. S., & Stouthamer, R. 2013. The Lesser of Two Weevils: Molecular-Genetics of Pest Palm Weevil Populations Confirm *Rhynchophorus vulneratus* (Panzer 1798) as a Valid Species Distinct from *R. ferrugineus* (Olivier 1790), and Reveal the Global Extent of Both. *PLoS ONE*, 8(10), 1–15.
- Sazali, S. N., Hazmi, I. R., Abang, F., Rahim, F., & Jemain, A. A. 2018. Morphometric study of the palm weevils, *rhynchophorus vulneratus* and *r. ferrugineus* (coleoptera: Curculionidae) in view of insular and mainland populations of Malaysia. *Pertanika Journal of Tropical Agricultural Science*, 41(3), 1329–1340.
- Sazali, S. N., Hazmi, I. R., Abang, F., Rahim, F., & Jemain, A. A. 2019. Population variation of the red stripe weevils, *Rhynchophorus vulneratus* (Coleoptera: Curculionidae) isolated by geographical limit. *Raffles Bulletin of Zoology*, 67(July), 378–384.
- Sivapragasm A, Ngalim A, Razali B, Sukaime MS. 2010. Field trapping of the adult red stripe weevil, *Rhynchophorus vulneratus* (Panzer) with an aggregation pheromone in a coconut ecosystem. *The Planter* 86: 173–180.
- Soroker, V., & Colazza, S. 2016. Handbook of Major Palm Pests: Biology and Management. *Handbook of Major Palm Pests: Biology and Management, December*, 1–317.
- Speight, M. R., Hunter, M. D., & Watt, A. D. 2008. *Ecology of Insects : Concepts and Application 2nd*. Wiley-Blackwell.
- Sugiyono. (2015). *Metode Penelitian Kuantitatif, Kualitatif, dan R&D*. Penerbit Alfabeta.

- Sukirno, S., Tufail, M., Rasool, K. G., & Aldawood, A. S. 2018a. Palm weevil diversity in Indonesia: Description of phenotypic variability in asiatic palm weevil, *rhynchophorus vulneratus* (Coleoptera: Curculionidae). *Journal of the Entomological Research Society*, 20(3), 1–22.
- Sukirno, S., Tufail, M., Rasool, K. G., & Aldawood, A. S. 2018b. Undescribed Color Polymorphism of the Asiatic Palm Weevil, *Rhynchophorus vulneratus* Panzer (Coleoptera: Curculionidae) in Indonesia: Biodiversity Study Based on COI Gene. *Florida Entomologist*, 101(4), 642.
- Tanyi Tambe, J., Riolo, P., Okolle, J. N., Isidoro, N., Fanciulli, P. P., & Dallai, R. 2013. Sexual size differences and colour polymorphism of *Rhynchophorus phoenicis* in the southwest region of Cameroon. *Bulletin of Insectology*, 66(1), 153–159.
- Tiglia, E. A., Vilela, E. F., Moura, J. I. L., & Anjos, N. 1998. Eficiência de armadilhas com feromônio de agregação e cana-de-açúcar na captura de *Rhynchophorus palmarum* (L.). *Anais Da Sociedade Entomológica Do Brasil*, 27(2), 177–183.
- Trisnadi, R. (2014). Kumbang Sagu (*Rhynchophorus* , sp) Penyebab Utama Kematian Pohon Kelapa. [Http://Dkpp.Probolinggakab.Go.Id/Wp-Content/Uploads/2018/02/Kumbang-Sagu2-1.Pdf](http://Dkpp.Probolinggakab.Go.Id/Wp-Content/Uploads/2018/02/Kumbang-Sagu2-1.Pdf) Diakses Pada 20 Januari 2020.
- Ul Haq, I., Shams, S., Khan, S., Khan, A., & Hameed, A. 2018. A novel report on morphological study of Red Palm Weevil (*Rhynchophorus ferrugineus*) from district Bannu KPK, Pakistan. *Cogent Food & Agriculture*, 4(1), 1–8.

Wattanapongsiri, A. 1966. a Revision of the Genera *Rhynchophorus* and *Dynamis* (Coleoptera: Curculionidae). *Department of Agriculture Science Bulletin*, 1(1)(June), 418.

Yulianto, Y., & Ernawati, D. 2015. Serangan *Rhynchophorus ferreginus* Di Wilayah Jawa Timur. *Buletin DITJENBUN Pertanian. BBPPT Surabaya*, 1–7.

Zverev, V., Kozlov, M. V., Forsman, A., & Zvereva, E. L. 2018. Ambient temperatures differently influence colour morphs of the leaf beetle *Chrysomela lapponica*: Roles of thermal melanism and developmental plasticity. *Journal of Thermal Biology*, 74(March), 100–109.