

DAFTAR PUSTAKA

- A, E., Utami, T. D. and Ramadan, D. (2017) 'Transfersomal gel containing green tea (*Camellia sinensis* L. kuntze) leaves extract: Increasing in vitro penetration', *Asian Journal of Pharmaceutical and Clinical Research*, 10(8), pp. 294–298. doi: 10.22159/ajpcr.2017.v10i8.19124.
- Ahmad, R. S. *et al.* (2014) 'Quantitative and Qualitative Portrait of Green Tea Catechins (Gtc) Through Hplc', *International Journal of Food Properties*, 17(7), pp. 1626–1636. doi: 10.1080/10942912.2012.723232.
- Ahmed, S. and Stepp, J. R. (2013) 'Green Tea: The Plants, Processing, Manufacturing and Production', *Elsevier Academic Press*. doi: 10.1016/B978-0-12-384937-3.00002-1.
- Ali, S. M. and Yosipovitch, G. (2013) 'Skin pH: from basic science to basic skin care', *Acta Dermato-Venereologica*, 93(3), pp. 261–267. doi: 10.2340/00015555-1531.
- Anwar, E., Inees, J. and Ramadan, D. (2018) 'Formulation of a Cream Containing Ethosomal Green Tea (*Camellia sinensis* L. Kuntze) Leaf Extracts for Improved Dermal Penetration', *International Journal of Applied Pharmaceutics*, 10(1), pp. 221–224. doi: 10.22159/ijap.2018.v10s1.49.
- Anwar, E., Ramadan, D. and Ardi, G. D. (2018) 'Novel Transethosome Containing Green Tea (*Camellia Sinensis* L. Kuntze) Leaf Extract for Enhanced Skin Delivery of Epigallocatechin Gallate: Formulation and In Vitro Penetration Test', *International Journal of Applied Pharmaceutics*, 10(1), pp. 299–302. doi: 10.22159/ijap.2018.v10s1.66.
- Anwar, E. and Rizkamiarty, S. (2020) 'Formulation and evaluation of cosmetic foundation using epigallocatechin gallate as a sun protection', *International Journal of Applied Pharmaceutics*, 12(Special Issue 1), pp. 130–134. doi: 10.22159/ijap.2020.v12s1.FF029.
- Archana, S. and Abraham, J. (2011) 'Comparative analysis of antimicrobial activity of leaf extracts from fresh green tea, commercial green tea and black tea on pathogens', *Journal of Applied Pharmaceutical Science*, 01(08), pp. 149–152.
- Aryani, R. *et al.* (2019) 'Effect of solid lipid nanoparticles system on the stability of Green Tea leaves (*Camellia sinensis* L. Kuntze) extract as sunscreen', *Journal of Physics: Conference Series*, 1375, p. 012078. doi: 10.1088/1742-6596/1375/1/012078.
- Avadhani, K. S. *et al.* (2017) 'Skin delivery of epigallocatechin-3-gallate (EGCG) and hyaluronic acid loaded nano-transfersomes for antioxidant and anti-aging effects in UV radiation induced skin damage', *Drug Delivery*, 24(1), pp. 61–74. doi: 10.1080/10717544.2016.1228718.

- Bae, J. *et al.* (2020) 'Activity of catechins and their applications', *Biomedical Dermatology*, 4(1), p. 8. doi: 10.1186/s41702-020-0057-8.
- Bartosikova, L. and Necas, J. (2018) 'Epigallocatechin gallate: A review', *Veterinarni Medicina*, 10, pp. 443–467. doi: 10.17221/31/2018-VETMED.
- Bhattacharya, S. and Sherje, A. P. (2020) 'Development of resveratrol and green tea sunscreen formulation for combined photoprotective and antioxidant properties', *Journal of Drug Delivery Science and Technology*, 60, p. 102000. doi: 10.1016/j.jddst.2020.102000.
- Bianchi, A., Marchetti, N. and Scalia, S. (2011) 'Photodegradation of (-)-epigallocatechin-3-gallate in topical cream formulations and its photostabilization', *Journal of Pharmaceutical and Biomedical Analysis*, 56(4), pp. 692–697. doi: 10.1016/j.jpba.2011.07.007.
- Chacko, S. M. *et al.* (2010) 'Beneficial effects of green tea: A literature review', *Chinese Medicine*, 5, p. 13. doi: 10.1186/1749-8546-5-13.
- Chamcheu, J. C. *et al.* (2018) 'Chitosan-based nanoformulated (-)-epigallocatechin-3-gallate (EGCG) modulates human keratinocyte-induced responses and alleviates imiquimod-induced murine psoriasiform dermatitis', *International Journal of Nanomedicine*, 13, pp. 4189–4206. doi: 10.2147/IJN.S165966.
- Chan, E. W. C. *et al.* (2011) 'Antioxidant and antibacterial properties of green, black, and herbal teas of *Camellia sinensis*', *Pharmacognosy Research*, 3(4), pp. 266–272. doi: 10.4103/0974-8490.89748.
- Chen, J. *et al.* (2017) 'Anti-skin-aging effect of epigallocatechin gallate by regulating epidermal growth factor receptor pathway on aging mouse model induced by d-Galactose', *Mechanisms of Ageing and Development*, 164, pp. 1–7. doi: 10.1016/j.mad.2017.03.007.
- Chang, R.-K. *et al.* (2012) 'Generic Development of Topical Dermatologic Products: Formulation Development, Process Development, and Testing of Topical Dermatologic Products', *The AAPS Journal*, 15(1), pp. 41–52. doi: 10.1208/s12248-012-9411-0.
- Chaturvedula, V. S. P. and Prakash, I. (2011) 'The aroma, taste, color and bioactive constituents of tea', *Journal of Medicinal Plants Research*, 5(11), pp. 2110–2124. doi: 10.5897/JMPR.9001187.
- Chaugule, A. *et al.* (2019) 'Extraction of Caffeine', *International Journal of Advanced Research in Chemical Science*, 6(9), pp. 11–19. doi: 10.20431/2349-0403.0609002.
- Domingo, D. S. *et al.* (2010) 'Anti-angiogenic effects of epigallocatechin-3-gallate in human skin', *International Journal of Clinical and Experimental Pathology*, 3(7), pp. 705–709.

- Dooley, A. *et al.* (2010) 'Modulation of collagen type I, fibronectin and dermal fibroblast function and activity, in systemic sclerosis by the antioxidant epigallocatechin-3-gallate', *Rheumatology (Oxford, England)*, 49(11), pp. 2024–2036. doi: 10.1093/rheumatology/keq208.
- Dube, A. *et al.* (2010) 'Effective use of reducing agents and nanoparticle encapsulation in stabilizing catechins in alkaline solution', *Food Chemistry*, 122(3), pp. 662–667. doi: 10.1016/j.foodchem.2010.03.027.
- Elcistia, R. and Zulkarnain, A. K. (2018) 'Formula Optimization of o/w Cream Combination Oxybenzone and Titanium Dioxide and Its In Vivo Activity Testing', *Majalah Farmaseutik*, 14(2), pp. 63–78. doi: 10.22146/farmaseutik.v14i2.42596.
- Fangueiro, J. F. *et al.* (2014) 'Validation of a high performance liquid chromatography method for the stabilization of epigallocatechin gallate', *International Journal of Pharmaceutics*, 475(1–2), pp. 181–190. doi: 10.1016/j.ijpharm.2014.08.053.
- Faramayuda, F., Alatas, F. and Desmiaty, Y. (2010) 'Formulasi Sediaan Losion Antioksidan Ekstrak Teh Hijau (*Camellia sinensis* L.)', *Majalah Obat Tradisional*, 15(3), pp. 105–111.
- Feng, H.-L. *et al.* (2014) 'Isolation and purification of condensed tannins from flamboyant tree and their antioxidant and antityrosinase activities', *Applied Biochemistry and Biotechnology*, 173(1), pp. 179–192. doi: 10.1007/s12010-014-0828-z.
- Ganceviciene, R. *et al.* (2012) 'Skin anti-aging strategies', *Dermato-endocrinology*, 4(3), pp. 308–319. doi: 10.4161/derm.22804.
- Garbossa, W. A. C. and Maia Campos, P. M. B. G. (2016) 'Euterpe oleracea, Matricaria chamomilla, and Camellia sinensis as promising ingredients for development of skin care formulations', *Industrial Crops and Products*, 83, pp. 1–10. doi: 10.1016/j.indcrop.2015.11.026.
- Gomes, E. C., Silva, A. N. and Oliveira, M. R. de (2012) *Oxidants, Antioxidants, and the Beneficial Roles of Exercise-Induced Production of Reactive Species, Oxidative Medicine and Cellular Longevity*. Hindawi. doi: <https://doi.org/10.1155/2012/756132>.
- Hong, Y.-H. *et al.* (2013) 'Tannase-converted green tea catechins and their anti-wrinkle activity in humans', *Journal of Cosmetic Dermatology*, 12(2), pp. 137–143. doi: 10.1111/jocd.12038.
- Hu, C. *et al.* (2015) 'Development and characterization of a self-double-emulsifying drug delivery system containing both epigallocatechin-3-gallate and α -lipoic acid', *Journal of Materials Science*, 50(20), pp. 6567–6577. doi: 10.1007/s10853-015-9194-7.

- Hu, C. *et al.* (2016) 'Transdermal solid delivery of epigallocatechin-3-gallate using self-double-emulsifying drug delivery system as vehicle: Formulation, evaluation and vesicle-skin interaction', *Journal of Biomaterials Applications*, 30(7), pp. 1080–1091. doi: 10.1177/0885328215617891.
- Huang, W.-Y. *et al.* (2013) 'Effects of Water Solutions on Extracting Green Tea Leaves', *The Scientific World Journal*, 2013, pp. 1–6. doi: <https://doi.org/10.1155/2013/368350>.
- Hussain, S. and Ashafaq, M. (2018) 'Epigallocatechin-3-Gallate (EGCG): mechanisms, perspectives and clinical applications in cervical cancer', *Journal of Cancer Prevention & Current Research*, 9(4), pp. 179–182. doi: 10.15406/jcpcr.2018.09.00345.
- Isnan, A. P. and Jufri, M. (2017) 'Formulation of Niosomal Gel Containing Green Tea Extract (*Camellia sinensis* L. Kuntze) Using Thin-Layer Hydration', *International Journal of Applied Pharmaceutics*, 9, pp. 38–43. doi: 10.22159/ijap.2017.v9s1.23_28.
- Jo, Y.-H. *et al.* (2012) 'Antioxidant, tyrosinase inhibitory, and acetylcholinesterase inhibitory activities of green tea (*Camellia sinensis* L.) seed and its pericarp', *Food Science and Biotechnology*, 21(3), pp. 761–768. doi: 10.1007/s10068-012-0099-9.
- Kaur, L. P. and Guleri, T. K. (2013) 'Topical Gel: A Recent Approach for Novel Drug delivery', *Asian Journal of Biomedical and Pharmaceutical Sciences*, 3(17), pp. 1–5.
- Kim, H.-L. *et al.* (2014) 'Promotion of full-thickness wound healing using epigallocatechin-3-O-gallate/poly (lactic-co-glycolic acid) membrane as temporary wound dressing', *Artificial Organs*, 38(5), pp. 411–417. doi: 10.1111/aor.12190.
- Khullar, R. *et al.* (2012) 'Formulation and evaluation of mefenamic acid emulgel for topical delivery', *Saudi pharmaceutical journal: SPJ: the official publication of the Saudi Pharmaceutical Society*, 20(1), pp. 63–67. doi: 10.1016/j.jsps.2011.08.001.
- Kumar, A. and Dixit, C. K. (2017) 'Methods for characterization of nanoparticles' *Advances in Nanomedicine for the Delivery of Therapeutic Nucleic Acids*. Elsevier, pp. 43–58. doi: 10.1016/B978-0-08-100557-6.00003-1.
- Koch, W. *et al.* (2020) 'The Role of Extracting Solvents in the Recovery of Polyphenols from Green Tea and Its Antiradical Activity Supported by Principal Component Analysis', *Molecules*, 25(9). doi: 10.3390/molecules25092173.
- Koch, W. *et al.* (2019) 'Applications of Tea (*Camellia sinensis*) and Its Active Constituents in Cosmetics', *Molecules*, 24(23). doi: 10.3390/molecules24234277.

- Lai-Cheong, J. E. and McGrath, J. A. (2017) 'Structure and function of skin, hair and nails', *Medicine*, 45(6), pp. 347–351. doi: 10.1016/j.mpmed.2017.03.004.
- Lambert, J. D. and Elias, R. J. (2010) 'The antioxidant and pro-oxidant activities of green tea polyphenols: a role in cancer prevention', *Archives of Biochemistry and Biophysics*, 501(1), pp. 65–72. doi: 10.1016/j.abb.2010.06.013.
- Lee, K. O., Kim, S. N. and Kim, Y. C. (2014) 'Anti-wrinkle Effects of Water Extracts of Teas in Hairless Mouse', *Toxicological Research*, 30(4), pp. 283–289. doi: 10.5487/TR.2014.30.4.283.
- Li, N. *et al.* (2012) 'Kinetic study of catechin stability: effects of pH, concentration, and temperature', *Journal of Agricultural and Food Chemistry*, 60(51), pp. 12531–12539. doi: 10.1021/jf304116s.
- Li, H. *et al.* (2016) 'Topical treatment of green tea polyphenols emulsified in carboxymethyl cellulose protects against acute ultraviolet light B-induced photodamage in hairless mice', *Photochemical & Photobiological Sciences*, 15(10), pp. 1264–1271. doi: 10.1039/C6PP00073H.
- M, S. *et al.* (2020) 'An overview on topical drug delivery system – Updated review', *International Journal of Research in Pharmaceutical Sciences*, 11(1), pp. 368–385. doi: 10.26452/ijrps.v11i1.1831.
- Mahmood, T., Akhtar, N. and Khan, B. A. (2010) 'The morphology, characteristics, and medicinal properties of *Camellia sinensis* tea', *Journal of Medicinal Plants Research*, 4(19), pp. 2028–2033. doi: 10.5897/JMPR10.010.
- Mahmood, T., Akhtar, N. and Moldovan, C. (2013) 'A comparison of the effects of topical green tea and lotus on facial sebum control in healthy humans', *Hippokratia*, 17(1), p. 64.
- Mahmood, T. *et al.* (2013) 'Fabrication, physicochemical characterization and preliminary efficacy evaluation of a W/O/W multiple emulsion loaded with 5% green tea extract', *Brazilian Journal of Pharmaceutical Sciences*, 49(2), pp. 341–349. doi: 10.1590/S1984-82502013000200016.
- Makvandi, P. *et al.* (2020) 'Natural Formulations Provide Antioxidant Complement to Hyaluronic Acid-Based Topical Applications Used in Wound Healing', *Polymers*, 12(8). doi: 10.3390/polym12081847.
- Malik, D. S., Mital, N. and Kaur, G. (2016) 'Topical drug delivery systems: a patent review', *Expert Opinion on Therapeutic Patents*, 26(2), pp. 213–228. doi: 10.1517/13543776.2016.1131267.
- Manea, A.-M., Vasile, B. S. and Meghea, A. (2014) 'Antioxidant and antimicrobial activities of green tea extract loaded into nanostructured lipid carriers', *Comptes Rendus Chimie*, 17(4), pp. 331–341. doi: 10.1016/j.crci.2013.07.015.

- McGrath, J. A. and Uitto, J. (2016) 'Structure and Function of the Skin', in *Rook's Textbook of Dermatology, Ninth Edition*. American Cancer Society, pp. 1–52. doi: 10.1002/9781118441213.rtd0002.
- Mereles, D. and Hunstein, W. (2011) 'Epigallocatechin-3-gallate (EGCG) for clinical trials: more pitfalls than promises?', *International Journal of Molecular Sciences*, 12(9), pp. 5592–5603. doi: 10.3390/ijms12095592.
- Min, K. and Kwon, T. K. (2014) 'Anticancer effects and molecular mechanisms of epigallocatechin-3-gallate', *Integrative Medicine Research*, 3(1), pp. 16–24. doi: 10.1016/j.imr.2013.12.001.
- Min, T. and Bartholomew, B. (2010) 'Theaceae', *Flora of China*. Available at: http://efloras.org/florataxon.aspx?flora_id=2&taxon_id=10882 (Accessed: 26 September 2020).
- Molyneux, P. (2004) 'The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant', *Songklanakarin Journal of Science and Technology (SJST)*, 26(2), pp. 211–219.
- Nadim, M. *et al.* (2014) 'Improvement of polyphenol properties upon glucosylation in a UV-induced skin cell ageing model', *International Journal of Cosmetic Science*, 36(6), pp. 579–587. doi: 10.1111/ics.12159.
- Namal Senanayake, S. P. J. (2013) 'Green tea extract: Chemistry, antioxidant properties and food applications – A review', *Journal of Functional Foods*, 5(4), pp. 1529–1541. doi: 10.1016/j.jff.2013.08.011.
- Namita, P., Mukesh, R. and Vijay, K. J. (2012) 'Camellia Sinensis (Green Tea): A Review', *Global Journal of Pharmacology*, 6(2), pp. 52–59.
- Nurwaini, S. and Sari, D. A. P. (2019) 'MASKER GEL PEEL-OFF EKSTRAK DAUN TEH HIJAU (*CAMELLIA SINENSIS* L.): SIFAT FISIK DAN AKTIVITAS ANTIOKSIDANNYA', *The 9th University Research Colloquium (Urecol)*, 9(1). Available at: <http://eproceedings.umpwr.ac.id/index.php/urecol9/article/view/587> (Accessed: 28 September 2020).
- Pai, V. V., Shukla, P. and Kikkeri, N. N. (2014) 'Antioxidants in dermatology', *Indian Dermatology Online Journal*, 5(2), pp. 210–214. doi: 10.4103/2229-5178.131127.
- Pandel, R. *et al.* (2013) 'Skin photoaging and the role of antioxidants in its prevention', *ISRN dermatology*, 2013, p. 930164. doi: 10.1155/2013/930164.
- Pando, D. *et al.* (2013) 'Nanodesign of olein vesicles for the topical delivery of the antioxidant resveratrol', *The Journal of Pharmacy and Pharmacology*, 65(8), pp. 1158–1167. doi: 10.1111/jphp.12093.

- Patel, A. and Velikov, K. (2013) 'Non-covalent interactions of green tea polyphenol: source of novel colloidal structures', *Nova Biomedical*, pp. 151–166.
- Patil, P. B., Datir, S. K. and Saudagar, R. B. (2019) 'A Review on Topical Gels as Drug Delivery System', *Journal of Drug Delivery and Therapeutics*, 9(3-s), pp. 989–994. doi: 10.22270/jddt.v9i3-s.2930.
- Phongpaichit, S. *et al.* (2007) 'Biological activities of extracts from endophytic fungi isolated from *Garcinia* plants', *FEMS immunology and medical microbiology*, 51(3), pp. 517–525. doi: 10.1111/j.1574-695X.2007.00331.x.
- Pires, F. *et al.* (2019) 'Polycaprolactone/Gelatin Nanofiber Membranes Containing EGCG-Loaded Liposomes and Their Potential Use for Skin Regeneration', *ACS Applied Bio Materials*, 2(11), pp. 4790–4800. doi: 10.1021/acsabm.9b00524.
- Pounikar, Y. *et al.* (2012) 'FORMULATION AND CHARACTERIZATION OF ALOE VERA COSMETIC HERBAL HYDROGEL', *International Journal of Pharmacy and Pharmaceutical Science*, 4, p. 2.
- Pauli, E. D. *et al.* (2014) 'Mixture design analysis of solvent extractor effects on epicatechin, epigallocatechin gallate, epigallocatechin and antioxidant activities of the *Camellia sinensis* L. leaves', *Analytical Chemistry Research*, 2, pp. 23–29. doi: 10.1016/j.ancr.2014.08.001.
- Prasanth, M. I. *et al.* (2019) 'A Review of the Role of Green Tea (*Camellia sinensis*) in Antiphotaging, Stress Resistance, Neuroprotection, and Autophagy', *Nutrients*, 11(2). doi: 10.3390/nu11020474.
- Puri, A., Nguyen, H. X. and Banga, A. K. (2016) 'Microneedle-mediated intradermal delivery of epigallocatechin-3-gallate', *International Journal of Cosmetic Science*, 38(5), pp. 512–523. doi: 10.1111/ics.12320.
- Radhakrishnan, R. *et al.* (2016) 'Encapsulation of biophenolic phytochemical EGCG within lipid nanoparticles enhances its stability and cytotoxicity against cancer', *Chemistry and Physics of Lipids*, 198, pp. 51–60. doi: 10.1016/j.chemphyslip.2016.05.006.
- Ramadon, D. (2017) 'Novel Transdermal Ethosomal Gel Containing Green Tea (*Camellia sinensis* L. Kuntze) Leaves Extract: Formulation and In vitro Penetration Study', *Journal of Young Pharmacists*, 9(3), pp. 336–340. doi: 10.5530/jyp.2017.9.67.
- Ramadon, D., Harme, M. and A, E. (2018) 'Formulation of transfersomal green tea (*Camellia sinensis* L. Kuntze) leaves extract cream and in vitro penetration study using franz diffusion cell', *Journal of Young Pharmacists*, 10(2), pp. s63–s68. doi: 10.5530/jyp.2018.2s.12.

- Rani, R. *et al.* (2014) 'Phytochemical, Pharmacological, and Beneficial Effects of Green Tea', *International Journal of Pharmacognosy and Phytochemical Research*, 6(3), pp. 420–426.
- Reuter, J. *et al.* (2010) 'Which plant for which skin disease? Part 1: Atopic dermatitis, psoriasis, acne, condyloma and herpes simplex', *Journal der Deutschen Dermatologischen Gesellschaft = Journal of the German Society of Dermatology: JDDG*, 8(10), pp. 788–796. doi: 10.1111/j.1610-0387.2010.07496.x.
- Reygaert, W. C. (2014) 'The antimicrobial possibilities of green tea', *Frontiers in Microbiology*, 5. doi: 10.3389/fmicb.2014.00434.
- Rivera-Gonzalez, G., Shook, B. and Horsley, V. (2014) 'Adipocytes in Skin Health and Disease', *Cold Spring Harbor Perspectives in Medicine*, 4(3), pp. a015271–a015271. doi: 10.1101/cshperspect.a015271.
- Roowi, S. *et al.* (2010) 'Green tea flavan-3-ols: colonic degradation and urinary excretion of catabolites by humans', *Journal of Agricultural and Food Chemistry*, 58(2), pp. 1296–1304. doi: 10.1021/jf9032975.
- Rutter, K. *et al.* (2003) 'Green Tea Extract Suppresses the Age-Related Increase in Collagen Crosslinking and Fluorescent Products in C57BL/6 Mice', *International journal for vitamin and nutrition research. Internationale Zeitschrift fur Vitamin- und Ernährungsforschung. Journal international de vitaminologie et de nutrition*, 73(6), pp. 453–460.
- Scalia, S., Marchetti, N. and Bianchi, A. (2013) 'Comparative evaluation of different co-antioxidants on the photochemical- and functional-stability of epigallocatechin-3-gallate in topical creams exposed to simulated sunlight', *Molecules (Basel, Switzerland)*, 18(1), pp. 574–587. doi: 10.3390/molecules18010574.
- Scalia, S., Trotta, V. and Bianchi, A. (2014) 'In vivo human skin penetration of (-)-epigallocatechin-3-gallate from topical formulations', *Acta Pharmaceutica (Zagreb, Croatia)*, 64(2), pp. 257–265. doi: 10.2478/acph-2014-0017.
- Sariyerli, G. S., Sakarya, O. and Akcadag, U. Y. (2018) 'Comparison tests for the determination of the viscosity values of reference liquids by capillary viscometers and stabinger viscometer SVM 3001', *International Journal of Metrology and Quality Engineering*, 9, p.7. doi: 10.1051/ijmqe/2018004.
- Saudagar, R. and Kute, S. B. (2013) 'Emulsified gel A Novel approach for delivery of hydrophobic drugs : An overview', *Journal of Advanced Pharmacy Education and Research*, 3(4), pp. 368–376.
- Scalia, S., Trotta, V. and Bianchi, A. (2014) 'In vivo human skin penetration of (-)-epigallocatechin-3-gallate from topical formulations', *Acta*

Pharmaceutica (Zagreb, Croatia), 64(2), pp. 257–265. doi: 10.2478/acph-2014-0017.

Sharma, A., Saini, S. and Rana, A. (2013) ‘Transdermal Drug Delivery System: A Review’, *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 4(1), pp. 286–292.

Sharma, N., Vashist, H. and Kumar, A. (2018) ‘Reputation of Herbal Drugs in Treatments of Liver Cancer’, *Innovat International Journal of Medical and Pharmaceutical Sciences*, 3(3), pp. 9–16.

Singh, B. N., Shankar, S. and Srivastava, R. K. (2011) ‘Green tea catechin, epigallocatechin-3-gallate (EGCG): mechanisms, perspectives and clinical applications’, *Biochemical pharmacology*, 82(12), pp. 1807–1821. doi: 10.1016/j.bcp.2011.07.093.

Singh, V. and Verma, D. K. (2014) ‘Processing Technology and Health Benefits of Green Tea’, *Popular Kheti*, 2(1), pp. 23–30.

Slamet, S. and U, W. (2019) ‘Optimasi Formulasi Sediaan Handbody Lotion Ekstrak Daun Teh Hijau (*Camellia sinensis* Linn)’, *Pena Jurnal Ilmu Pengetahuan dan Teknologi*, 33(1), pp. 53–57. doi: 10.31941/jurnalpena.v33i1.844.

Song, X. *et al.* (2013) ‘Effect of high-pressure homogenization on particle size and film properties of soy protein isolate’, *Industrial Crops and Products*, 43, pp. 538–544. doi: 10.1016/j.indcrop.2012.08.005.

Srivastava, A. K. *et al.* (2013) ‘Synthesis of PLGA nanoparticles of tea polyphenols and their strong in vivo protective effect against chemically induced DNA damage’, *International Journal of Nanomedicine*, 8, pp. 1451–1462. doi: 10.2147/IJN.S26364.

Sultana, S. S. *et al.* (2014) ‘Emulgel - A Novel Surrogate Approach for Transdermal Drug Delivery System’, *Indo American Journal of Pharmaceutical Research*, 4(2231), pp. 5250–5265.

Tanwar, H. and Sachdeva, R. (2016) ‘Transdermal Drug Delivery System: A review’, *International Journal of Pharmaceutical Sciences and Research*, 7(6), pp. 2274–2290.

Teixé-Roig, J. *et al.* (2018) ‘The Effect of Sodium Carboxymethylcellulose on the Stability and Bioaccessibility of Anthocyanin Water-in-Oil-in-Water Emulsions’, *Food and Bioprocess Technology*, 11(12), pp. 2229–2241. doi: 10.1007/s11947-018-2181-7.

Vishnoi, H., Bodla, R. B. and Kant, R. (2018) ‘Green Tea (*Camellia sinensis*) and Its Antioxidant Property: A Review’, *International Journal of Pharmaceutical Sciences and Research*, 9(5), pp. 1723–1736.

- Wang, S. Q., Balagula, Y. and Osterwalder, U. (2010) 'Photoprotection: a review of the current and future technologies', *Dermatologic Therapy*, 23(1), pp. 31–47. doi: 10.1111/j.1529-8019.2009.01289.x.
- Wisuitiprot, W. *et al.* (2011) 'In vitro human skin permeation and cutaneous metabolism of catechins from green tea extract and green tea extract-loaded chitosan microparticles', *International Journal of Cosmetic Science*, 33(6), pp. 572–579. doi: 10.1111/j.1468-2494.2011.00673.x.
- Yallapu, M. M. *et al.* (2010) 'Fabrication of curcumin encapsulated PLGA nanoparticles for improved therapeutic effects in metastatic cancer cells', *Journal of Colloid and Interface Science*, 351(1), pp. 19–29. doi: 10.1016/j.jcis.2010.05.022.
- Yapar, E. A., Ynal, O. and Erdal, M. S. (2013) 'Design and in vivo evaluation of emulgel formulations including green tea extract and rose oil', *Acta Pharmaceutica (Zagreb, Croatia)*, 63(4), pp. 531–544. doi: 10.2478/acph-2013-0037.
- Zhang, W. *et al.* (2017) 'Sucrose esters improve the colloidal stability of nanoethosomal suspensions of (-)-epigallocatechin gallate for enhancing the effectiveness against UVB-induced skin damage', *Journal of Biomedical Materials Research. Part B, Applied Biomaterials*, 105(8), pp. 2416–2425. doi: 10.1002/jbm.b.33785.
- Zink, A. and Traidl-Hoffmann, C. (2015) 'Green tea in dermatology--myths and facts', *Journal der Deutschen Dermatologischen Gesellschaft = Journal of the German Society of Dermatology: JDDG*, 13(8), pp. 768–775. doi: 10.1111/ddg.12737.