

DAFTAR PUSTAKA

- Austin, M. P. (2002). Spatial prediction of species distribution: an interface between ecological theory and statistical modelling. *Ecological Modelling*, 101–11.
- Balmford, A. A. (2002). Economic Reasons for Conserving Wild Nature. *Science*, 950-953.
- Brook, S. P. (2011). *Extinction of the Javan Rhinoceros (Rhinoceros sondaicus) from Vietnam*. WWF Report Vietnam.
- Colin P. Groves and David M. Leslie, J. (2011). *Rhinoceros sondaicus* (Perissodactyla: Rhinocerotidae). *American Society of Mammalogists*, 190-208.
- Elisabeth S. Bakker, J. L.-C. (2016). Effect of megafauna extinctions on woody plants. *National Academy of Sciences*, (hal. 847-855).
- Elith, J., & Leathwick, J. (2009). Conservation prioritisation using species distribution. Dalam A. W. In Moilanen, *Spatial Conservation Prioritization: Quantitative Methods and Computational Tools* (hal. 70–93). Oxford: Oxford University Press.
- Elith, J., Graham, H. C., & Anderson, R. (2006). Novel methods improve prediction of species distributions from occurrence data. *Ecography*, 129–151.
- Franklin, J. (2009). *Mapping species distributions: spatial inference and prediction*. New York: Cambridge University Press.
- Grubb, P. (2005). *Mammal Species of the World. A Taxonomic and Geographic Reference*. Baltimore: Johns Hopkins University Press.
- Hall, L. K. (1997). The habitat concept and a plea for standard terminology. *Wildlife Society Bulletin*, 173-182.
- Hanley, J., & McNeil, B. (1982). The Meaning and Use of the Area Under a Receiver Operating Characteristic (ROC) Curve. *Radiology*, 143(1), 29–36.
- Haryono, M. M. (2016). *Population and Habitat Viability Assessment for the Javan Rhino*. Apple Valley, MN: IUCN/SSC Conservation Breeding Specialist Group.
- Ian Heywood, S. C. (2006). *An introduction to geographical information systems*. New York: Pentrice Hall.

- Isbell, F., Gonzalez, A., Loreau, M., Cowles, J., Díaz, S., Hector, A., & Mace, G. M. (2017). Linking the influence and dependence of people on biodiversity across scales. *Nature*, 65-72.
- IUCN. (2008). *Rhinoceros sondaicus*. *The IUCN Red List of Threatened Species*.
- Mittermeier RA, G. P. (2005). *Hotspots Revisited: Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions*. New York: Conservation International.
- Moore. (1988). Topographic effects on the distribution of surface soil water and the location of ephemeral gullies. *Transactions of the American Society of Agricultural Engineers*, 1098–1107.
- Phillips, S. J., Anderson, & Schapire. (2006). *Maximum Entropy* modeling of species geographic distributions. *Ecological Modelling*, 231–259.
- Rahmat, M. (2013). Javan Rhino (*Rhinoceros sondaicus* Desmarest 1822) Utilization Distribution and Habitat Selection in Ujung Kulon National Park. *JMHT*.
- Ramono, W. I. (2009). *Report on a second habitat assessment for the Javan rhinoceros (*Rhinoceros sondaicus sondaicus*) within the island of Java*. Florida: International Rhino Foundation.
- Schenkel R., S. L. (1969). The Javan rhinoceros (Rh. sondaicus Desm.) in Ujung Kulon Nature Reserve. Its ecology and behaviour. Field study 1967 and 1968. *Acta Tropica*, 97–135.
- Vitousek, P. M., Mooney, H. A., Lubchenco, J., & Melillo, J. M. (1997). Human Domination of Earth's Ecosystems. *Science*, 494-499.
- Wood, E. M., Pidgeon, A. M., Radeloff, V. C., & Keuler, N. S. (2012). Image texture as a remotely sensed measure of vegetation structure. *Remote Sensing of Environment*, 516-526.