

- [1] R. Magalhães Siqueira, K. Feio Soares Richard, J. Ferreira do Nascimento, A. P. Santana Musse, A. E. Belo Torres, D. Cristina Silva de Azevedo, and M. Bastos-Neto, “Simulation of CO₂/CH₄ high pressure separation on microporous activated carbon,” *Chem. Eng. Commun.*, vol. 206, no. 11, pp. 1425–1436, 2019, doi: 10.1080/00986445.2018.1547713.
- [2] T. Remy, E. Gobechiya, D. Danaci, S. A. Peter, P. Xiao, L. Van Tendeloo, S. Couck, J. Shang, C. E. A. Kirschhock, R. K. Singh, J. A. Martens, G. V. Baron, P. A. Webley, and J. F. M. Denayer, “Biogas upgrading through kinetic separation of carbon dioxide and methane over Rb- and Cs-ZK-5 zeolites,” *RSC Adv.*, vol. 4, no. 107, pp. 62511–62524, 2014, doi: 10.1039/c4ra12460j.
- [3] S. Khalili, B. Khoshandam, and M. Jahanshahi, “A comparative study of CO₂ and CH₄ adsorption using activated carbon prepared from pine cone by phosphoric acid activation,” *Korean J. Chem. Eng.*, vol. 33, no. 10, pp. 2943–2952, 2016, doi: 10.1007/s11814-016-0138-y.
- [4] J. A. Delgado, M. A. Uguina, J. L. Sotelo, B. Ruíz, and M. Rosário, “Carbon Dioxide/Methane Separation by Adsorption on Sepiolite,” *J. Nat. Gas Chem.*, vol. 16, no. 3, pp. 235–243, 2007, doi: 10.1016/S1003-9953(07)60054-1.
- [5] G. K. Parshetti, S. Chowdhury, and R. Balasubramanian, “Biomass derived low-cost microporous adsorbents for efficient CO₂ capture,” *Fuel*, vol. 148, pp. 246–254, 2015, doi: 10.1016/j.fuel.2015.01.032.
- [6] B. Yuan, X. Wu, Y. Chen, J. Huang, H. Luo, and S. Deng, “Adsorption of CO₂, CH₄, and N₂ on Ordered mesoporous carbon: Approach for greenhouse gases capture and biogas upgrading,” *Environ. Sci. Technol.*, vol. 47, no. 10, pp. 5474–5480, 2013, doi: 10.1021/es4000643.

- [7] N. Alvarez-Gutiérrez, S. García, M. V. Gil, F. Rubiera, and C. Pevida, “Dynamic Performance of Biomass-Based Carbons for CO₂ /CH₄ Separation. Approximation to a Pressure Swing Adsorption Process for Biogas Upgrading,” *Energy and Fuels*, vol. 30, no. 6, pp. 5005–5015, 2016, doi: 10.1021/acs.energyfuels.6b00664.
- [8] Y. Jiang, J. Ling, P. Xiao, Y. He, Q. Zhao, Z. Chu, Y. Liu, Z. Li, and P. A. Webley, “Simultaneous biogas purification and CO₂ capture by vacuum swing adsorption using zeolite NaUSY,” *Chem. Eng. J.*, vol. 334, pp. 2593–2602, 2018, doi: 10.1016/j.cej.2017.11.090.
- [9] S. Sircar, “Pressure swing adsorption page4.pdf,” *Ind. Eng. Chem.*, vol. 41, pp. 1389–1392, 2002.
- [10] M. G. Plaza, S. García, F. Rubiera, J. J. Pis, and C. Pevida, “Post-combustion CO₂ capture with a commercial activated carbon: Comparison of different regeneration strategies,” *Chem. Eng. J.*, vol. 163, no. 1–2, pp. 41–47, 2010, doi: 10.1016/j.cej.2010.07.030.
- [11] C. O. Edmund, M. S. Christopher, and D. K. Pascal, “Characterization of palm kernel shell for materials reinforcement and water treatment,” *J. Chem. Eng. Mater. Sci.*, vol. 5, no. 1, pp. 1–6, 2014, doi: 10.5897/jcem2014.0172.
- [12] I. Prasetyo, Rochmadi, E. Wahyono, and T. Ariyanto, “Controlling synthesis of polymer-derived carbon molecular sieve and its performance for CO₂/CH₄ separation,” *Eng. J.*, vol. 21, no. 4, pp. 83–94, 2017, doi: 10.4186/ej.2017.21.4.83.
- [13] A. Jayaraman, A. S. Chiao, J. Padin, R. T. Yang, and C. L. Munson, “Kinetic separation of methane/carbon dioxide by molecular sieve carbons,” *Sep. Sci. Technol.*, vol. 37, no. 11, pp. 2505–2528, 2002, doi: 10.1081/SS-120004450.
- [14] X. Song, L. Wang, X. Ma, and Y. Zeng, “Adsorption equilibrium and thermodynamics of CO₂ and CH₄ on carbon molecular sieves,” *Appl. Surf. Sci.*, vol.

- [15] I. Prasetyo and D. D. Do, “Adsorption rate of methane and carbon dioxide on activated carbon by the semi-batch constant molar flow rate method,” *Chem. Eng. Sci.*, vol. 53, no. 19, pp. 3459–3467, 1998, doi: 10.1016/S0009-2509(98)00146-8.
- [16] Y. Shen, W. Shi, D. Zhang, P. Na, and B. Fu, “The removal and capture of CO₂ from biogas by vacuum pressure swing process using silica gel,” *J. CO₂ Util.*, vol. 27, no. July, pp. 259–271, 2018, doi: 10.1016/j.jcou.2018.08.001.
- [17] G. Klein, “Principles of adsorption and adsorption processes,” *Reactive Polymers, Ion Exchangers, Sorbents*, vol. 4, no. 1. p. 62, 1985, doi: 10.1016/0167-6989(85)90037-6.
- [18] L. Song and L. V. C. Rees, “Frequency response measurements of diffusion in microporous materials,” *Mol. Sieves - Sci. Technol.*, vol. 7, no. September 2007, pp. 235–276, 2008, doi: 10.1007/3829_2007_013.
- [19] D. B. Shah and D. M. Ruthven, “Measurement of zeolitic diffusivities and equilibrium isotherms by chromatography,” *AIChE J.*, vol. 23, no. 6, pp. 804–809, 1977, doi: 10.1002/aic.690230605.
- [20] M. Goddard and D. M. Ruthven, “Sorption and diffusion of C₈ aromatic hydrocarbons in faujasite-type zeolites III. Self-diffusivities by tracer exchange,” *Zeolites*, vol. 6, no. 6, pp. 445–448, 1986, doi: 10.1016/0144-2449(86)90028-X.
- [21] D. D. Do, *Adsorption Analysis: Equilibria and Kinetics.*, vol. 2. 1998.
- [22] R. P. Ribeiro, T. P. Sauer, F. V. Lopes, R. F. Moreira, C. A. Grande, and A. E. Rodrigues, “Adsorption of CO₂, CH₄, and N₂ in activated carbon honeycomb monolith,” *J. Chem. Eng. Data*, vol. 53, no. 10, pp. 2311–2317, 2008, doi: 10.1021/je800161m.
- [23] E. Ryckebosch, M. Drouillon, and H. Vervaeren, “Techniques for transformation of

doi: 10.1016/j.biombioe.2011.02.033.

- [24] S. O. Masebinu, A. Aboyade, and E. Muzenda, “Enrichment of Biogas for Use as Vehicular Fuel : A Review of the Upgrading Techniques,” *Int. J. Res. Chem. , Metall. Civ. Eng.*, vol. 1, no. 1, pp. 89–98, 2014.
- [25] J. Niesner, D. Jecha, and P. Stehlík, “Biogas upgrading technologies: State of art review in european region,” *Chem. Eng. Trans.*, vol. 35, pp. 517–522, 2013, doi: 10.3303/CET1335086.
- [26] R. Kadam and N. L. Panwar, “Recent advancement in biogas enrichment and its applications,” *Renew. Sustain. Energy Rev.*, vol. 73, no. January, pp. 892–903, 2017, doi: 10.1016/j.rser.2017.01.167.
- [27] V. Vrbová and K. Ciahotný, “Upgrading Biogas to Biomethane Using Membrane Separation,” *Energy and Fuels*, vol. 31, no. 9, pp. 9393–9401, 2017, doi: 10.1021/acs.energyfuels.7b00120.
- [28] G. Xu, Y. Lv, J. Sun, H. Shao, and L. Wei, “Recent Advances in Biochar Applications in Agricultural Soils: Benefits and Environmental Implications,” *CLEAN - Soil, Air, Water*, vol. 40, no. 10, pp. 1093–1098, Oct. 2012, doi: 10.1002/clen.201100738.
- [29] M. Younas, M. Sohail, L. L. Kong, M. J. K. Bashir, and S. Sethupathi, “Feasibility of CO₂adsorption by solid adsorbents: a review on low-temperature systems,” *Int. J. Environ. Sci. Technol.*, vol. 13, no. 7, pp. 1839–1860, 2016, doi: 10.1007/s13762-016-1008-1.
- [30] Suhas, P. J. M. Carrott, and M. M. L. Ribeiro Carrott, “Lignin – from natural adsorbent to activated carbon: A review,” *Bioresour. Technol.*, vol. 98, no. 12, pp. 2301–2312, Sep. 2007, doi: 10.1016/j.biortech.2006.08.008.
- [31] F. Rodríguez-Reinoso, M. Molina-Sabio, and M. T. González, “The use of steam and

CO₂ as activating agents in the preparation of activated carbons,” *Carbon N. Y.*, vol.

33, no. 1, pp. 15–23, 1995, doi: 10.1016/0008-6223(94)00100-E.

- [32] G. Singh, K. S. Lakhi, S. Sil, S. V. Bhosale, I. Y. Kim, K. Albahily, and A. Vinu, “Biomass derived porous carbon for CO₂ capture,” *Carbon N. Y.*, vol. 148, pp. 164–186, 2019, doi: 10.1016/j.carbon.2019.03.050.
- [33] H. Teng and L. Y. Hsu, “High-Porosity Carbons Prepared from Bituminous Coal with Potassium Hydroxide Activation - Industrial & Engineering Chemistry Research (ACS Publications),” *Ind. Eng. Chem. Res.*, pp. 2947–2953, 1999, [Online]. Available:
<http://pubs.acs.org/doi/abs/10.1021/ie990101%2B%5Cnpapers2://publication/uuid/B9B327B5-A382-40A4-8EAE-CEA389D7C951>.
- [34] S. M. Manocha, “Porous Carbon,” *Porous Carbons*, vol. 28, no. April, pp. 335–348, 2003.
- [35] A. M. Silvestre-Albero, A. Wahby, J. Silvestre-Albero, F. Rodríguez-Reinoso, and W. Betz, “Carbon molecular sieves prepared from polymeric precursors: Porous structure and hydrogen adsorption properties,” *Ind. Eng. Chem. Res.*, vol. 48, no. 15, pp. 7125–7131, 2009, doi: 10.1021/ie900091n.
- [36] D. Cazorla-Amorós, J. Alcañiz-Monge, M. A. De La Casa-Lillo, and A. Linares-Solano, “CO₂ as an adsorptive to characterize carbon molecular sieves and activated carbons,” *Langmuir*, vol. 14, no. 16, pp. 4589–4596, 1998, doi: 10.1021/la980198p.
- [37] S. Cho, D. Lee, and Y. S. Lee, “Separation of biomass using carbon molecular sieves treated with hydrogen peroxide,” *J. Ind. Eng. Chem.*, vol. 21, pp. 278–282, 2015, doi: 10.1016/j.jiec.2013.12.113.
- [38] M. B. Kim, Y. S. Bae, D. K. Choi, and C. H. Lee, “Kinetic separation of landfill gas by a two-bed pressure swing adsorption process packed with carbon molecular sieve:

Nonisothermal operation,” *Ind. Eng. Chem. Res.*, vol. 45, no. 14, pp. 5050–5058, 2006, doi: 10.1021/ie0511074.

- [39] M. Miltner, A. Makaruk, and M. Harasek, “Review on available biogas upgrading technologies and innovations towards advanced solutions,” *J. Clean. Prod.*, vol. 161, pp. 1329–1337, 2017, doi: 10.1016/j.jclepro.2017.06.045.
- [40] M. S. Shafeeyan, W. M. A. W. Daud, A. Shamiri, and N. Aghamohammadi, “Modeling of Carbon Dioxide Adsorption onto Ammonia-Modified Activated Carbon: Kinetic Analysis and Breakthrough Behavior,” *Energy and Fuels*, vol. 29, no. 10, pp. 6565–6577, 2015, doi: 10.1021/acs.energyfuels.5b00653.
- [41] S. Sircar, J. R. Hufton, and A. Products, “Why Does the Linear Driving Force Model for Adsorption Kinetics Work ?,” pp. 137–147, 2000.
- [42] R. S. Todd and P. A. Webley, “Pressure Drop in a Packed Bed under Nonadsorbing and Adsorbing Conditions,” *Ind. Eng. Chem.*, vol. 44, pp. 7234–7241, 2005.
- [43] S. M. Manocha, “Porous carbons,” *Sadhana*, vol. 28, no. 1–2, pp. 335–348, Feb. 2003, doi: 10.1007/BF02717142.
- [44] I. Prasetyo, N. I. F. Mukti, R. B. Cahyono, A. Prasetya, and T. Ariyanto, “Nanoporous Carbon Prepared from Palm Kernel Shell for CO₂/CH₄ Separation,” *Waste and Biomass Valorization*, no. 0123456789, 2020, doi: 10.1007/s12649-020-01006-4.
- [45] A. R. Hidayu and N. Muda, “Preparation and Characterization of Impregnated Activated Carbon from Palm Kernel Shell and Coconut Shell for CO₂ Capture,” *Procedia Eng.*, vol. 148, pp. 106–113, 2016, doi: 10.1016/j.proeng.2016.06.463.
- [46] N. A. Rashidi and S. Yusup, “Production of palm kernel shell-based activated carbon by direct physical activation for carbon dioxide adsorption,” *Environ. Sci. Pollut. Res.*, vol. 26, no. 33, pp. 33732–33746, Nov. 2019, doi: 10.1007/s11356-018-1903-8.
- [47] V. K. Singh and E. Anil Kumar, “Measurement and analysis of adsorption isotherms

10.1016/j.applthermaleng.2015.10.052.

- [48] G. Fraux, A. Boutin, A. H. Fuchs, and F. X. Coudert, “On the use of the IAST method for gas separation studies in porous materials with gate-opening behavior,” *Adsorption*, vol. 24, no. 3, pp. 233–241, 2018, doi: 10.1007/s10450-018-9942-5.
- [49] K. S. Walton and D. S. Sholl, “Predicting multicomponent adsorption: 50 years of the ideal adsorbed solution theory,” *AIChE J.*, vol. 61, no. 9, pp. 2757–2762, Sep. 2015, doi: 10.1002/aic.14878.
- [50] J. Chen, L. S. Loo, and K. Wang, “An Ideal Adsorbed Solution Theory (IAST) Study of Adsorption Equilibria of Binary Mixtures of Methane and Ethane on a Templated Carbon,” *J. Chem. Eng. Data*, vol. 56, no. 4, pp. 1209–1212, Apr. 2011, doi: 10.1021/je101099c.
- [51] A. Silvestre-Albero, S. Rico-Francés, F. Rodríguez-Reinoso, A. M. Kern, M. Klumpp, B. J. M. Etzold, and J. Silvestre-Albero, “High selectivity of TiC-CDC for CO₂/N₂ separation,” *Carbon N. Y.*, vol. 59, pp. 221–228, Aug. 2013, doi: 10.1016/j.carbon.2013.03.012.
- [52] H. Wei, S. Deng, B. Hu, Z. Chen, B. Wang, J. Huang, and G. Yu, “Granular Bamboo-Derived Activated Carbon for High CO₂ Adsorption: The Dominant Role of Narrow Micropores,” *ChemSusChem*, vol. 5, no. 12, pp. 2354–2360, Dec. 2012, doi: 10.1002/cssc.201200570.
- [53] N. Álvarez-Gutiérrez, S. García, M. V. Gil, F. Rubiera, and C. Pevida, “Dynamic Performance of Biomass-Based Carbons for CO₂/CH₄ Separation. Approximation to a Pressure Swing Adsorption Process for Biogas Upgrading,” *Energy & Fuels*, vol. 30, no. 6, pp. 5005–5015, Jun. 2016, doi: 10.1021/acs.energyfuels.6b00664.
- [54] N. Wakao and T. Funazkri, “Effect of fluid dispersion coefficients on particle-to-fluid

Chemical Engineering Science, vol. 33, no. 10. pp. 1375–1384, 1978, doi: 10.1016/0009-2509(78)85120-3.

- [55] A. Malek and S. Farooq, “Kinetics of hydrocarbon adsorption on activated carbon and silica gel,” *AIChE J.*, vol. 43, no. 3, pp. 761–776, 1997, doi: 10.1002/aic.690430321.
- [56] X. Hu and D. D. Do, “Multicomponent adsorption kinetics of hydrocarbons onto activated carbon: Contribution of micropore resistance,” *Chem. Eng. Sci.*, vol. 48, no. 7, pp. 1317–1323, Apr. 1993, doi: 10.1016/0009-2509(93)81012-K.
- [57] N. Al-Janabi, R. Vakili, P. Kalumpasut, P. Gorgojo, F. R. Siperstein, X. Fan, and P. McCloskey, “Velocity variation effect in fixed bed columns: A case study of CO₂ capture using porous solid adsorbents,” *AIChE J.*, vol. 64, no. 6, pp. 2189–2197, Jun. 2018, doi: 10.1002/aic.16135.
- [58] S. H. Ha, K. H. Kim, S. A. Younis, and X. Dou, “The interactive roles of space velocity and particle size in a microporous carbon bed system in controlling adsorptive removal of gaseous benzene under ambient conditions,” *Chem. Eng. J.*, vol. 401, p. 126010, 2020, doi: 10.1016/j.cej.2020.126010.
- [59] R. Sabouni, H. Kazemian, and S. Rohani, “Mathematical modeling and experimental breakthrough curves of carbon dioxide adsorption on metal organic framework CPM-5,” *Environ. Sci. Technol.*, vol. 47, no. 16, pp. 9372–9380, 2013, doi: 10.1021/es401276r.
- [60] A. Möller, R. Eschrich, C. Reichenbach, J. Guderian, M. Lange, and J. Möllmer, “Dynamic and equilibrium-based investigations of CO₂-removal from CH₄-rich gas mixtures on microporous adsorbents,” *Adsorption*, vol. 23, no. 2–3, pp. 197–209, 2017, doi: 10.1007/s10450-016-9821-x.
- [61] Y. Sun, G. Yang, J. Zhang, Y. Wang, and M. Yao, “Activated Carbon Preparation

- [62] H. Azarabadi and K. S. Lackner, “A sorbent-focused techno-economic analysis of direct air capture,” *Appl. Energy*, vol. 250, pp. 959–975, Sep. 2019, doi: 10.1016/j.apenergy.2019.04.012.
- [63] J. Zhang, P. A. Webley, and P. Xiao, “Effect of process parameters on power requirements of vacuum swing adsorption technology for CO₂ capture from flue gas,” *Energy Convers. Manag.*, vol. 49, no. 2, pp. 346–356, 2008, doi: 10.1016/j.enconman.2007.06.007.
- [64] M. T. Ho, G. W. Allinson, and D. E. Wiley, “Reducing the cost of CO₂ capture from flue gases using pressure swing adsorption,” *Ind. Eng. Chem. Res.*, vol. 47, no. 14, pp. 4883–4890, 2008, doi: 10.1021/ie070831e.
- [65] R. B. Cahyono, A. N. Rozhan, N. Yasuda, T. Nomura, H. Purwanto, and T. Akiyama, “Carbon deposition using various solid fuels for ironmaking applications,” *Energy and Fuels*, vol. 27, no. 5, pp. 2687–2692, 2013, doi: 10.1021/ef400322w.