



DAFTAR PUSTAKA

A. Daftar Pustaka

- Albrektsen, G., Heuch, I., Løchen, M., Thelle, D., Wilsgaard, T., Njølstad, I., & Bønaa, K. (2016). Lifelong Gender Gap in Risk of Incident Myocardial Infarction. *JAMA Internal Medicine*, 176(11), p.1673. <https://doi.org/10.1001/jamainternmed.2016.54.151>
- Altmann DB, Racz M, Battleman DS, Bergman G, Spokojny A, Hannan EL, Sanborn TA. (1996). Reduction In Angioplasty Complications After The Introduction Of Coronary Stents: Results From A Consecutive Series of 2242 Patients. *Am Heart J*, 132(3):503-7. [https://doi.org/10.1016/S0002-8703\(96\)90231-8](https://doi.org/10.1016/S0002-8703(96)90231-8)
- Akbar H, Foth C, Kahloon RA, et al. Acute Myocardial Infarction ST Elevation (STEMI) [Updated 2020 Jun 24]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK532281/>
- Assessment of the safety and Efficacy of a New Treatment Strategy with Percutaneous Coronary Intervention. (2006). Primary versus tenecteplase-facilitated percutaneous coronary intervention in patients with ST-segment elevation acute myocardial infarction (ASSENT-4 PCI): randomised trial. *The Lancet*, 367(9510), 569–578. [https://doi.org/10.1016/s0140-6736\(06\)68147-6](https://doi.org/10.1016/s0140-6736(06)68147-6)
- Bär, F., Vainer, J., Stevenhagen, J., Neven, K., Aalbregt, R., Oude Ophuis, T., Wellens, H. (2000). Ten-Year Experience With Early Angioplasty In 759 Patients With Acute Myocardial Infarction. *Journal of the American College of Cardiology*, 36(1), 51–58. [https://doi.org/10.1016/s0735-1097\(00\)00718-x](https://doi.org/10.1016/s0735-1097(00)00718-x)
- Barrett-Connor, E. (1997). Sex Differences in Coronary Heart Disease. *Circulation*, 95(1), pp.252-264. <https://doi.org/10.1161/01.CIR.95.1.252>
- Body, R. (2018). Acute Coronary Syndromes Diagnosis, version 2.0: Tomorrow's Approach to Diagnosing Acute Coronary Syndromes?. *Turkish Journal of Emergency Medicine*, 18(3), pp.94-99. <https://doi.org/10.1016/j.tjem.2018.05.005>
- Boey, E., Gay, G. M. W., Poh, K.-K., & Lee, C.-H. (2016). Visit-to-visit variability in LDL- and HDL-cholesterol is associated with adverse events after ST-segment elevation myocardial infarction: A 5-year follow-up study. *Atherosclerosis*, 244, 86–92. <https://doi.org/10.1016/j.atherosclerosis.2015.10.110>



Bonaca M. P., Bhatt D. L., Fish M. P., & Im K. (2015). Long-Term Use of Ticagrelor in Patients with Prior Myocardial Infarction. *N Engl J Med*, 372(19):1791–1800. <https://doi.org/10.1056/NEJMoa1500857>

Bozbay, M., Uyarel, H., Cicek, G., Oz, A., Keskin, M., Murat, A., Eren, M. (2016). CHA2DS2-VASc Score Predicts In-Hospital and Long-Term Clinical Outcomes in Patients With ST-Segment Elevation Myocardial Infarction Who Were Undergoing Primary Percutaneous Coronary Intervention. *Clinical and Applied Thrombosis/Hemostasis*, 23(2), 132–138. <https://doi.org/10.1177/1076029616646874>

Budiman, B., Sihombing, R., & Pradina, P. (2017). Hubungan Dislipidemia, Hipertensi Dan Diabetes Melitus Dengan Kejadian Infark Miokard Akut. *Jurnal Kesehatan Masyarakat Andalas*, 10(1), p.32. <http://jurnal.fkm.unand.ac.id/index.php/jkma/>

Carro, A., & Juan C. (2010). Myocardial Infarction in the Elderly. *Aging and Disease*, 2(1), pp.116-137. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3295051/pdf/ad-2-2-116.pdf>

Choi, B., Rha, S., Yoon, S., Choi, C., Lee, M., & Kim, S. (2019). Association of Major Adverse Cardiac Events up to 5 Years in Patients With Chest Pain Without Significant Coronary Artery Disease in the Korean Population. *Journal of the American Heart Association*, 8(12). <https://doi.org/10.1161/JAHA.118.01054.11>

Chua, S. K., Shyu, K. G., & Hung, H. F. (2014). Gender and Age Differences in Short- and Long-Term Outcomes Following Primary Percutaneous Coronary Intervention for ST-Elevation Myocardial Infarction. *Acta Cardiologica Sinica*, 30(4), 274–283.

Diagnosis and Classification of Diabetes Mellitus. (2004). *Diabetes Care*, 27(Supplement 1), pp.S5-S10.

De Luca, G., Ernst, N., van't Hof, A. W. J., Ottenvanger, J. P., Hoornje, J. C. A., Gosselink, A. T. M., Suryapranata, H. (2006). Predictors and clinical implications of early reinfarction after primary angioplasty for ST-segment elevation myocardial infarction. *American Heart Journal*, 151(6), 1256–1259. <https://10.1016/j.ahj.2005.06.047>

De Luca, G., Dirksen, M. T., Spaulding, C., Kelbæk, H., Schalij, M., Thuesen, L., Stone, G. W. (2014). Impact of hypertension on clinical outcome in STEMI patients undergoing primary angioplasty with BMS or DES. *International Journal of Cardiology*, 175(1), 50–54,1. <https://doi.org/10.1016/j.ijcard.2014.04.180>

De Luca G, Parodi G, Sciagrà R, et al. (2013). Impact of hypertension on infarct size in ST elevation myocardial infarction patients undergoing primary

angioplasty. *J Hypertens.* 2013;31(12):2433-2437.
<https://doi.org/10.1097/HJH.0b013e328364cbee>

Dönges, K., Schiele, R., Gitt, A., Wienbergen, H., Schneider, S., Zahn, R., Senges, J. (2001). Incidence, determinants, and clinical course of reinfarction in-hospital after index acute myocardial infarction (results from the pooled data of the maximal individual therapy in acute myocardial infarction [mitra], and the myocardial infarction registry [MIR]). *The American Journal of Cardiology*, 87(9), 1039–1044. [https://doi.org/10.1016/s0002-9149\(01\)01458-8](https://doi.org/10.1016/s0002-9149(01)01458-8)

Ellis SG, Tendera M, de Belder MA, van Boven AJ. (2008). Facilitated PCI in patients with ST-elevation myocardial infarction. *N Engl J Med*, 22;358(21), 2205-17. <https://doi.org/10.1056/NEJMoa0706816>

Faslur Rahuman, M. B., Jayawardena, J. B., Francis, G. R., Mahboob, N., Kumara A.H.T., W., Wijesinghe, A., De Silva, P. A. (2017). A Comparison Of Rescue And Primary Percutaneous Coronary Interventions For Acute ST Elevation Myocardial Infarction. *Indian Heart Journal*, 69, S57–S62. <https://doi.org/10.1016/j.ihj.2017.02.019>

Fokkema, M. L., van der Vleuten, P. A., Vlaar, P. J., Svilaas, T., & Zijlstra, F. (2009). Incidence, predictors, and outcome of reinfarction and stent thrombosis within one year after primary percutaneous coronary intervention for ST-elevation myocardial infarction. *Catheterization and Cardiovascular Interventions*, 73(5), 627–634. <https://doi.org/10.1002/ccd.21887>

Frangogiannis, N. (2015). Pathophysiology of Myocardial Infarction. *Comprehensive Physiology*, pp.1841-1875. <https://doi.org/10.1002/cphy.c150006>

Ganassin, F. P., de Carvalho Cantarelli, M. J., Castello, H. J., Gonçalves, R., Ribeiro, E. K. P., de Freitas Guimaraes, J. B., de Almeida, R. S. (2013). In-Hospital Outcomes on Patients Submitted to Primary Percutaneous Coronary Intervention Versus Rescue. *Revista Brasileira de Cardiologia Invasiva (English Edition)*, 21(2), 133–139. [https://doi.org/10.1016/s2214-1235\(15\)30121-6](https://doi.org/10.1016/s2214-1235(15)30121-6)

Goss, F., Brachmann, J., & Levenson, B. (2017). High adherence to therapy and low cardiac mortality and morbidity in patients after acute coronary syndrome systematically managed by office-based cardiologists in Germany: 1-year outcomes of the ProAcor Study. *Vascular Health and Risk Management*, Volume 13, 127–137. <https://doi.org/10.2147/vhrm.s119490>

Harjai, K. J., Stone, G. W., Boura, J., Mattos, L., Chandra, H., Cox, D., Grines, C. (2003). Comparison of outcomes of diabetic and nondiabetic patients undergoing primary angioplasty for acute myocardial infarction. *The*



American Journal of Cardiology, 91(9), 1041–1045.
[https://doi.org/10.1016/s0002-9149\(03\)00145-0](https://doi.org/10.1016/s0002-9149(03)00145-0)

Harjola, V., Parissis, J., Bauersachs, J., Brunner-La Rocca, H., Bueno, H., Čelutkienė, J., & Yilmaz, M. B. (2020). Acute coronary syndromes and acute heart failure: a diagnostic dilemma and high-risk combination. A statement from the Acute Heart Failure Committee of the Heart Failure Association of the European Society of Cardiology. *European Journal of Heart Failure*. <https://doi.org/10.1002/ejhf.1831>

Harrison, T., Kasper, D., Longo, D., Fauci, A., Hauser, S., Jameson, J., & Loscalzo, J. (2013). *Harrison's Cardiovascular Medicine*. 2nd ed. Boston: McGraw-Hill Education, pp.385-425.

Huber, C. A., Meyer, M. R., Steffel, J., Blozik, E., Reich, O., & Rosemann, T. (2018). Post-Myocardial Infarction (MI) Care: Medication Adherence for Secondary Prevention After MI in a Large Real-World Population. *Clinical Therapeutics*. <https://doi.org/10.1016/j.clinthera.2018.11.012>

Hur, S. H., Won, K. B., Kim, I. C., Bae, J. H., Choi, D. J., Ahn, Y. K., Park, J. S., Kim, H. S., Choi, R. K., Choi, D., & DIAMOND investigators. (2016). Comparison of 2-year clinical outcomes between diabetic versus nondiabetic patients with acute myocardial infarction after 1-month stabilization: Analysis of the prospective registry of DIAMOND (DIabetic acute myocardial infarctiON Disease) in Korea: an observational registry study. *Medicine*, 95(25), e3882.
<https://doi.org/10.1097/MD.0000000000003882>

Ibrahim, A., Riddell, T., & Devireddy, C. (2014). Acute Myocardial Infarction. *Crit Care Lin*, 30, p.341-346. <http://dx.doi.org/10.1016/j.ccc.2014.03.010>

Ismail, M. D., Jalalonmuhalil, M., Azhari, Z., Mariapun, J., Lee, Z. V., Zainal Abidin, I., Wan Ahmad, W. A., Zuhdi, A., & NCVD-PCI investigators. (2018). Outcomes of STEMI patients with chronic kidney disease treated with percutaneous coronary intervention: the Malaysian National Cardiovascular Disease Database - Percutaneous Coronary Intervention (NCVD-PCI) registry data from 2007 to 2014. *BMC cardiovascular disorders*, 18(1), 184. <https://doi.org/10.1186/s12872-018-0919-9>

Jan, S., Lee, S., Sawhney, J., Ong, T., Chin, C., Kim, H., Krittayaphong, R., Nhan, V., Itoh, Y., & Huo, Y. (2016). Catastrophic health expenditure on acute coronary events in Asia: a prospective study. *Bull World Health Organ*, pp.193-200. <http://dx.doi.org/10.2471/BLT.15.158303>

Keeley, E. C., Boura, J. A., & Grines, C. L. (2003). Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *The Lancet*, 361(9351), 13–20. [https://doi.org/0.1016/s0140-6736\(03\)12113-7](https://doi.org/0.1016/s0140-6736(03)12113-7)



Kelompok Staf Medik Jantung. (2017). *Panduan Praktek Klinis: Infark Miokard dengan Elevasi Segmen ST*. Yogyakarta: RSUP Dr. Sardjito.

Kelompok Staf Medik Jantung. (2019). *Panduan Praktek Klinis: Infark Miokard dengan Non-Elevasi Segmen ST*. Yogyakarta: RSUP Dr. Sardjito

Kementerian Kesehatan Republik Indonesia. (2018). Riset Kesehatan Dasar. Diunduh dari http://www.kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-riskesdas-2018_1274.pdf

Kernis, S. J., Harjai, K. J., Stone, G. W., Grines, L. L., Boura, J. A., Yerkey, M. W., Grines, C. L. (2003). The incidence, predictors, and outcomes of early reinfarction after primary angioplasty for acute myocardial infarction. *Journal of the American College of Cardiology*, 42(7), 1173–1177. [https://doi.org/10.1016/s0735-1097\(03\)00920-3](https://doi.org/10.1016/s0735-1097(03)00920-3)

Kip, K., Hollabaugh, K., Marroquin, O., & Williams, D. (2008). The Problem With Composite End Points in Cardiovascular Studies. *Journal of the American College of Cardiology*, 51(7), pp.701-707. <https://doi.org/10.1016/j.jacc.2007.10.034>

Kozma, C., Dickson, M., Phillips, A., & Meletiche, D. (2013). Medication Possession Ratio: Implications of Using Fixed and Variable Observation Periods in Assessing Adherence with Disease-Modifying Drugs in Patients with Multiple Sclerosis. *Patient Preference and Adherence*, 7, pp. 509-516. <https://doi.org/10.2147/PPA.S40736>

Lu, L., Liu, M., Sun, R., Zheng, Y., & Zhang, P. (2015). Myocardial Infarction: Symptoms and Treatments. *Cell Biochemistry and Biophysics*, 72(3), pp.865-867. <http://dx.doi.org/10.1007/s12013-015-0553-4>

Makki, N., Brennan, T., & Girotra, S. (2013). Acute Coronary Syndrome. *Journal of Intensive Care Medicine*, 0(0), pp.1-15. <http://doi.org/10.1177/0885066613503294>

Mal, K., Awan, I., & Shaukat, F. (2019). Evaluation of Risk Factors Associated with Reinfarction: A Multicenter Observational Study. *Cureus*, 11(11), pp.1-6. <http://doi.org/10.7759/cureus.6063>

Mauri, L., Kereiakes, D. J., Yeh, R. W., Driscoll-Shempp, P., Cutlip, D. E., Steg, P. G., & Normand, S. L. (2014). Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *The New England journal of medicine*, 371(23), 2155–2166. <https://doi.org/10.1056/NEJMoa1409312>

Modi,K., Soos, M., Mahajan, K. (2019). *Stent Thrombosis*. Dikunjungi pada Fabruari 14, 2020, dari <https://www.ncbi.nlm.nih.gov/books/NBK441908/>



Otten, A. M., Maas, A. H., Ottervanger, J. P., Kloosterman, A., van 't Hof, A. W., Dambrink, J. H., Gosselink, A. T., Hoornje, J. C., Suryapranata, & Zwolle Myocardial Infarction study Group. (2013). Is the difference in outcome between men and women treated by primary percutaneous coronary intervention age dependent? Gender difference in STEMI stratified on age. *European Heart Journal Acute Cardiovascular Care*, 2(4), 334–341. <https://doi.org/10.1177/2048872612475270>

Pan, F., Chernew, E., & Fendrick, M. (2008). Impact of fixed-dose combination drugs on adherence to prescription medications. *Journal of general internal medicine*, 23(5), 611–614. <https://doi.org/10.1007/s11606-008-05414-x>

Park, J. S., Cha, K. S., Shin, D., Lee, D. S., Lee, H. W., Oh, J. H., Choi, J. H., Lee, H. C., Hong, T. J., Lee, S. H., Kim, J. S., & Korea Working Group on Myocardial Infarction Investigators. (2015). Prognostic Significance of Presenting Blood Pressure in Patients With ST-Elevation Myocardial Infarction Undergoing Percutaneous Coronary Intervention. *American journal of hypertension*, 28(6), 797–805. <https://doi.org/10.1093/ajh/hpu230>

Perhimpunan Dokter Spesialis Kardiovaskular Indonesia. (2018). *Pedoman Tata Laksana Sindrom Koroner Akut*. Jakarta: Pengurus Pusat Pedoman Tata Laksana Sindrom Koroner Akut.

Perkumpulan Endokrinologi Indonesia. (2015). *Konsensus Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 Di Indonesia*. Jakarta: Pengurus Besar Perkumpulan Endokrinologi Indonesia.

Perkumpulan Endokrinologi Indonesia. (2015). *Panduan Pengelolaan Dislipidemia*. Jakarta: Pengurus Besar Perkumpulan Endokrinologi Indonesia.

Picariello, C., Lazzeri, C., Attana, P., Chiostri, M., Gensini, G., & Valente, S. (2011). The Impact of Hypertension on Patients with Acute Coronary Syndromes. *International Journal of Hypertension*, 11, pp.1-7. <https://doi.org/10.4061/2011/563657>

Polanska-Skrzypczyk, M., Karcz, M., Kepka, S. (2013). Prognostic value of renal function in STEMI patients treated with primary PCI: ANIN Registry. *British Journal of Cardiology*. <https://doi.org/10.5837/bjc.2013.17>

Que, B., Wang, C., Ai, H., Zhang, X., Wang, M., & Nie, S. (2016). Residual Dyslipidemia Leads to Unfavorable Outcomes in Patients with Acute Coronary Syndrome after Percutaneous Coronary Intervention. *Stem Cells International*, 2016, 1–5. <https://doi.org/10.1155/2016/6175948>

Radovanovic, D., Maurer, L., Bertel, O., Witassek, F., Urban, P., Stauffer, J.-C., ... Erne, P. (2016). Treatment and outcomes of patients with recurrent



myocardial infarction: A prospective observational cohort study. *Journal of Cardiology*, 68(6), 498–503. <https://doi.org/10.1016/j.jcc.2015.11.013>

Reinstadler, S. J., Stiermaier, T., Eitel, C., Saad, M., Metzler, B., de Waha, S., Fuernau, G., Desch, S., Thiele, H., & Eitel, I. (2016). Antecedent hypertension and myocardial injury in patients with reperfused ST-elevation myocardial infarction. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*, 18(1), 80. <https://doi.org/10.1186/s12968-016-0299-1>

Reinstadler SJ, Stiermaier T, Eitel C, et al. (2017). Relationship between diabetes and ischaemic injury among patients with revascularized ST-elevation myocardial infarction. *Diabetes Obes Metab.* 2017;19(12):1706-1713. <https://doi.org/10.1111/dom.13002>

Saleh, M. & Ambrose, J. A. (2018). Understanding myocardial infarction. *F1000Research*, 7, pp1-8. <http://doi.org/10.12688/f1000research.15096.1>

Saltzman, A. J., Stone, G. W., Claessen, B. E., Narula, A., Leon-Reyes, S., Weisz, G., Mehran, R. (2011). Long-Term Impact of Chronic Kidney Disease in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention. *JACC: Cardiovascular Interventions*, 4(9), 1011–1019. <https://doi.org/10.1016/j.jcin.2011.06.012>

Sanchis-Gomar, F., Perez-Quilis, C., Leischik, R., & Lucia, A. (2016). Epidemiology of Coronary Heart Disease and Acute Coronary Syndrome. *Annals of Translational Medicine*, 4(13), pp.1-12. <http://dx.doi.org/10.21037/atm.2016.06.33>

Sattar, S., Ahmed, N., Akhter, Z., Aijaz, S., Lakhani, S., Malik, R., & Pathan, A. (2019). In-Hospital outcomes in acute coronary syndrome patients with concomitant severe chronic kidney disease undergoing percutaneous coronary intervention. *Pakistan journal of medical sciences*, 35(2), 291–296. <https://doi.org/10.12669/pjms.35.2.276>

Sim, D. S., Jeong, M. H., Ahn, Y., Kim, Y. J., Chae, S. C., Hong, T. J., Park, S. J. (2016). Pharmacoinvasive Strategy Versus Primary Percutaneous Coronary Intervention in Patients With ST-Segment–Elevation Myocardial Infarction. *Circulation: Cardiovascular Interventions*, 9(9), e003508. <https://doi.org/10.1161/circinterventions.115.003508>

Smith, N., Negrelli, M., Manek, B., Hawes, M., & Viera, J. (2015). Diagnosis and Management of Acute Coronary Syndrome: An Evidence-Based Update. *J Am Board Fam*, 28(2), pp.283-293. <http://doi.org/10.3122/jabfm.2015.02.140189>

Stone, S. G., Serrao, G. W., Mehran, R., Tomey, M. I., Witzenbichler, B., Guagliumi, G., Stone, G. W. (2014). Incidence, Predictors, and Implications of Reinfarction After Primary Percutaneous Coronary Intervention in ST-



UNIVERSITAS
GADJAH MADA

Prediktor Infark Miokard Berulang Pascainfark Miokard Akut dengan Elevasi Segmen ST yang Dilakukan

Intervensi Koroner Perkutan di RSUP Dr. Sardjito

PUTRI NOVANTIKA L, Dr. dr. Lucia Kris Dinarti, Sp.PD, Sp.JP (K); dr. Dhite Bayu Nugroho, M.Sc., PhD

Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Segment-Elevation Myocardial Infarction: The Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction Trial. *Circulation: Cardiovascular Interventions*, 7(4), 54,13–551.

<https://doi.org/10.1161/circinterventions.113.001360>

Thygesen, K., Alpert, J., Jaffe, A., Simoons, M., Chaitman, B., & White, H. (2012). Third Universal Definition of Myocardial Infarction. *Circulation*, 126(16), pp.2020-2035. <https://doi.org/10.1161/CIR.0b013e31826e1058>

Tian, L., Yang, Y., Zhu, J., Liu, L., Liang, Y., Li, J., & Yu, B. (2016). Impact of Previous Stroke on Short-Term Myocardial Reinfarction in Patients With Acute ST Segment Elevation Myocardial Infarction. *Medicine*, 95(6), e2742. <https://doi.org/10.1097/md.0000000000002742>

Tillmanns, H., Waas, W., Voss, R., Grempels, E., Hölschermann, H., Haberbosch, W., & Waldecker, B. (2005). *Gender Differences in the Outcome of Cardiac Interventions*. *Herz Kardiovaskuläre Erkrankungen*, 30(5), 375–389. <https://doi.org/10.1007/s00059-005-2716-3>

Turpie, A. (2006). Burden of Disease: Medical and Economic Impact of Acute Coronary Syndromes. *The American Journal of Managed Care*, 12(Supplement 16), pp.S430-S434. <https://www.ncbi.nlm.nih.gov/m/pubmed/17203987>

World Health Organisation. (2013) *Adherence to Long-Term Therapies: Evidence for Action*. Geneva: World Health Organisation.

World Health Organisation. (2016). *International Statistical Classification of Diseases and Related Health Problems*. 10th rev. France: WHO Library Cataloguing-in-Publication Data.