



## References

- Al-Farsi, M. A. and Lee, C. Y. (2008) ‘Optimization of phenolics and dietary fibre extraction from date seeds’, *Food Chemistry*, 108(3), pp. 977–985. doi: 10.1016/j.foodchem.2007.12.009.
- Alam, M. N., Bristi, N. J. and Rafiquzzaman, M. (2013) ‘Review on in vivo and in vitro methods evaluation of antioxidant activity’, *Saudi Pharmaceutical Journal*, 21(2), pp. 143–152. doi: 10.1016/j.jsps.2012.05.002.
- Allwood, J. W., Ellis, D. I. and Goodacre, R. (2008) ‘Metabolomic technologies and their application to the study of plants and plant-host interactions’, *Physiologia Plantarum*, 132(2), pp. 117–135. doi: 10.1111/j.1399-3054.2007.01001.x.
- Avula, B., Joshi, V., Wang, Y. H., Jadhav, A. N. and Khan, I. A. (2008) ‘Quantitative Determination of Ecdysteroids in *Sida rhombifolia* L. and various other *Sida* Species Using LC-UV, and their Anatomical Characterization’, *Natural Product Communications*, 3(5), pp. 705–710.
- Aybastier, Ö., Isik, E., Sahin, S. and Demir, C. (2013) ‘Optimization of ultrasonic-assisted extraction of antioxidant compounds from blackberry leaves using response surface methodology’, *Industrial Crops and Products*, 44, pp. 558–565. doi: 10.1016/j.indcrop.2012.09.022.
- Cacace, J. E. and Mazza, G. (2003) ‘Mass transfer process during extraction of phenolic compounds from milled berries’, *Journal of Food Engineering*, 59(4), pp. 379–389. doi: 10.1016/S0260-8774(02)00497-1.
- Chaves, O. S., Gomes, R. A., Claudia, A., Tomaz, D. A., Fernandes, M. G., Mendes, G., Agra, M. D. F., Braga, V. A. and Souza, M. D. F. V. D. (2013) ‘Secondary Metabolites from *Sida rhombifolia* L. (Malvaceae) and the Vasorelaxant Activity of Cryptolepinone’, pp. 2769–2777. doi: 10.3390/molecules18032769.
- Chemat, F. and Khan, M. K. (2011) ‘Applications of ultrasound in food technology: Processing, preservation and extraction’, *Ultrasonics - Sonochemistry*, 18(4), pp. 813–835. doi: 10.1016/j.ulsonch.2010.11.023.
- Chun, O.K., Kim, D.O. and Lee, C. Y. (2003). *Superoxide Radical Scavenging Activity of the Major Polyphenols in Fresh Plums*. 8067–8072.
- Dai, J. and Mumper, R. J. (2010) ‘Plant phenolics: Extraction, analysis and their antioxidant and anticancer properties’, *Molecules*, 15(10), pp. 7313–7352. doi: 10.3390/molecules15107313.
- Destefanis, G., Barga, M. T., Brugia Paglia, A. and Tassone, S. (2000) ‘The use of principal component analysis (PCA) to characterize beef’, *Meat Science*, 56(3), pp. 255–259. doi: 10.1016/S0309-1740(00)00050-4.
- Dhalwal, K., Deshpande, Y. S. and Purohit, A. P. (2007) ‘Evaluation of in vitro antioxidant activity of *Sida rhombifolia* (L.) Ssp. *retusa* (L.)’, *Journal of Medicinal Food*, 10(4), pp. 683–688. doi: 10.1089/jmf.2006.129.
- Dinda, B., Das, N., Dinda, S., Dinda, M. and SilSarma, I. (2015) ‘The genus *Sida* L . – A traditional medicine : Its ethnopharmacological , phytochemical and pharmacological data for commercial exploitation in herbal drugs industry’, *Journal of Ethnopharmacology*, 176, pp. 135–176. doi:



10.1016/j.jep.2015.10.027.

- Easmin, S., Sarker, M. Z. I., Ghafoor, K., Ferdosh, S., Jaffri, J., Ali, M. E., Mirhosseini, H., Al-Juhaimi, F. Y., Perumal, V. and Khatib, A. (2017) 'Rapid investigation of  $\alpha$ -glucosidase inhibitory activity of Phaleria macrocarpa extracts using FTIR-ATR based fingerprinting', *Journal of Food and Drug Analysis*, 25(2), pp. 306–315. doi: 10.1016/j.jfda.2016.09.007.
- Ellis, D. I., Broadhurst, D., Clarke, S. J. and Goodacre, R. (2005) 'Rapid identification of closely related muscle foods by vibrational spectroscopy and machine learning', *Analyst*, 130(12), pp. 1648–1654. doi: 10.1039/b511484e.
- Ellis, D. I., Broadhurst, D. and Goodacre, R. (2002) 'Rapid and quantitative detection of the microbial spoilage of beef by Fourier transform infrared spectroscopy and machine learning', *Analytica Chimica Acta*, 514(2), pp. 193–201. doi: 10.1016/j.aca.2004.03.060.
- Ellis, D. I., Dunn, W. B., Griffin, J. L., Allwood, J. W. and Goodacre, R. (2007) 'Metabolic fingerprinting as a diagnostic tool', *Pharmacogenomics*, 8(9), pp. 1243–1266. doi: 10.2217/14622416.8.9.1243.
- Ellis, D. I. and Goodacre, R. (2006) 'Metabolic fingerprinting in disease diagnosis: Biomedical applications of infrared and Raman spectroscopy', *Analyst*, 131(8), pp. 875–885. doi: 10.1039/b602376m.
- Ferreira, S. L. C., Bruns, R. E., Ferreira, H. S., Matos, G. D., David, J. M., Brandao, G. C., da Silva, E. G. P., Portugal, L. A., dos Reis, P. S., Souza, A. S. and dos Santos, W. N. L. (2007) 'Box-Behnken design: An alternative for the optimization of analytical methods', 597, pp. 179–186. doi: 10.1016/j.aca.2007.07.011.
- Ferro, M. D., Mazzutti, S., Vitali, L., Muller, C. M. O. and Ferreira, S. R. S. (2019) 'Integrated extraction approach to increase the recovery of antioxidant compounds from *Sida rhombifolia* leaves', *The Journal of Supercritical Fluids*, 149, pp. 10–19. doi: 10.1016/j.supflu.2019.03.013.
- Galvan D'Alessandro, L., Kriaa, K., Nikov, I. and Dimitrov, K. (2012) 'Ultrasound assisted extraction of polyphenols from black chokeberry', *Separation and Purification Technology*, 93, pp. 42–47. doi: 10.1016/j.seppur.2012.03.024.
- Ghasemzadeh, A., Jaafar, H. Z. E., Karimi, E. and Rahmat, A. (2014) 'Optimization of ultrasound-assisted extraction of flavonoid compounds and their pharmaceutical activity from curry leaf (*Murraya koenigii* L.) using response surface methodology', *BMC Complementary and Alternative Medicine*, 14(1). doi: 10.1186/1472-6882-14-318.
- Gupta, M., Paul, S., Karmakar, N., Sasmal, S. and Chowdhury, S. (2016) 'Free Radical Scavenging Activity of *Sida cordifolia* Linn. Extracts Measured by Hydrogen Peroxide, DPPH, ABTS and Ferric Reducing Antioxidant Methods', *International Journal of Current Research in Biosciences and Plant Biology*, 3(8), pp. 114–122. doi: 10.20546/ijcrbp.2016.308.018.
- Heberle, J. and Zscherp, C. (1996) 'ATR/FT-IR difference spectroscopy of biological matter with microsecond time resolution', *Applied Spectroscopy*, 50(5), pp. 588–596. doi: 10.1366/0003702963905907.
- Heydari Majd, M., Rejaei, A., Salar Bashi, D., Ali Mortazavi, S. and Bolourian, S. (2014) 'Optimization of ultrasonic-assisted extraction of phenolic compounds



- from bovine pennyroyal (*Phlomidoschema parviflorum*) leaves using response surface methodology', *Industrial Crops and Products*, 57, pp. 195–202. doi: 10.1016/j.indcrop.2014.03.031.
- Irakli, M., Chatzopoulou, P. and Ekateriniadou, L. (2018) 'Optimization of ultrasound-assisted extraction of phenolic compounds: Oleuropein, phenolic acids, phenolic alcohols and flavonoids from olive leaves and evaluation of its antioxidant activities', *Industrial Crops and Products*, 124(July), pp. 382–388. doi: 10.1016/j.indcrop.2018.07.070.
- Iswantini, D., Darusman, L. K. and Hidayat, R. (2009) 'Indonesian Sidaguri (*Sida rhombifolia* L.) as antigout and inhibition kinetics of flavonoids crude extract on the activity of xanthine oxidase.', *Journal of Biological Sciences*, 9, pp. 504–508. doi: 10.3923/jbs.2009.504.508.
- Jakobek, L., Boc, M., & Barron, A. R. (2015). Optimization of Ultrasonic-Assisted Extraction of Phenolic Compounds from Apples. *Food Analytical Methods*, 8(10), 2612–2625. <https://doi.org/10.1007/s12161-015-0161-3>
- Johnson, H. E., Broadhurst, D., Kell, D. B., Theodorou, M. K., Merry, R. J. and Griffith, G. W. (2004) 'High-Throughput Metabolic Fingerprinting of Legume Silage Fermentations via Fourier Transform Infrared Spectroscopy and Chemometrics', 70(3), pp. 1583–1592. doi: 10.1128/AEM.70.3.1583.
- Khoddami, A., Wilkes, M. A. and Roberts, T. H. (2013) 'Techniques for analysis of plant phenolic compounds', *Molecules*, 18(2), pp. 2328–2375. doi: 10.3390/molecules18022328.
- Lee, J. E., Lee, B. J., Chung, J. O., Hwang, J. A., Lee, S. J., Lee, C. H. and Hong, Y. S. (2010) 'Geographical and Climatic Dependencies of Green Tea (*Camellia sinensis*) Metabolites: A <sup>1</sup>H NMR-Based Metabolomics Study', *Journal of Agricultural and Food Chemistry*, 58(19), pp. 10582–10589. doi: 10.1021/jf102415m.
- Li, H., Gong, X., Wang, Z., Pan, C., Zhao, Y., Gao, X. and Liu, W. (2019) 'Multiple fingerprint profiles and chemometrics analysis of polysaccharides from *Sarcandra glabra*', *International Journal of Biological Macromolecules*, 123, pp. 957–967. doi: 10.1016/j.ijbiomac.2018.11.103.
- Liu, C. W., Wang, Y. C., Lu, H. C. and Chiang, W. D. (2014) 'Optimization of ultrasound-assisted extraction conditions for total phenols with anti-hyperglycemic activity from *Psidium guajava* leaves', *Process Biochemistry*, 49(10), pp. 1601–1605. doi: 10.1016/j.procbio.2014.06.009.
- Mah, S. H., Teh, S. S. and Ee, G. C. L. (2017) 'Anti-inflammatory, anti-cholinergic and cytotoxic effects of *Sida Rhombifolia*', *Pharmaceutical Biology*, 55(1), pp. 920–928. doi: 10.1080/13880209.2017.1285322.
- Mason, T. J. (1997) 'Ultrasound in synthetic organic chemistry', *Chemical Society Reviews*, 26(6), pp. 443–451. doi: 10.1039/cs9972600443.
- Miller, J. N. and Miller, J. C. (2005) *Chemometrics for Analytical Chemistry, Analytical Chemistry*.
- Muñiz-Márquez, D. B., Martinez-Avila, G. C., Wong-Paz, J. E., Belmares-Cerda, R., Rodriguez-Herrera, R. and Aguilar, C. N. (2013) 'Ultrasound-assisted extraction of phenolic compounds from *Laurus nobilis* L. and their antioxidant activity', *Ultrasonics Sonochemistry*, 20(5), pp. 1149–1154. doi:



- 10.1016/j.ulstsonch.2013.02.008.
- Muritala, H. F., Akolade, J. O., Akande, S. A., Abdulazeez, A. T., Aladodo, R. A., & Bello, A. B. (2018). Antioxidant and alpha-amylase inhibitory potentials of *Cocos nucifera* husk. *Food Science and Nutrition*, 6(6), 1676–1683. <https://doi.org/10.1002/fsn3.741>
- Naes, T., Baardseth, P., Helgesen, H. and Isaksson, T. (1996) ‘Multivariate techniques in the analysis of meat quality’, *Meat Science*, 43(SUPPL. 1), pp. 135–149. doi: 10.1016/0309-1740(96)00061-7.
- Navas, M., Dan, M. and Gopalakrishnan, P. (2013) ‘Comparative root anatomy of the species under *Sida rhombifolia* complex ( Malvaceae )’, *Pharmacognosy Journal*, 5(6), pp. 269–274. doi: 10.1016/j.phcogj.2013.09.004.
- Nikalje, G. C., Kumar, J., Nikam, T. D. and Suprasannaa, P. (2019) ‘FT-IR profiling reveals differential response of roots and leaves to salt stress in a halophyte *Sesuvium portulacastrum* (L.) L.’, *Biotechnology Reports*, 23, p. e00352. doi: 10.1016/j.btre.2019.e00352.
- Nour, V., Trandafir, I. and Cosmulescu, S. (2016) ‘Optimization of ultrasound-assisted hydroalcoholic extraction of phenolic compounds from walnut leaves using response surface methodology’, *Pharmaceutical Biology*, 54(10), pp. 2176–2187. doi: 10.3109/13880209.2016.1150303.
- Panda, D. and Manickam, S. (2019) ‘applied sciences Cavitation Technology — The Future of Greener Extraction Method : A Review on the Extraction of Natural Products and Process Intensification Mechanism and Perspectives’. doi: 10.3390/app9040766.
- Pollini, L., Tringaniello, C., Ianni, F., Blasi, F., Manes, J. and Cossignani, L. (2020) ‘Impact of ultrasound extraction parameters on the antioxidant properties of *Moringa oleifera* leaves’, *Antioxidants*, 9(4), pp. 1–14. doi: 10.3390/antiox9040277.
- Prommuak, C., De-Eknamkul, W. and Shotipruk, A. (2008) ‘Extraction of flavonoids and carotenoids from Thai silk waste and antioxidant activity of extracts’, *Separation and Purification Technology*, 62(2), pp. 444–448. doi: 10.1016/j.seppur.2008.02.020.
- Rifai, Y., Djabir, Y. Y. and Alam, G. (2020) ‘Chemometric Analysis of Arbutin Derivatives from *Paederia foetida* and *Vitis vinifera* with Fourier Transform Infrared ( FTIR )’, 12(3), pp. 436–441.
- Rivera-Mondragón, A., Broeckx, G., Bijntebier, S., Naessens, T., Fransen, E., Kiekens, F., Caballero-George, C., Vander Heyden, Y., Apers, S., Pieters, L. and Fouber, K. (2019) ‘Ultrasound-assisted extraction optimization and validation of an HPLC-DAD method for the quantification of polyphenols in leaf extracts of *Cecropia* species’, *Scientific Reports*, 9(1), pp. 1–16. doi: 10.1038/s41598-018-37607-2.
- Şahin, S. and Şamlı, R. (2013) ‘Optimization of olive leaf extract obtained by ultrasound-assisted extraction with response surface methodology’, *Ultrasonics Sonochemistry*, 20(1), pp. 595–602. doi: 10.1016/j.ulstsonch.2012.07.029.
- Saifullah, M., McCullum, R., McCluskey, A. and Vuong, Q. (2020) ‘Comparison of conventional extraction technique with ultrasound assisted extraction on



- recovery of phenolic compounds from lemon scented tea tree (*Leptospermum petersonii*) leaves', *Helijon*, 6(4), p. e03666. doi: 10.1016/j.heliyon.2020.e03666.
- Setford, P. C., Jeffery, D. W., Grbin P. R. and Muhlack, R. A. (2017) 'Factors affecting extraction and evolution of phenolic compounds during red wine maceration and the role of process modelling', *Trends in Food Science and Technology*, 69, pp. 106–117. doi: 10.1016/j.tifs.2017.09.005.
- Setyaningsih, W., Duros, E., Palma, M., and Barroso, C. G. (2015) 'Optimization of the ultrasound-assisted extraction of melatonin from red rice (Oryza sativa) grains through a response surface methodology', *APPLIED ACOUSTICS*, pp. 1–7. doi: 10.1016/j.apacoust.2015.04.001.
- Setyaningsih, W., Saputro, I. E., Carrera, C. A. and Palma, M. (2019) 'Optimisation of an ultrasound-assisted extraction method for the simultaneous determination of phenolics in rice grains', *Food Chemistry*, 288(October 2018), pp. 221–227. doi: 10.1016/j.foodchem.2019.02.107.
- Shirzad, H., Niknam, V., Taheri, M. and Ebrahimzadeh, H. (2017) 'Ultrasound-assisted extraction process of phenolic antioxidants from Olive leaves: a nutraceutical study using RSM and LC–ESI–DAD–MS', *Journal of Food Science and Technology*, 54(8), pp. 2361–2371. doi: 10.1007/s13197-017-2676-7.
- Subramanya, M. D., Pai, S. R., Upadhyaya, V., Ankad, G. M., Bhagwat, S. S. and Hegde, H. V. (2015) 'Total polyphenolic contents and in vitro antioxidant properties of eight *Sida* species from Western Ghats, India', *Journal of Ayurveda and Integrative Medicine*, 6(1), pp. 24–28. doi: 10.4103/0975-9476.146544.
- Tao, Y., Wu, D. and Zhang, Q. (2013) 'Ultrasound-assisted extraction of phenolics from wine lees: Modeling, optimization and stability of extracts during storage'. doi: 10.1016/j.ulsonch.2013.09.005.
- Tiwari, B. K. (2015) 'Ultrasound: A clean, green extraction technology', *TrAC - Trends in Analytical Chemistry*, 71, pp. 100–109. doi: 10.1016/j.trac.2015.04.013.
- Upadhyay, R., Nachiappan, G. and Mishra, H. N. (2015) 'Ultrasound-assisted extraction of flavonoids and phenolic compounds from *Ocimum tenuiflorum* leaves', *Food Science and Biotechnology*, 24(6), pp. 1951–1958. doi: 10.1007/s10068-015-0257-y.
- Wang, L. and Weller, C. L. (2006) 'Recent advances in extraction of nutraceuticals from plants', *Trends in Food Science and Technology*, 17(6), pp. 300–312. doi: 10.1016/j.tifs.2005.12.004.
- Wang, Y., Avula, B., Jadhav, A. N., Smillie, T. J. and Khan, I. A. (2008) 'Structural characterization and identification of ecdysteroids from *Sida rhombifolia* L. in positive electrospray ionization by tandem mass spectrometry', pp. 2413–2422. doi: 10.1002/rcm.
- William-Allwood, J. J., Ellis, D. I., Heald, J. K. Goodacre, R. and Mur, L. A. J. (2006) 'Metabolomic approaches reveal that phosphatidic and phosphatidyl glycerol phospholipids are major discriminatory non-polar metabolites in responses by *Brachypodium distachyon* to challenge by *Magnaporthe grisea*',



*Plant Journal*, 46(3), pp. 351–368. doi: 10.1111/j.1365-313X.2006.02692.x.

- Winder, C. L. and Goodacre, R. (2004) ‘Comparison of diffuse-reflectance absorbance and attenuated total reflectance FT-IR for the discrimination of bacteria’, *Analyst*, 129(11), pp. 1118–1122. doi: 10.1039/b408169b.
- Xia, E. Q., Ai, X. X., Zang, S. Y., Guan, T. T., Xu, X. R. and Li, H. B. (2011) ‘Ultrasound-assisted extraction of phillyrin from *Forsythia suspensa*’, *Ultrasonics Sonochemistry*, 18(2), pp. 549–552. doi: 10.1016/j.ultsonch.2010.09.015.

Yang, F. Chu, T., Zhang, Y., Liu, X., Sun, G. and Chen, Z. (2020) ‘Quality assessment of licorice (*Glycyrrhiza glabra* L.) from different sources by multiple fingerprint profiles combined with quantitative analysis, antioxidant activity and chemometric methods’, *Food Chemistry*, 324(April), p. 126854. doi: 10.1016/j.foodchem.2020.126854.

Zamora-Gasga, V. M. Serafín-García, M. S., Sánchez-Burgos, J. A., Estrada, R. M. V. and Sáyago-Ayerdi, S. G.. (2017) ‘Optimization of Ultrasonic-Assisted Extraction of Antioxidant Compounds from Starfruit (*Averrhoa carambola* L) Leaves’, *Journal of Food Processing and Preservation*, 41(4), pp. 1–10. doi: 10.1111/jfpp.13093.