



DAFTAR PUSTAKA

- Abdulla, J., Barlera, S., Latini, R., Kjoller-Hansen, L., Sogaard, P., Christensen, E., Kober, L. and Torp-Pedersen, C. (2007). A systematic review: Effect of angiotensin converting enzyme inhibition on left ventricular volumes and ejection fraction in patients with a myocardial infarction and in patients with left ventricular dysfunction. *European Journal of Heart Failure*, 9(2), pp.129-135.
- Afari, M. and Bhat, T. (2016). Neutrophil to lymphocyte ratio (NLR) and cardiovascular diseases: an update. *Expert Review of Cardiovascular Therapy*, 14(5), pp.573-577.
- Akpek, M., Kaya, M.G., Lam, Y.Y., Sahin, O., Elcik, D., Celik, T., Ergin, A. and Gibson, C.M. (2012). Relation of Neutrophil/Lymphocyte Ratio to Coronary Flow to In-Hospital Major Adverse Cardiac Events in Patients With ST-Elevated Myocardial Infarction Undergoing Primary Coronary Intervention. *The American Journal of Cardiology*, 110(5), pp.621–627.
- Angkananard, T., Anothaisintawee, T., Ingsathit, A., McEvoy, M., Silapat, K., Attia, J., Sritara, P. and Thakkinstian, A., 2019. Mediation Effect of Neutrophil Lymphocyte Ratio on Cardiometabolic Risk Factors and Cardiovascular Events. *Scientific Reports*, 9(1).
- Angkananard, T., Anothaisintawee, T., McEvoy, M., Attia, J. and Thakkinstian, A., 2018. Neutrophil Lymphocyte Ratio and Cardiovascular Disease Risk: A Systematic Review and Meta-Analysis. *BioMed Research International*, 2018, pp.1-11.
- Arbel, Y., Shacham, Y., Ziv-Baran, T., Laufer Perl, M., Finkelstein, A., Halkin, A., et al. (2014). Higher Neutrophil/Lymphocyte Ratio Is Related to Lower Ejection Fraction and Higher Long-term All-Cause Mortality in ST-Elevation Myocardial Infarction Patients. *Canadian Journal of Cardiology*, 30(10), pp.1177-1182.
- Bagaswoto, H., (2015). Sistem Skoring Berdasarkan Gambaran Elektrokardiogram Untuk Memprediksi Jenis Gagal Jantung Pada Pasien Gagal Jantung Kronik. *Tesis*. Universitas Gadjah Mada.
- Balta, S., Celik, T., Mikhailidis, D.P., Ozturk, C., Demirkol, S., Aparci, M. and Iyisoy, A. (2015). The Relation Between Atherosclerosis and the Neutrophil–Lymphocyte Ratio. *Clinical and Applied Thrombosis/Hemostasis*, 22(5), pp.405–411.
- Bellenger, N., Burgess, M., Ray, S., Lahiri, A., Coats, A., Cleland, J. and Pennell, D., (2000). Comparison of left ventricular ejection fraction and volumes in heart failure by echocardiography, radionuclide ventriculography and cardiovascular



magnetic resonance. Are they interchangeable?. *European Heart Journal*, 21(16), pp.1387-1396.

Bentzon, J., Otsuka, F., Virmani, R. and Falk, E. (2014). Mechanisms of Plaque Formation and Rupture. *Circulation Research*, 114(12), pp.1852-1866.

Budzianowski, J., Pieszko, K., Burchardt, P., Rzeźniczak, J. and Hiczkiewicz, J. (2017). The Role of Hematological Indices in Patients with Acute Coronary Syndrome. *Disease Markers*, 2017, pp.1-9.

Carrick, D., Haig, C., Maznyczka, A.M., Carberry, J., Mangion, K., Ahmed, N., Yue May, V.T., McEntegart, M., Petrie, M.C., Eteiba, H., Lindsay, M., Hood, S., Watkins, S., Davie, A., Mahrouse, A., Mordi, I., Ford, I., Radjenovic, A., Welsh, P., Sattar, N., Wetherall, K., Oldroyd, K.G. and Berry, C. (2018). Hypertension, Microvascular Pathology, and Prognosis After an Acute Myocardial Infarction. *Hypertension*, 72(3), pp.720–730.

Chen, C., Cong, B., Wang, M., Abdullah, M., Wang, X., Zhang, Y., et al. (2018). Neutrophil to lymphocyte ratio as a predictor of myocardial damage and cardiac dysfunction in acute coronary syndrome patients. *Integrative Medicine Research*, 7(2), pp.192-199.

Chen, C., Zhao, H. and Zhang, Y., 2019. Correlation between neutrophil-to-lymphocyte ratio and kidney dysfunction in undiagnosed hypertensive population from general health checkup. *The Journal of Clinical Hypertension*, 22(1).

Chen, Z.-W., Yu, Z.-Q., Yang, H.-B., Chen, Y.-H., Qian, J.-Y., Shu, X.-H. and Ge, J.-B. (2016). Rapid predictors for the occurrence of reduced left ventricular ejection fraction between LAD and non-LAD related ST-elevation myocardial infarction. *BMC Cardiovascular Disorders*, 16(1).

Costanzo, L. (2014). *Physiology*. 5th ed. Philadelphia: Saunders Elsevier.

Dahlan, M. (2014). *Statistik untuk Kedokteran dan Kesehatan: Deskriptif, Bivariat, dan Multivariat, Dilengkapi Aplikasi Menggunakan SPSS*. 6th ed. Jakarta: Epidemiologi Indonesia.

Dubey, G., Verma, S. and Bahl, V. (2017). Primary percutaneous coronary intervention for acute ST elevation myocardial infarction: Outcomes and determinants of outcomes: A tertiary care center study from North India. *Indian Heart Journal*, 69(3), pp.294-298.

Duffy, B.K., Gurm, H.S., Rajagopal, V., Gupta, R., Ellis, S.G., Bhatt, D.L. (2006). Usefulness of an elevated neutrophil to lymphocyte ratio in predicting long-term mortality after percutaneous coronary intervention. *Am J Cardiol*. 2006;97:993–996.



- Dong, C., Wang, Z. and Chen, S. (2018). Neutrophil to lymphocyte ratio predict mortality and major adverse cardiac events in acute coronary syndrome: A systematic review and meta-analysis. *Clinical Biochemistry*, 52, pp.131-136.
- Ehl, N.F., Kühne, M., Brinkert, M., Müller-Brand, J. and Zellweger, M.J. (2011). Diabetes reduces left ventricular ejection fraction-irrespective of presence and extent of coronary artery disease. *European Journal of Endocrinology*, 165(6), pp.945–951.
- Erre, G., Paliogiannis, P., Castagna, F., Mangoni, A., Carru, C., Passiu, G. and Zinelli, A., (2018). Meta-analysis of neutrophil-to-lymphocyte and platelet-to-lymphocyte ratio in rheumatoid arthritis. *European Journal of Clinical Investigation*, 49(1), p.e13037.
- Ethier, J., Desautels, D., Templeton, A., Shah, P. and Amir, E. (2017). Prognostic role of neutrophil-to-lymphocyte ratio in breast cancer: a systematic review and meta-analysis. *Breast Cancer Research*, 19(1).
- Foley, T., Mankad, S., Anavekar, N., Bonnichsen, C., Miller, M., Morris, T. and Araoz, P., (2012). Measuring Left Ventricular Ejection Fraction – Techniques and Potential Pitfalls. *European Cardiology Review*, 8(2), p.108.
- Forget, P., Khalifa, C., Defour, J., Latinne, D., Van Pel, M. and De Kock, M. (2017). What is the normal value of the neutrophil-to-lymphocyte ratio?. *BMC Research Notes*, 10(1).
- Foth, C. and Mountfort, S. (2019). *Acute Myocardial Infarction ST Elevation (STEMI)*. [online] Ncbi.nlm.nih.gov. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK532281/> [Accessed 8 Feb. 2020].
- Gulbagci, B., Sametoglu, F., Cengiz, H. and Varim, C., (2020). Evaluation of the relationship of disease activity with neutrophile to lymphocyte ratio and platelet to lymphocyte ratio in patients with peptic ulcer and gastritis. *International Journal of Research in Medical Sciences*, 8(11), p.3850.
- Guo, X., Zhang, S., Zhang, Q., Liu, L., Wu, H., Du, H., Shi, H., Wang, C., Xia, Y., Liu, X., Li, C., Sun, S., Wang, X., Zhou, M., Huang, G., Jia, Q., Zhao, H., Song, K. and Niu, K., (2015). Neutrophil:lymphocyte ratio is positively related to type 2 diabetes in a large-scale adult population: a Tianjin Chronic Low-Grade Systemic Inflammation and Health cohort study. *European Journal of Endocrinology*, 173(2), pp.217-225.
- Han, Y., Yang, T., Kim, D., Jin, H., Chung, S., Seo, J., et al. (2013). Neutrophil to Lymphocyte Ratio Predicts Long-Term Clinical Outcomes in Patients with ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. *Korean Circulation Journal*, 43(2), p.93.
- Hareen, T., Bhaskaran, A. and Jaswanthi, A., (2016). Neutrophil to lymphocyte ratio in diagnosing acute cholecystitis: a retrospective cohort study in a tertiary rural hospital. *International Surgery Journal*, 4(1), p.372.



Hendriks, T., van Dijk, R., Alsabaan, N.A. *et al.* (2020). Active Tobacco Smoking Impairs Cardiac Systolic Function. *Sci Rep*, 10.

Hong, D., Choi, K., Song, Y., Lee, J., Park, T., Yang, J., Hahn, J., Choi, J., Choi, S., Kim, S., Choe, Y., Kim, E., Chang, S., Lee, S., Oh, J. and Gwon, H., (2019). Prognostic implications of post-percutaneous coronary intervention neutrophil-to-lymphocyte ratio on infarct size and clinical outcomes in patients with acute myocardial infarction. *Scientific Reports*, 9(1).

Ibanez, B., James, S., Agewall, S., Antunes, M., Bucciarelli-Ducci, C., Bueno, H., *et al.* (2017). 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *European Heart Journal*, 39(2), pp.119-177.

Jahic, E. (2017). Experience and Outcomes of Primary Percutaneous Coronary Intervention for Patients with ST-Segment Elevation Myocardial Infarction of Tertiary Care Center in Bosnia and Herzegovina. *Medical Archives*, 71(3), p.183.

Jan, S., Lee, S., Sawhney, J., Ong, T., Chin, C., Kim, H., *et al.* (2016). Catastrophic health expenditure on acute coronary events in Asia: a prospective study. *Bulletin of the World Health Organization*, 94(3), pp.193-200.

Jayaraj, J., Davatyan, K., Subramanian, S. and Priya, J., 2018. Epidemiology of Myocardial Infarction. *Myocardial Infarction*.

Kalra, S., Bhatt, H. and Kirtane, A. (2018). Stenting in Primary Percutaneous Coronary Intervention for Acute ST-Segment Elevation Myocardial Infarction. *Methodist Debakey Cardiovascular Journal*, 14(1).

Karakas, M.S., Korucuk, N., Tosun, V., Altekin, R.E., Koç, F., Ozbek, S.C., Ozel, D. and Ermis, C. (2016). Red cell distribution width and neutrophil-to-lymphocyte ratio predict left ventricular dysfunction in acute anterior ST-segment elevation myocardial infarction. *Journal of the Saudi Heart Association*, 28(3), pp.152–158.

Kazemi, T., Mohseni, J., Maleki, M. and Beydokhti, H. (2017). A systematic review on the prevalence of acute myocardial infarction in Iran. *Heart Views*, 18(4), p.125.

Kementerian Kesehatan Republik Indonesia (2018). *Hasil Utama Riskesdas 2018*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Republik Indonesia.

Khaled, S. and Matahen, R. (2018). Cardiovascular risk factors profile in patients with acute coronary syndrome with particular reference to left ventricular ejection fraction. *Indian Heart Journal*, 70(1), pp.45-49.

Kırış, T., Avcı, E. and Çelik, A. (2018). Combined value of left ventricular ejection fraction and the Model for End-Stage Liver Disease (MELD) score for



predicting mortality in patients with acute coronary syndrome who were undergoing percutaneous coronary intervention. *BMC Cardiovascular Disorders*, 18(1).

Kosaraju, A. and Makaryus, A. (2020). *Left Ventricular Ejection Fraction*. [online] Ncbi.nlm.nih.gov. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK459131/> [Accessed 12 Feb. 2020].

Lilly, L., 2016. *Pathophysiology Of Heart Disease: A Collaborative Project Of Medical Students And Faculty*. 6th ed. Philadelphia: Lippincott, Williams & Wilkins.

Loyeau, A., Benamer, H., Bataille, S., Tepper, S., Boche, T., Lamhaut, L., et al., 2018. Evolution of ST-Elevation Acute Myocardial Infarction Prevalence by Gender Assessed Age Pyramid Analysis—The Piramyd Study. *Journal of Clinical Medicine*, 7(12), p.509.

McNair, P.W., Bilchick, K.C. and Keeley, E.C. (2019). Very late presentation in ST elevation myocardial infarction: Predictors and long-term mortality. *IJC Heart & Vasculature*, 22, pp.156–159.

Montecucco, F., Carbone, F. and Schindler, T. (2015). Pathophysiology of ST-segment elevation myocardial infarction: novel mechanisms and treatments. *European Heart Journal*, 37(16), pp.1268-1283.

Mozaffarian, D., Benjamin, E., Go, A., Arnett, D., Blaha, M., Cushman, et al. (2016). Heart Disease and Stroke Statistics—2016 Update. *Circulation*, 133(4).

Nito, I., Waspadji, S., Harun, S. and Markum, H., (2004). Correlation Between Cortisol Levels and Myocardial Infarction Mortality Among Intensive Coronary Care Unit Patients during First Seven Days in Hospital. *Acta Med Indonesia*, 36(1), pp.8-14.

Núñez, J., Núñez, E., Bodí, V., Sanchis, J., Mainar, L., Miñana, G., Fácila, L., Bertomeu, V., Merlos, P., Darmofal, H., Palau, P. and Llácer, A., (2010). Low lymphocyte count in acute phase of ST-segment elevation myocardial infarction predicts long-term recurrent myocardial infarction. *Coronary Artery Disease*, 21(1), pp.1-7.

Nugraha, I.W., Hartopo, A.B. and Taufiq, N. (2020). Wire Crossing Time Correlate with Left Ventricular End-Diastolic Pressure in Patients with ST Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. *Indonesian Journal of Cardiology*.

Nursidiq, A., Purwaningtyas, N. and Wasyanto, T., (2018). Association Between Neutrophil to Lymphocyte Ratio and Left Ventricle Global Longitudinal Strain in Acute Myocardial Infarction. *ACI (Acta Cardiologia Indonesiana)*, 4(2), pp.80-87.

Ozaki, Y., Imanishi, T., Tanimoto, T., Teraguchi, I., Nishiguchi, T., Orii, M., Shiono, Y., Shimamura, K., Yamano, T., Ino, Y., Yamaguchi, T., Kubo, T. and



- Akasaka, T., (2014). Effect of Direct Renin Inhibitor on Left Ventricular Remodeling in Patients With Primary Acute Myocardial Infarction. *International Heart Journal*, 55(1), pp.17-21.
- Peng, Y., Li, Y., He, Y., Wei, Q., Xie, Q., Zhang, L., et al. (2018). The role of neutrophil to lymphocyte ratio for the assessment of liver fibrosis and cirrhosis: a systematic review. *Expert Review of Gastroenterology & Hepatology*, 12(5), pp.503-513.
- Perhimpunan Dokter Spesialis Kardiovaskular Indonesia (2015). *Pedoman Tatalaksana Sindrom Koroner Akut Edisi Ketiga*. Centra Communications.
- PERKENI (2015). *Konsensus Pengelolaan Dan Pencegahan Diabetes Melitus Tipe 2 Di Indonesia*. 1st ed. PB PERKENI.
- Perhimpunan Dokter Spesialis Kardiovaskular Indonesia (2015). *Pedoman Tatalaksana Hipertensi pada Penyakit Kardiovaskular*. 1st ed.
- Perhimpunan Dokter Spesialis Kardiovaskular Indonesia. (2016). *Panduan Praktik Klinis (PPK) dan Clinical Pathway (CP) Penyakit Jantung dan Pembuluh Darah*. 1st Ed.
- Rosales, C. (2018). Neutrophil: A Cell with Many Roles in Inflammation or Several Cell Types?. *Frontiers in Physiology*, 9.
- Roth, G., Johnson, C., Abajobir, A., Abd-Allah, F., Abera, S., Abyu, G., et al. (2017). Global, Regional, and National Burden of Cardiovascular Diseases for 10 Causes, 1990 to 2015. *Journal of the American College of Cardiology*, 70(1), pp.1-25.
- Serrao, G., Lansky, A., Mehran, R. and Stone, G. (2018). Predictors of Left Ventricular Ejection Fraction Improvement After Primary Stenting in ST-Segment Elevation Myocardial Infarction (from the Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction Trial). *The American Journal of Cardiology*, 121(6), pp.678-683.
- Steg, P. G., James, S. K., Badano, L. P., Borger, M. A., Blomstorm-Lundqvist, C., Mario, C., et al. (2012). ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation The Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology. *European Heart Journal*. 33:2569–2619.
- Tahto, E., Jadric, R., Pojskic, L. and Kicic, E. (2017). Neutrophil-to-lymphocyte Ratio and Its Relation with Markers of Inflammation and Myocardial Necrosis in Patients with Acute Coronary Syndrome. *Medical Archives*, 71(5), p.312.
- Templeton, A., McNamara, M., Šeruga, B., Vera-Badillo, F., Aneja, P., Ocaña, A., et al. (2014). Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Solid



UNIVERSITAS
GADJAH MADA

Hubungan antara Neutrophil-to-Lymphocyte Ratio terhadap Fraksi Ejeksi Ventrikel Kiri pada Pasien Infark Miokardium Akut dengan Elevasi Segmen ST yang Dilakukan Intervensi Koroner Perkutan Primer

KARUNIA WIDHI A P, Budi Yuli Setianto; Hendry Purnasidha Bagaswoto

Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Tumors: A Systematic Review and Meta-Analysis. *JNCI: Journal of the National Cancer Institute*, 106(6).

Vaduganathan, M., Ambrosy, A., Greene, S., Mentz, R., Subacius, H., Maggioni, A., et al. (2012). Predictive Value of Low Relative Lymphocyte Count in Patients Hospitalized for Heart Failure With Reduced Ejection Fraction. *Circulation: Heart Failure*, 5(6), pp.750-758.

Vogel, B., Claessen, B., Arnold, S., Chan, D., Cohen, D., Giannitsis, E., et al. (2019). ST- segment elevation myocardial infarction. *Nature Review*, 5(39).

Wang, Q., Ma, J., Jiang, Z. and Ming, L. (2018). Prognostic Value of Neutrophil-To-Lymphocyte Ratio and Platelet-To-Lymphocyte Ratio in Acute Pulmonary Embolism: A Systematic Review and Meta-Analysis. *Minerva Medica*, 37(1), pp.4-11.

Who.int. (2020). *The top 10 causes of death*. [online] Available at:

<https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>
[Accessed 8 Feb. 2020].