

Daftar Pustaka

- Ades, P.A., Keteyian, J.S., Balady, G.J., Houston-Miller, N. 2014. Cardiac Rehabilitation Exercise and Self Care for Chronic Heart Failure. *jchf*.540-547
- Abassia, Z., Karramb, T., Ellahamc, S., Winaver, J. and Hoffman, A. (2004) Implications of the natriuretic peptide system in the pathogenesis of heart failure: diagnostic and therapeutic importance. *Pharmacol Therap.* 102, 223–241
- Alphonsus C, Govender P, Rodseth R, Biccard B. 2019, The Role of Cardiac rehabilitation using exercise to decrease natriuretic peptide levels in non-surgical patients: a systematic review. *Perioperative Medicine* 8:14
- Arena, R., Myers, J., William, M.A., Gulati, M., Kligfield, P., Balady, G.J., Collins, E., Fletcher, G. 2007. Assessment of Functional Capacity in Clinical and Research Settings. *pub4*. CD007130
- Balady, G.J., Williams, M.A., Ades, P.A., Bittner, V., Comoss, P., Foody, J.M. 2007. Core components of cardiac rehabilitation/secondary prevention programs: update. *Circulation.* 115(20):2675–82.
- Beswick, A.D., Rees, K., Griebisch, I., Taylor, F.C., Burke, M., West, R.R. 2004. Provision, uptake and cost of cardiac rehabilitation programmes: improving services to under-represented groups. *Health Technology Assessment.* 8(41):1–166.
- Berent R, von Duvillard SP, Crouse SF, Auer J, Green JS, Sinzinger H. 2009, Short term cardiac rehabilitation reduces B-Type Natriuretic peptides. *Eur J Cardiovasc Prev Rehabil* 16:603-608
- BPJS Kesehatan. 2018. Panduan Praktis Pelayanan Kesehatan. Jakarta: BPJS Kesehatan
- Buckingham, S.A., Taylor, R.S., Jolly, K., Zawada, A., Dean, S.G., Cowie, A., Norton, R.J., Dalal, H.M. 2016. Home-based versus centre-based cardiac rehabilitation: Abridged Cochrane systematic review and meta-analysis. *openhrt*.000463
- Chamberlain, R.S., Sond, J., Mahendraraj, K., Lau, C.S., Siracuse, B. L. 2018 Determining 30-day readmission risk for heart failure patients: the Readmission After Heart Failure scale. *Int J Gen Med*;11:127-141
- Charloux, A., Piquard, F., Doutreleau, S., Brandenberger, G. and Geny, B. (2003) Mechanisms of renal hyporesponsiveness to ANP in heart failure. *Eur. J. Clin. Invest.* 33, 769–778
- Choi BG, Rha SW., Yoon SG., Choi CU., Lee MW 2019. Association of Major Adverse Cardiac Events up to 5 Years in Patients With Chest Pain Without Significant Coronary Artery Disease in The Korean Population. *J Am Hear Assoc*18;8(12)

- Clark, R.A., Conway, A., Poulsen, V., Keech, W., Tirimacco, R., Tideman, P. 2015. Alternative models of cardiac rehabilitation: a systematic review. *Eur J Prev Cardiol* .22(1):35-74
- Currell, R., Urquhart, C., Wainwright, P., Lewis, R. 2000. Telemedicine versus face to face patient care: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev*. 2:CD002098
- Dewi IK, Aminuddin M, Zulkarnain BS. 2016. Analysis of Change in NT-PrBNP after angiotensin receptor bloker therapy in patient with heart failure. *Fol Med Indo*. Vol 52.
- Di Angelantonio E, Chowdhury R, Sarwar N, Ray KK, Gobin R, Saleheen D et al. 2009 B-type natriuretic peptides and cardiovascular risk: systematic review and meta-analysis of 40 prospective studies. *Circ Cardiovasc QualOutcomes*120:2177–2187.
- Felker G, Whellan D, Kraus W, Clare R, Zannad F, DOnahue M et al. 2009, N-Terminal Pro-Brain Natriuretic Peptide and exercise capacity in chronic heart failure: data from HF-Action study. *Am Heart J*. 158(4 0): S37-S44
- Fleg, J.L., Cooper, L.S., Levine, B.D., Pfeffer, M.A., Piña, I.L., 2016. Exercise Training as Therapy for Heart Failure: Current Status and Future Directions. *circheartfailure*.113.001420
- Francis,G.S.,Benedict,C.,Johnstone,D.E.,Kirlin,P.C.,Nicklas, J., Liang, C.S., Kubo, S.H., Rudin-Toretsky, E. and Yusuf, S. 1990 Comparison of neuroendocrine activation in patients with left ventricular dysfunction with and without congestive heart failure. A substudy of the Studies of Left Ventricular Dysfunction (SOLVD). *Circulation* 82, 1724–1729
- Fu TC, Huang SC, Hsu, CC, Wang CH, Wang, JS, 2013. Cardiac Rehabilitation in Patients with Heart Failure. *Acta Cardiol Sin*. 30:353-359
- Gardner RS, Henderson G, McDonagh TA. 2005, The prognostic use of right heart catheterization data in patients with advanced heart failure: how relevant are invasive procedures in the risk stratification of advanced heart failure in the era of neurohormones? *J Heart Lung Transplant* 24: 303–9.
- Goldsmith, S. 2004 Interactions between the sympathetic nervous system and the RAAS in heart failure. *Curr Heart Fail Rep* 1: 45–50^[SEP]
- Giallauria F, De lorenzo A, Pileri F, Manakos M, Lucci R, Psaroudaki M et al. 2006. Reduction of N Terminal-ProBNP with exercise based cardiac rehabilitation in patient with left ventricular dysfunction after myocardial infarction *Eur Jour Card Prev Rehab*: 13, 625-632
- Guazzi M, Dickstein K, Vicenzi M, Arena,R. 2009. 6-Minute walk test and cardiopulmonary exercise testing in patient with chronic heart failure. *Circ Heart Fail*.;549-555
- Gundogdu F, Bozkurt E, Kizilunc A, Sevimli S, Arslan S, Gurlertop Y. 2007. The effect of beta blocker (carvedilol) therapy on terminal pro BN{ levels and echocardiographic findings in patients with congestive heart failure. *pubmed.gov* 24(2):113-7

- Hwang, R, Bruning, J, Morris, NR, Mandrusiak, A, Russel, T. Home-based telerehabilitation is not inferior to a centre-based program in patients with chronic heart failure: a randomised trial. *JPhysioter.* 2017. Apr;63(2):101-107
- Imran, H. M., Baig, M., Erqou, S., Taveira, T. H., Shah, N. R., Morrison, A., Choudhary, G., Wu, W. C. 2019. Home-Based Cardiac Rehabilitation Alone and Hybrid With Center-Based Cardiac Rehabilitation in Heart Failure: A Systematic Review and Meta-Analysis. *J Am Heart Assoc.* 8(16), e012779.
- Januzzi JL, van Kimmenade R, Lainchbury J, Bayes-Genis A, Ordonez- Llanos J, Santalo-Bel M, et al. 2006. NT-proBNP testing for diagnosis and short-term prognosis in acute destabilized heart failure: an international pooled analysis of 1256 patients: the international collaborative of NT-proBNP study. *Eur Heart J*;27:330–7 [L¹ SEP]
- Jolly, K., Taylor, R.S., Lip, G.Y. 2009. A randomized trial of the addition of home-based exercise to specialist heart failure nurse care: the Birmingham Rehabilitation Uptake Maximisation study for patients with Congestive Heart Failure (BRUM-CHF) study. *Eur J Heart Fail*;11:205–13
- Lam, C.,S.,P. 2015. Heart failure in Southeast Asia: facts and numbers. *ESC Heart Failure*; 2:46–49.
- Lam, C.,S.,P., 2019. The Role of NT-ProBNP in Clinical Evaluation of Heart Failure. *J Chin Med Assoc*; 82:447-451
- Levin, E., Gardner, D. and Samson, W. (1998) Natriuretic peptides. *N Engl J Med* 339: 321–328.
- Linke, A., Schoene, N., Gielen, S., Bs, C., Erbs, S., Schuler, G., Hambrecht, R. 2001. Endothelial Dysfunction in Patients With Chronic Heart Failure : Systemic Effects of Lower-Limb Exercise Training. *JAmCollCardiol.* 37. S0735-1097(00)01108-6
- Karamanoglu, M., Bennett, T., Ståhlberg, M. 2011. Estimation of cardiac output in patients with congestive heart failure by analysis of right ventricular pressure waveforms. *Biomed Eng Online*;10:36.
- Knapton, M. 2017. The BACPR Standards and Core Components for Cardiovascular Disease Prevention and Rehabilitation. *BritishCardiovascularSociety*.3rdedition
- Krupicka, J., Janota, T., Kasalova, Z. and Hradec, J. 2009 Natriuretic peptides: physiology, pathophysiology and clinical use in heart failure. *Physiol Res* 58: 171–177.
- Magri, P., Rao, M.A., Cangianiello, S., Bellizzi, V., Russo, R., Mele, A.F., Andreucci, M., Memoli, B., De Nicola, L. and Volpe, M. 1998 Early impairment of renal hemodynamic reserve in patients with asymptomatic heart failure is restored by angiotensin II antagonism. *Circulation* 98, 2849–2854
- Magnusson M., Melander O., Israelsson B., Grubb A., Groop L., Jovinge S. 2004.Elevated Plasma levels of NT proBNP in patients with diabetes without overt cardiovascular disease. *Diab care.* 1929-1935

- Mampuya, W.M., 2012. Cardiac rehabilitation past, present and future: an overview. *Cardiovasc.Diagn.Ther.* 2, 38–49.
- Marin-Grez, M., Fleming, J.T. and Steinhausen, M. 1986. Atrial natriuretic peptide causes pre-glomerular vasodilatation and post-glomerular vasoconstriction in rat kidney. *Nature* 324, 473–476 CrossRef PubMed ^[1]_{SEP}
- Masoudi F, 2012. Functional Capacity in Heart Failure: Which Test, and for What Purpose?. *J Am Coll Cardiol* Nov 21
- Masson S, Latini R, Anand IS, Barlera S, Angelici L, Vago T, et al.; Val- HeFT Investigators. 2008. Prognostic value of changes in N-terminal pro-brain natriuretic peptide in val-heft (valsartan heart failure trial). *J Am Coll Cardiol* ;52:997–1003. ^[1]_{SEP}
- McMurray, J.J., Adamopoulos, S., Anker, S.D., Auricchio, A., Bohm, M., Dickstein, K., Falk, V., Filippatos, G., Fonseca, C., Gomez-Sanchez, M.A. et al. 2012. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur. J. Heart Fail.* 14, 803–869
- Mesquita, ET., Jorge AJL., Rabelo LM., Souza Jr, CV., 2017. Understanding Hospitalization in Patients with heart failure. *IntJour Card Sci*, 30(1):81-90
- Mueller C., McDonald K., deBoer R., Maisel A., Cleland JGF., Kozhuharov N., 2019. Heart Failure Association of the European Society of Cardiology practical guidance of the use of natriuretic peptide concentration. *Eur Jour of Heart Fail.* 715-731
- Naughton J. Exercise training for patients with coronary artery disease. Cardiac rehabilitation revisited. *Sports Med* 1992;14:304-19. ^[1]_{SEP}
- Oremus M, Don-Wauchope A, McKelvie R, Santaguida PL, Hill S, Balion C *et al.* 2014. BNP and NT-ProBNP as prognostic markers in persons with chronic stable heart failure. *Heart Fail rev* DOI 10.1007/s10741-014-9439-6
- Ozkara A, Turgut F, Selcoki Y, Karanfil A, Metin MR, Kanbay M. 2007. Probrain natriuretic peptide for assessment of efficacy in heart failure treatment. *adv ther.* 24(6):1233-9
- Pearson M, King N, Smart N. 2018. Effect of exercise therapy on established and emerging circulating biomarker in patient with heart failure: a systematic review and meta-analysis. *Open hrt*: <http://dx.doi.org/10.1136/openhrt-2018-000819>
- Piepoli, M., Dickstein, K., Piepoli, M.F., Conraads, V., Corra, U., Francis, D.P., Jaarsma, T., McMurray, J., Pieske, B. 2014. Exercise training in heart failure : from theory to practice . A consensus document of the Heart Failure Association and the European Association for Cardiovascular Prevention and Rehabilitation.*eurjhf*. hfr017
- Piotrowicz, E., Baranowski1, R., Bilinska1, M., Stepnowska1, M., Piotrowska, M., Wo Jcik, A., Korewicki, J., Chojnowska, L., Malek, L.A., Klotowski, M., Piotrowski, W., Piotrowicz, W. 2010. A new model of home-based

- telemonitored cardiac rehabilitation in patients with heart failure: effectiveness, quality of life, and adherence. *European Journal of Heart Failure*. 12, 164–171 doi:10.1093/eurjhf/hfp181
- Piotrowicz, E., Piepoli M, Jaarsma T., Lambinau E, Coats A, Schmid JP *et al*, 2016. Telerehabilitation in heart failure patients: The Evidence and Pitfalls. *International Journal of Cardiology*,408-413
- Ponikowski, P., Voors, A.A., Anker, S.D., Bueno, H., Cleland, J.G.F., Coats, A.J.S., Falk, V. 2016. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *European Heart Journal*. Volume 37, Issue 27, pages 2129-2200
- Radi, B., Santoso, A., Siswanto, B., Mansyur, M., Ibrahim, N., Kusmana, D. 2017. Early Exercise Program for Patients with Heart Failure after Hospital Discharge. *International Journal of Physical Medicine & Rehabilitation*. 05. 10.4172/2329-9096.1000392
- Reddy, Y.N. V, Borlaug, B.A. 2016. Heart Failure With Preserved Ejection Fraction. *Curr. Probl. Cardiol*. 41, 145–188. *cpcardiol*.2015.12.002
- Reyes, E.B., Ha, J.W., Firdaus, I. 2016. Heart failure across Asia: same healthcare burden but differences in organization of care. *Int J Cardiol*; 223:163–167.
- Rivera M, Visconti RT, Salvador A, Bertomeu V, Miro V, Burgos F,. 2004. NT-ProBNP levels and hypertension. Their importance in the diagnosis of heart failure. *Rev esp De Card*. Vol5. pages 396-402
- Rohrbach, G., Schopfer, D.W., Krishnamurthi, N., Pabst, M., Bettencourt, M., Loomis, J., Whooley, M.A. 2017. The Design and Implementation of a Home-Based Cardiac Rehabilitation Program. *Fed. Pract*. 34, 34–39.
- Salah K, Stienen S., Pinto YM, Eurlings LW, Metra M, Bayes-Genis, A. 2019. Prognosis and NT-ProBNP in heart failure patients with preserved versus reduced ejection fraction. *Heart BMJ*. 105:15
- Santoso, A., Maulana, Ri., Alzahra, F., Prameswari, H., Ambari, A., Hartopo, A., Arso, I., Radi, B. 2020. The Effects of Aerobic Exercise on N-terminal Pro-B-type Natriuretic Peptide and Cardiopulmonary Function in Patients With Heart Failure: A Meta-Analysis of Randomised Clinical Trials. *Heart, Lung and Circulation*. 10.1016/j.hlc.2020.05.098
- Savarese, G., Lund, L.H. 2017. Global Public Health Burden of Heart Failure. *Card Fail Rev*;3(1):7-11.
- Savarese, G., Hage C., Orsini, N., Dahlstrom U., Perrone-Filardi P., Rosano GMC *et al*. 2016. Reduction in NT ProBNP levels are associated with lower mortality and heart failure hospitalization rates in patient with heart failure with mid-range and preserved ejection fraction. *Circ Heart Fail*.,Nov 9(11)
- Scalvini S., Grossetti F., Paganoni AM, Rovere MTL, Pedretti R., 2019. Impact of in-hospital cardiac rehabilitation on mortality and readmission in heart failure: A population study in Lombardy, Italy from 2005 to 2012>, *Eur Heart Journ of Prev Card*, Vol 26 (8) 808-817

- Siswanto, B.B., Sunanto, Munawar, M., Kusmana, D., Hanafiah, A., Waspadji, S. 2006. Predictor of mortality and rehospitalization of acute decompensated heart failure at six months follow up. *Crit Care & Shock.* ;9:61-7
- Smart NA, Meyer T, Butterfield JA, Faddy SC, Passino C, Malfatto M *et al*, 2011. Individual patient meta-analysis of exercise training effects on systemic brain natriuretic peptide expression in heart failure. *Eur Jour of Prev Card.* 19(3) 428-435
- Smart NA, Steele M, 2010. Systematic review of the effect of aerobic and resistant training on systemic brain natriuretic peptide (BNP and NT Pro BNP in heart failure patients. *Int Jour Card.* 260-265
- Soelistijo, S.A., Lindarto, D., Decroli, E., Permana, H., Sucipto, K.W., Kusnadi, Y., Budiman, Ikhsan, R., Sasiarini, L., Sanusi, H., 2019. Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 Dewasa di Indonesia. PB Perkeni
- Sokhanvar S, Shekhi M, Mazlomzadeh S, Golmohammadi Z. 2011. The Relationship between serum NT-ProBNP levels and prognosis in patients with Heart Failure. *Jour of Card and Thor Res.* 3(2): 57-61
- Steiner J, Guglin, M, 2008. BNP or NTpro BNP? A clinician's perspective. *Int Jour Card.* doi:10.1016/j.ijcard.2007.12.093
- Sullivan MJ, Higginbotham MB, Cobb FR. Exercise training in patients with severe left ventricular dysfunction. Hemodynamic and metabolic effects. *Circulation* 1988;78:506-15
- Sugiyono. 2007. Statistika untuk penelitian.. Penerbit Alfabeta Bandung.
- Squires RW, Gau GT, Miller TD. 1990. Cardiovascular rehabilitation: status, Mayo Clin Proc 1990;65:731-55 [SEP]
- Steinen S, Salah K, Dickhoff C, Carubelli V. 2015. N-Terminal Pro-B-Type Natriuretic Peptide (NT-proBNP) Measurements Until a 30% Reduction Is Attained During Acute Decompensated Heart Failure Admissions and Comparison With Discharge NT-proBNP Levels: Implications for In-Hospital Guidance of Treatment. *Jour of Card fail* 21(11)
- Taylor, R.S, Brown, A., Ebrahim, S., Jolliffe, J., Noorani, H., Rees K. 2004. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *American Journal of Medicine.* 15;116(10):682-92
- Taylor, R.S., Dalal, H., Jolly, K., Moxham, T., Zawada, A. 2014. Europe PMC Funders Group Home-based versus centre-based cardiac rehabilitation. *Cochrane Database Syst Rev* 1–64.14651858.CD007130
- Thibault, G., Amiri, F. and Garcia, R. (1999) Regulation of natriuretic peptide secretion by the heart. *Annu. Rev. Physiol.* 61, 193–217
- Thomas, R.J., Beatty, A.L., Beckie, T.M., Brewer, L.P.C., Brown, T.M., Forman, D.E., Franklin, B.A., Keteyian, S.J., Kitzman, D.W., Regensteiner, J.G., Sanderson, B.K., Whooley, M.A. 2019. Home-Based Cardiac Rehabilitation: A Scientific Statement From the American Association of Cardiovascular and Pulmonary Rehabilitation, the American Heart

- Association, and the American College of Cardiology. *J. Am. Coll. Cardiol.* 74, 133–153.
- Trimarco, B., Lembo, G., De Luca, N., Volpe, M., Ricciardelli, B., Condorelli, G., Rosiello, G. and Condorelli, M. (1989) Blunted sympathetic response to cardiopulmonary receptor unloading in hypertensive patients with left ventricular hypertrophy. A possible compensatory role of atrial natriuretic factor. *Circulation* 80, 883–892
- Tsutamoto, T. 2006. Relationship Between Renal Function and Plasma Brain Natriuretic Peptide in Patients With Heart Failure, *Journal of the American College of Cardiology*. Vol. 47, No. 3; p582-586
- Tucker, W.J., Lijauco, C.C., Nelson, M.D., Sarma, S., Mbbs, N. 2017. Mechanisms of the Improvement in Peak VO₂ with Exercise Training in Heart Failure with Reduced or Preserved Ejection Fraction. *Heart. Lung Circ. j.hlc.07.002*
- Van Tol BA, Huijisman RJ, Kroon DW, Schotthorst M, Kwakkel G, 2006. Effects of exercise training on cardiac performance, exercise capacity and quality of life in patients with heart failure: a meta-analysis. *Eur J Heart Fail.*8(8):841-50
- Verdiani V, Ognibene A, Rutili MS, Lombardo C. 2008. NT-ProBNP reduction percentage during hospital stay predicts long-term mortality and readmission in heart failure patients. *Journ of Card Med* 9(7):694-9
- Volterrani, M., Iellamo, F. 2016. Cardiac Rehabilitation in Patients with Heart Failure: New Perspectives in Exercise Training. *Card Fail Rev*;2(1):63-68.
- Wardani. 2019. 83 persen pengguna internet Indonesia pakai Whatsapp. <https://m.liputan6.com/tekno/read/4113678/83-persen-pengguna-internet-indonesia-pakai-whatsapp>. 18 Nov 2019, 17.00
- Wise, F.M. 2007. Exercise based cardiac rehabilitation in chronic heart failure. *Aust. Fam. Physician* 36, 1019–1024
- Zhong, B, 2009. How to calculate sample size in randomized controlled trial? *Journal of thoracic disease*, Vol 1 51-4