

DAFTAR PUSTAKA

- Adamson, E.A.S., Hurwood, D.A., & Mather, P.B., 2010. A reappraisal of the evolution of Asian snakehead fishes (Pisces, *Channidae*) using molecular data from multiple genes and fossil calibration. *Molecular Phylogenetics and Evolution*. 56(2): 707–717.
- Adamson, E.A.S., Hurwood, D.A., & Mather, P.B. 2012. Insights into historical drainage evolution based on the phylogeography of the chevron snakehead fish (*Channa striata* Bloch, 1793) in the Mekong Basin. *Freshwater Biology*. 57(11): 2211–2229.
- Ahmadi A., 2018. The length-weight relationship and condition factor of the threatened snakehead (*Channa striata* Bloch, 1793) from Sungai Batang River, Indonesia. *Polish Journal of Natural Sciences*. 33(4): 607-623.
- Ahmadi. 2019. Morphometric characteristic and growth patterns of Climbing perch (*Anabas testudineus*) from Sungai Batang River, Indonesia. *International Journal of Hydrology*. Vol. 3(4): 270-277.
- Aida S.N. 2018. Studi komparasi jenis makanan ikan gabus (*Channa striata*) di rawa banjir Lubuk Lampam, Sumatera Selatan. *Prosiding Seminar Nasional Ikan ke 8. Masyarakat Iktiologi Ikan*.
- Ali A.B. 1999. Aspects of the reproductive biology of female snakehead (*Channa striata* Bloch, 1793) obtained from irrigated rice agroecosystem, Malaysia. *Hydrobiologia*. Vol. 411: 71-77.
- Aliyu-Paiko, M. & Hashim, R. 2012. Effects of substituting dietary fish oil with crude palm oil and palm fatty acid distillate on growth, muscle fatty acid composition and the activities of hepatic lipogenic enzymes in snakehead (*Channa striatus*, Bloch 1793) fingerling. *Aquaculture Research*. 2(43): 767–776.
- Alvoidodinasyari R., Pribadi E.S., Soejoedono R.D. 2019. Kadar protein terlarut dalam albumin ikan gabus (*Channa striata* (Bloch, 1793) dan *Channa micropeltes*) asal Bogor. *Jurnal Veteriner*. Vol. 20(3):436-444.
- Ammar J.A., Kamal M.M., Sulistiono. 2018. Keragaman ikan di Dabau Cala, Kabupaten Musi Banyuasin, Sumatera Selatan Posiding Seminar Nasional Ikan ke-8 Masyarakat Iktiologi Indonesia. 195-199.
- Ambak, M.A., Bolong A-M. A., Ismail P. & MinhTam B. 2006. Genetic variation of snakehead fish (*Channa striata* Bloch, 1793) populations using Random Amplified Polymorphic DNA. *Journal of Biotechnology*. 5(1): 104–110.
- Anggoro S., RudiYanti S., & Rahmawati Y., 2013. Domestikasi ikan kerapu macan (*Epinephelus fuscoguttatus*) melalui optimalisasi media dan pakan. *Management of Aquatic Resources Journal*. 2(3): 119-127.
- Arikunto, S. 2010. *Prosedur penelitian suatu pendekatan praktik*. Jakarta: Rineka Cipta. pp. 413.
- Arechavala-Lopez P., Sanchez-Jerez P., Bayle-Sempere J.T., Sfakianakis D.G., & Somarakis S. 2011. Morphological differences between wild and farmed Mediterranean fish. *Hydrobiologia*. 679(1): 217-231.
- Artemov A.V., Mugue N.S., Rastorguev S.M., Zhenilo S., Mazur A.M., Tsygankova S.V., Boulygina E.U., Kaplun D., Nedoluzhko A.V., Medvedeva Y.A., & Prokhortchouk E.B. 2017. Genome-Wide DNA Methylation Profiling Reveals Epigenetic Adaptation of Stickleback to Marine and Freshwater Conditions. *Molecular Biology and Evolution*. 34(9): 2203-2213.
- Asiah N., Junianto, Yustiati A., dan Sukendi. 2018. Morfometrik dan meristik ikan kelabau (*Osteochilus melanopleurus*) dari Sungai Kampar, Provinsi Riau. *Jurnal Perikanan dan Kelautan*. Vol. 23(1):47-56.
- Asyari. 2007. Pentingnya labirin bagi ikan rawa. *Bawal*. 1(5): 161-167.
- Aydin S. 2015. A short history, principles, and types of ELISA, and our laboratory experience with peptide/protein analyses using ELISA. *Peptides*. 72: 4-15.

- Azrita & Syandri H. 2013. Fecundity, egg diameter and food *Channa lucius* cuvier in different waters. *Journal of Fisheries and Aquaculture*. Vol 4(3): 115-120.
- Azrita, Syandri H., Dahelmi, Syaifullah & Nugroho E. 2013. Karakterisasi morfologi ikan bujuk (*Channa lucius*) pada perairan Danau Singkarak Sumatera Barat, Rawa Banjiran Tanjung Jabung Timur Jambi dan Rawa Banjiran Kampar Riau. *Natur Indonesia*, 15(1), pp.1–8.
- Basak R., Roy A., & Rai U. 2016. Seasonality of reproduction in male spotted mirrel *Channa punctatus*: correlation of environmental variables and plasma sex steroid with histological changes in testis. *Fish Physiol. Ciochem*. Vol. 42:1249-1258.
- BPBAT Mandiangin. 2014. Naskah akademik ikan gabus haruan (*Channa striata* Bloch 1793) hasil domestikasi. Balai Perikanan Budidaya Air Tawar (BPBAT) Mandiangin Direktorat Jenderal Perikanan Budidaya Kementerian Kelautan dan Perikanan.
- Benziger A., Philip S., Raghavan R., Al P.H.A., Sukumaran M., Thairan J.C., Dahanukar N., Baby F., Peter R., Devi K.R., Radhakrishnan K.V., Haniffa M.A., Britz R., & Antunes A. 2011. Unraveling a 146 years old taxonomic puzzle: Validation of Malabar Snakehead, species-status and its relevance for Channid systematics and evolution. *PLoS ONE*. 6(6): 1–12.
- Bernatchez L. 2016. On the maintenance of genetic variation and adaptation to environmental change: considerations from population genomics in fishes. *Journal of Fish Biology*. 89(6): 1-38.
- Bhat, A.A., Haniffa M.A., Milton M.J., Paray B.A., Divya P.R. & Gopalakrishnan A. 2014. Genetic variation of striped snakehead (*Channa striatus* Bloch, 1793) populations using Random Amplified Polymorphic DNA (RAPD) markers. *International J. Biodiversity and conservation*. 6(5):363–372.
- Bich T.T.N, Tri D.Q., Yi-Ching C., Khoa H.D. 2020. Productivity and economic viability of snakehead *Channa striata* culture using an aquaponics approach. *Aquacultural Engineering*. 89.
- Biswas T.K., Bhattacharya T. K., Narayan A. D., Badola S., Kumar P., Sharma A. 2003. Growth hormone Gene Polymorphism and Its Effect on Birth Weight in Cattle and Buffalo. *Animal Breeding and Genetics*. 16(4): 494-497.
- Boonkusol D., Junshum P., & Panprommin K. 2020. Gonadosomatic index, oocyte development and fecundity of the snakehead fish (*Channa striata* Bloch, 1793) in natural river of Singburi Province, Thailand. *Pakistan Journal of Biological Sciences*. Vol. 23(10):1-8.
- BPBATM (Balai Perikanan Budidaya Air Tawar Mandiangin). 2014. Naskah akademik ikan gabus haruan (*Channa striata* Bloch 1793) hasil domestikasi. Kementerian Kelautan dan Perikanan Jakarta. pp: 67.
- Bruce S., Pedron S., Mehner T., Lauridsen T.L., Argillier C., Winfield I.J., Volta P., Emmrich M., Hesthagen T., Holmgren K., Benejam L., Kelly F., Krause T., Palm A., Rask M., & Jeppesen E. 2013. Fish diversity in European lakes: geographical factors dominate over anthropogenic pressures. *Freshwater Biology*. 58(9): 1779-1793.
- Burnawi & Pamungkas, Y.P., 2015. Komposisi jenis pakan alami ikan gabus (*Channa striata* Bloch, 1793) di Danau Cala, Kabupaten Musi Banyuasin Propinsi Sumatera Selatan. *Buletin Teknik Litkayasa Sumberdaya dan Penangkapan*. 13(2): 71–72.
- Calduch-Giner J.A., Mingarro M., de Celis S.V., Boujard D., & Perez-Sanchez J. 2003. Molecular cloning and characterization of gilthead sea bream (*Sparus aurata*) growth hormone receptor (GHR). Assessment of alternative splicing. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*. 136(1):1-13.
- Cardin S.X., Friedland K.D., & Waldman J.R. 2005. Stock identification Methods-Applications in Fishery Science. Chapter 1-Stock identification methods: An overview. Academic Press. Pp:3-6.

- Carvalho G.R. 1993. Evolutionary aspects of fish distribution: genetic variability and adaptation. *Journal of Fish Biology*. 43(5A): 53-73.
- Chasanah, E., M. Nurilmala, A.R. Purnamasari & D.Fihriani 2015. Komposisi kimia, kadar albumin dan bioaktivitas ekstrak protein ikan gabus (*Channa striata*) alam dan hasil budidaya. *Jurnal Pascapanen dan Bioteknologi Kelautan dan Perikanan*. 10(2):123-132.
- Cheng, F. Zhao, S. Schmidt, B.V. Ye, L. & Hallerman, E.M. 2017. Morphological but no genetic differentiation among fragmented populations of *Hemiculter leucisculus* (Actinopterygii, Cyprinidae) from a lake complex in the middle Yangtze, China. *Hydrobiologia* 809:185–200
- Cizidziel, J.V., Hinners T.A., Heithmar E.M., Pollard J.E., & Cross C.L. 2002. Mercury concentrations in fish from Lake Mead USA, related to fish size, condition, trophic level, location, and consumption risk. *Water, Air, and Soil Pollution*, 135, 355.
- Cui W., Liu N., Zhang X., Zhang Y., Qu L., Yan H., Lan X., Dong W., & Pan C. 2019. A 20-bp insertion/deletion (indel) polymorphism within the CDC25A gene and its associations with growth traits in goat. *Arch. Animal Breeding*. Vol. 62(1): 353-360.
- Datta S.N., Vannet I.K., Asha D. & Geeta J. 2013. Estimation of length-weight relationship and condition factor of spotted snakehead *Channa punctata* (Bloch) under different feeding regimens. *Springer Plus*. 2 (436): 2-5.
- De Faria C.U., Magnabosco C.U., Reyes A.L., Lobo R.B., Bezerra L.A.F., & Sainz R.B. 2007. of growth traits in Nelore cattle (*Bos indicus*). *Genetic and Molecular Biology*. 30(3): 753-760.
- De-Santis C., & Jerry D.R. 2007. Candidate growth gene in finfish-Where should we be looking?. *Aquaculture*. 272(1-4): 22-38.
- Devlin R.H., Sakhrani D., Tymchuk W.E., Rise M.L., & Goh B. 2009. Domestication and growth hormone transgenesis cause similar changes in gene expression in coho salmon (*Oncorhynchus kisutch*). *PNAS*. 106(9): 3047-3052.
- Dharmadi, Kartamihardja E.S., Utomo A.D., & Oktaviani D. 2009. Komposisi dan fluktuasi hasil tangkapan tugu di sungai Lempuing, Sumatera Selatan. *Jurnal Penelitian Perikanan Indonesia*. 15(20): 105-112.
- Djumanto, A. Murjiyanti, N. Azlina, A. Nurulitaerka, & A. Dwiramadhani. 2019. Reproductive biology of striped snakehead, *Channa striata* (Bloch, 1793) in Lake Rawa Pening, Central Java. *Jurnal Iktiologi Indonesia*. 19(3): 475-490.
- Djumanto, Setyobudi E., Simanjuntak C.P.H., & Rahardjo M.F. 2020. Estimating the spawning and growth of striped snakehead *Channa striata* Bloch, 1793 in Lake Rawa Pening Indonesia. *Nature*. 10: 19830.
- Diskominfo (Dinas Komunikasi dan Informasi). 2015. Peta Wilayah Kabupaten Kampar. Diakses dari <https://kominfosandi.kamparkab.go.id/peta-wilayah/#:~:text=Ketinggian%20tempat%2015%20%E2%80%93%20475%20meter%20di%20atas%20permukaan%20laut.> (29 Desember 2020).
- Dinesh K.R., Lim T.M., Chan W.K., & Phang V.P.E. 1996. Genetic variation inferred from RAPD fingerprinting in three species of tilapia. *Aquaculture International*. 4: 19-30.
- Dumalagan F.A., Garcines J.V., & Boyles L.Z. 2017. Reproductive biology, length-weight relationship, and condition factor of *Channa striata* (Bloch, 1793) tributaries of Lake Kilobidan, Agusan Marsh, Philippines. 2017. *International Journal of Computing, Communications, and Instrumentation Engineering*. Vol. 4(1): 78-81.
- Dunham, R.A. 2004. *Aquaculture and fisheries bio- technology: genetic approach*. CABI Publishing. Cambridge MA. 367 p.
- Duong T-Y., Nguyen T-T., & Pham T-L. 2017. Morphological differentiation among cultured and wild *Clarias macrocephalus*, *C. macrocephalus* x *C. gariepinus* hybrids, and their parental species in the Mekong delta, Viet Nam. *International Journal of Fisheries and Aquatic Studies*. Vol. 5(1): 233-240.



- Duong T.-Y., Sophorn U., Chheng P., So N., Tran T.-H.T., Nguyen N.-T.T., Pomeroy R., & Egna H. 2019. Genetic diversity and structure of striped snakehead (*Channa striata*) in the lower Mekong Basin: implications for aquaculture and fisheries management. *Fisheries Research*. Vol.218: 166-173.
- Dwiyedi A.K., & Dubey V.K. 2013. Advancements in morphometric differentiation: a review on stock identification among fish populations. *Review in Fish Biology and Fisheries*. 23 (4): 557-594.
- Ebbesson L.O.E., & Braithwaite V.A. 2012. Environmental effects on fish neural plasticity and cognition. 81(7): 2151-2174.
- Elmer, K.R., Kusche, H., Lehtonen, T.K., & Meyer, A. 2010. Local variation and parallel evolution: Morphological and genetic diversity across a species complex of neotropical crater lake cichlid fishes. *J. Biological Sciences*. 365: 1763-1782.
- Eprilurahman R., H.A. Asti., S. Hadisusanto, D.S. Yudha, Trijoko, R.S. Ramadani, F.X.S. Pranoto, I.A. Muhtianda. 2018. *Kekayaan fauna Gianyar, Bali*. Gadjah Mada University Press, Yogyakarta. 133 p.
- Excoffier, L. & Lischer H.E.L. 2010. Arlequin suite ver 3.5: A new series programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*. 10: 564-567
- Faizal I., Alilah R.S., Amarullah M.H., Megawati N. Sutanti, & Alimuddin. 2012. Produksi protein rekombinan hormon pertumbuhan ikan kerapu. *Jurnal Riset Akuakultur*. 7(2): 231-235.
- Fang M., Nie Q., Luo C., Zhang D., & Zhang X. 2006. An 8 bp indel in exon 1 of Ghrelin gene associated with chicken growth. *Domestic animal endocrinology*. 32: 216-225.
- Fang M., Nie Q., & Luo C. 2010. Associations of GHSR gene polymorphisms with chicken growth and carcass traits. *Molecular biology reproduction*. 37:423-428.
- Farid D.M. 2018. Hubungan Panjang Berat, Frekuensi Kejadian Makanan, dan Faktor Kondisi Ikan Gabus (*Channa striata*) di Waduk Gondang, Kabupaten Lamongan, Jawa Timur. Thesis, Universitas Brawijaya.
- Feliatra F., Yoswaty D., Lukystyowati I., & Hasyimi W. 2015. Karakteristik bakteri yang diisolasi dari saluran pencernaan udang galah (*Macrobrachium rosenbergii*, de Man) dan udang windu (*Panaes monodon* Fabricus) secara molekuler. *Aquacultura Indonesiana*. 15(1): 11–19.
- Feng. X., Yu X., & Tong J. 2014. Novel single nucleotide polymorphisms of the insulin-like growth factor-I gene and their associations with growth traits in common carp (*Cyprinus carpio* L.). *International journal of Molecular Science*. 15: 22471-22482.
- Ferdausi, H.J., N.C. Roy, M.J. Ferdous, M.A. Hossain, & M.M. Hasan. 2015. Reproductive biology of striped snakehead (*Channa striata* Bloch, 1793) from natural wetlands of Sylhet, Bangladesh. *Ann. Vet. Anim. Sci.*, 2: 162-169.
- Firlianty, E. Suprayitno, H. Nursyam, Hardoko & A. Mustafa. 2013. Chemical composition and acid profile of *Channidae* collected from Kalimantan, Indonesia. *International journal of science and technology*. 2(4): 25-31.
- Firmat, C., U.K. Schliwewn, M. Losseau & P. Alibert. 2012. Body shape differentiation at global and local geographic scale in the invasive cichlid *Oreochromis mossambicus*. *Biological J. of the Linnean Soc.* 105: 369-381.
- Fischer J., 2013. Fish identification tools for biodiversity and fisheries assessments: Review and guidance for decision-makers. *FAO Fisheries and Aquaculture Technical Paper*. Rome, Italia. pp. 107.
- Fitzpatrick S.W., Gerberich J. C., Kronenberger J. A., Angeloni L.M., & Funk W.C. 2014. Locally adapted traits maintained in the face of high gene flow. *Ecology Letters*. 18(1): 1-11.
- Froese, R. 2006. Cube law, condition factor and weight–length relationships: History, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22: 241–253.

- Geneaid, 2016. Geneaid DNA Isolation Kit. Instruction manual for research use. Geneaid Biotech Ltd.
- Gencheva D., & Stoyanova S. 2018. Polymorphisms of the candidate genes associated with growth traits in common carp (*Cyprinus carpio* L.). *Agricultural Sciences*. 10(23): 1-6.
- Ghaedi A., Kabir M.A., & Hashim R. 2013. Oocyte development and fecundity of snakehead murrel, *Channa striatus* (Bloch 1793) in captivity. *Asian Fisheries science*. Vol. 26: 39-51.
- Giacalone V.M., D'Anna G., Badalamenti F., & Pipitone C. 2010. Weight-length relationships and condition factor trends for thirty-eight fish species in trawled and untrawled areas off the coast of northern Sicily (central Mediterranean Sea). *Journal of Applied Ichthyology*. 26: 954-957).
- Gleni H.H., & Rudhy G. 2013. Peningkatan produktivitas budidaya ikan di lahan gambut. *J. Media Akuakultur*. 8 (1): 13-21.
- Graur D., & Li W-H. 2000. *Fundamentals of molecular evolution*. 2nd Edition. Sinauer Associates, Inc. Publisher. Pp. 479.
- Gusrina, 2014. *Genetika dan reproduksi ikan*, Yogyakarta: Deepublish. 263 p.
- Gustiano R, Oktaviani T., Soelistyowati D.T., Kusmini I.I., Wahyutomo, & Huwoyon G.H. 2013. Analisis ragam genotip RAPD dan fenotip truss morfometrik pada tiga populasi ikan gabus *Channa striata* (Bloch, 1793) (Bloch , 1793). *J. Ilmu-Ilmu Hayati*. 12(3): 325–333.
- Habibie S.A., Djumanto, & Murwantoko. 2018. Polychromatic sexual dimorphism and redescription species of red devil *Amphilophus Amarillo* [Stauffer & McKaye, 2002] in Sermo Reservoir, Yogyakarta. *Jurnal Iktiologi Indonesia*. 18(1): 69-86.
- Haniffa MA, Nagarajan M, Gopalakrishnan A. 2006. Length–weight relationships of *Channa punctata* (Bloch, 1793) from Western Ghats Rivers of Tamil Nadu. *Journal of Applied Ichthyology*. 22(4):308-309.
- Hartl D.L., & Clark G.C. 1997, *Principles of Population Genetics*. Sinauer Associates, Sunderland. Sinauer Associates Inc. pp. 682.
- Hien, T.T.T., Thi Be T., Lee C.M., & Bengston D.A. 2015. Development of formulated diets for snakehead (*Channa striata* (Bloch, 1793) and *Channa micropeltes*): Can phytase and taurine supplementation increase use of soybean meal to replace fish meal?. *Aquaculture*. 448: 334–340.
- Hien, T.T.T., Phu M.T., Cam Tu T.L., Tien N.V., & Minh Duc P. 2016. Effect of replacing fish meal with soya protein concentrate on growth, feed efficiency and digestibility in diets for snakehead. *Aquaculture research*. 1(8): 1–8.
- Hien T.T.T., Tam B.M., Tu T.L.C., and Bengtson D.A. 2017. Weaning methods using formulated feed for snakehead (*Channa striata* and *Channa micropeltes*) larvae. *Aquaculture research*: 1-9.
- Holden, M.J. & D.F.S., Raitt. 1974. *Manual of fisheries science. Part 2. Methodes of Resource Investigation and Their Application*. FAO Fish. Technical Paper, 214 p.
- Hossain, Azmad M., Sohel M., Mariya A., Fazley R.A., Sultana S., Arifur R.Md., Mahub M., Jakiul I.Md., Hossain H.M.M., and Mosarof M. 2015. Ovarian biology of spotted snakehead (*Channa punctatus*) from natural wetlands of Sylhet, Bangladesh. *Annals of Veterinary and Animal Science*. Vol. 2(3): 64-76.
- Hu, X., Li, C., & Shi, L. 2012. A novel 79-bp insertion/deletion polymorphism in 3'-flanking region of IGF-I gene is associated with growth-related traits in common carp (*Cyprinus carpio* L.). *Aquaculture Research*. 44(10): 1632–1638.
- Hua K., Koppe W., & Fotanilas R. 2019. Effects of dietary protein and lipid levels on growth, body composition and nutrient utilization of *Channa striata* (Bloch, 1793). *Aquaculture*. Vol. 501: 368-373.
- Huang C-Y & Lin H-C. 2010. The effect of acidity on gill variations in the aquatic air-breathing fish, *Trichogaster lalius*. *Comparative Biochemistry and Physiology, Part A*. 158: 61-71.

- Ibanez-Agurre A.L., Cabral-Solis E., Gallardo-Cabello M., & Espino-Barr E. 2006. Comparative morphometric of two populations of Mugil Curema (Pisces: Mugilidae) on the Atlantic and Mexican Pacific coasts. *Scientia Marina*. 70(1): 139-145.
- Imbert, E. & Lefevre. 2003. Dispersal and gene flow of *Populus nigra* (Salicaceae) along a dynamic river syst. *J. of Ecol.* 91: 447-456.
- Irawan B. 2018. *Genetika: Penjelasan mekanisme pewarisan sifat*. Airlangga University Press. Surabaya. pp: 312.
- Irmawati. 2016. *Genetika Populasi Ikan*. Andi Offset. Yogyakarta. Pp: 244.
- Irmawati, Tresnati J., Nadiarti and Fachruddin. 2019. Sex Differentiation and Gonadal Development of striped snakehead (*Channa striata* Bloch, 1793) Bloch, 1793). IOP Conf. Sereis: Earth and Environmental Science. IOP Publishing. 253: 1-8.
- Iqbal M. 2011. Ikan-ikan di hutan rawa gambut Merang-Kepayang dan Sekitarnya. Merang REDD Project (MRPP), Palembang, Sumatera Selatan Indonesia. Pp: 92.
- Jamaluddin, J.A.F., Pau, T.M. & Siti-Azizah, M.N. 2011. Genetic structure of the snakehead murrel, *Channa striata* (Bloch, 1793) (*Channidae*) based on the cytochrome c oxidase subunit i gene: Influence of historical and geomorphological factors. *Genetics and Molecular Biology*. 34(1):152–160.
- Jaya-Ram A., Ishak S.D., Enyu Y-L., Kuah M-K., Wong K-L., & Shu-Chien A.C. 2011. Molecular cloning and ontogenic mRNA expression of fatty acid desaturase in the carnivorous striped snakehead fish (*Channa striata*). *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 158(4): 415-422.
- Jing Ni, Feng You, Jianhe Xu, Dongdong Xu, Aiyun W., Zhihao Wu, Yongli Xu, and Penijun Z. 2012. Single nucleotide polymorphisms in intron 1 and intron 2 of Larimichtys crocea growth hormone gene are correlated with growth traits. *Chinese Journal of Oceanology and Limnology*. 3(2): 279-285.
- Jaya-Ram, A., Ishak, S. D., Enyu, Y. L., Kuah, M. K., Wong, K. L., & Shu-Chien, A. C. (2011). Molecular cloning and ontogenic mRNA expression of fatty acid desaturase in the carnivorous striped snakehead fish (*Channa striata*). *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 158(4), 415–422.
- Jisr N., Younes G., Sukhn C., & El-Dakdouki M.H. 2018. Length-weight relationship and relative condition factor of fish inhabiting the marine area of the Eastern Mediterranean city, Tripoli-Lebanon. *Egyptian Journal of Aquatic Research*. Vol. 44: 299-305.
- John S. L., & Southgate P.C. 2007. *Aquaculture farming aquatic animals and plants*. 2nd Edition. Blackwell Publishing. 648 p.
- Jonsson B., N. Jonsson, & A.G. Finstad. 2012. Effects of temperature and food quality on age and size at maturity in ectotherms: an experimental test with Atlantic salmon. *Journal of Animal Ecology*. 82(1): 201-210.
- Jorgensen, H.B.H., Pertoldi C., Hansen M.M., Ruzzante D.E., & Loeschcke V. 2008. Genetic and environmental correlates of morphological variation in a marine fish the case of Baltic Sea herring (*Clupea harengus*). *Canadian Journal of Fisheries and Aquatic Sciences*, 65, 389-400.
- Jubaedah^a D., Kamal M.M., Muchsin I., & Hariyadi S., 2015. Karakteristik kualitas air dan estimasi resiko ekobiologi herbisida di perairan rawa banjiran lubuk lampam, Sumatera Selatan. *J. Manusia dan Lingkungan*. 22(1): 12-21.
- Jubaedah^b D., S. Hariyadi, I. Muchsin, & M.M. Kamal. 2015. Water quality index of floodplain River Lubuk Lampam South Sumatera Indonesia. *International Journal of Environmental Science and Development*. 6(4): 252-258.
- Kaban S., Asyari, Fatah K., Marini M., Nanda T., Burnawi, Nasution D.H., & Mersi. 2015. Laporan teknis: Kajian tingkat degradasi potensi sumberdaya ikan di Sungai Batanghari, Jambi. Balai Penelitian Perikanan Perairan Umum Palembang, Kementerian Kelautan dan Perikanan. 73 p.



UNIVERSITAS
GADJAH MADA

- Kakkaeo, M., Chittapalapong, T. & Villanueva, M., 2004. Food habits, daily ration and relative food consumption in some fish populations in Ubolratana Reservoir, Thailand. *Asian Fish Sci.* 17: 249–259.
- Kalita M, Choudhury H, Saikia A, Sarma D. Length–weight relationships of four endemic snakeheads (*Channa andrao* Britz, 2013, *Channa aurantimaculata* Musikasinthorn, 2000, *Channa bleheri* Vierke, 1991, and *Channa stewartii* (Playfair, 1876)] from the Brahmaputra drainage, northeast India. *J Appl Ichthyol.* 00:1–2
- Kartamihardja E.S. 2015. Fish Stock Enhancement and Restocking of the Inland Waters of Indonesia: Lessons Learned. *Fish People.* 13(3): 25-32.
- Kashyap A., Awasthi M., Arshad M., & Serajuddin M. 2015. Length-weight, length-length relationship and condition factor of freshwater murrel, *Channa punctatus* from Northern and Eastern Region of India. *Journal of Fish and Marine Sciences.* Vol. 7(3):164-170.
- Kashyap, A., Awasthi, M., & Serajuddin, M. 2016. Phenotypic Variation in Freshwater Murrel, *Channa punctatus* (Bloch , 1793) from Northern and Eastern Regions of India Using Truss Analysis. *International Journal of Zoology*, 2016, 1–6.
- Katharine, D. Gisele, P. & Pinha, D. 2017. Dispersal mode and flooding regime as drivers of benthic metacommunity structure in a Neotropical floodplain. *Hydrobiologia* 788:131–141.
- Kelley J.L., P.M. Davies, S.P. Collin, & P.F. Grierson. 2017. Morphological plasticity in a native freshwater fish from semiarid Australia in response to variable water flows. *Ecology and Evolution.* 7(16): 6595-6605.
- Kementerian Kelautan dan Perikanan. 2011. Statistik perikanan tangkap di Indonesia tahun 2010. Direktorat Jenderal Perikanan Tangkap. 134 p.
- Kementerian Kelautan dan Perikanan. 2018: Produktivitas Perikanan Indonesia. Diakses pada 20 Desember 2020 <https://kkp.go.id/wp-content/uploads/2018/01/KKP-Dirjen-PDSPKP-FMB-Kominfo-19-Januari-2018.pdf>
- Khan M.A., Khan S., & Miyan K. 2012. Length-weight relationship of giant snakehead, *Channa marulius* and stinging catfish, *Heteroneustes fossilis* from the River Ganga, India. *Journal of Applied Ichthyology.* Vol. 28(154-155).
- Khumari S.A., & Kumar N.S.R. 2015. Length-weight relationship and condition factor in *Channa punctatus* (Bloch) from Hussaingsagar Lake, Hyderabad, Andhra Pradesh, India. *Aquatic Ecosyste: Biodiversity, Ecology and Conservation.* Springer India: 283-286.
- Kikko T., Usuki T., Ishizaki D., Kai Y., & Fujioka Y. 2014. Relationship of egg and hatchling size to incubation temperature in the multiple-spawning fish *Gnathopogon caeruleus* (Honmoroko). *Environmental Biology of Fishes.* 98: 1.151-1.161.
- Kimura M. 1980. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16:111-120.
- King R.S. 2015. Cluster analysis and data analysis data mining: An introduction. David Pallai Publisher. USA. pp: 300.
- Keivany Y. & Zamani-Faradonbe M. 2017. Length-weight and length-length relationships for seven fish species from the Zohreh River, Iran. *Journal of Applied Ichthyology.* Vol. 33(3): 1-3.
- Kocour M., & Kohlmann. 2011. Growth hormone gene polymorphisms in tench, *Tinca tinca* L. *Aquaculture.* 310: 298-304.
- Kocour M., & Kohlmann K. 2014. Distribution of five growth hormone gene haplogroups in wild and cultured tench, *Tinca tinca* L., populations. *Applied ichthyology.* 30(1):22-28.
- Konan, K.M. Adepo-Gourene, A.B. Outtara, A. Nyingy, W.D. & Gourene, G. 2010. Morphometric variation among male populations of freshwater shrimp *Macrobrachium vollehovenii* Herklots, 1851 from Côte d'Ivoire Rivers. *Fisheries Research* 103:1-8.
- Koundal A., Dhanze R., & Sharma I. 2014. Length-weight relationship, condition factor, and

- relative growth patterns of *Channa punctata* (Bloch) from Himachal Pradesh, India. Zoo's Print. Vol. 29(1): 25-29.
- Kumar R., Mohanty U.L., & Pillai B.R. 2021. Effect of hormonal stimulation on captive broodstock maturation, induced breeding and spawning performance of striped snakehead, *Channa striata* (Bloch, 1793). *Animal Reproduction Science*. 224
- Kusmini, I.I., Prakoso, V.A. & Kusdiarti, 2015. Keragaman fenotipe truss morfometrik dan genotipe ikan gabus (*Channa striata* Bloch, 1793) dari Jawa Barat, Sumatera Selatan dan Kalimantan Tengah. *J. Riset Akuakultur*. 10(4): 501–510.
- Laily N. 2006. Identifikasi jenis-jenis ikan teleostei yang tertangkap nelayan di wilayah perairan pesisir kota Semarang. Skripsi Prodi Biologi. Fakultas MIPA, Universitas Negeri Semarang. 44 p.
- Lakra, W.S., Goswami M., Gopalakrishnan A., Singh D.P., Singh NS., & Nagpure N.S. 2010. Genetic relatedness among fish species of genus *Channa* using mitochondrial DNA genes. *Biochemical Systematics and Ecology*. 38(6): 1212–1219.
- Lei M., Luo C., Peng X., Fang M., Nie Q., Zhang D., Yang G., & Zhang X. 2007. Polymorphism of Growth-Related Genes Associated with Fatness and Muscle Fiber Traits in Chickens. *Poultry science*. 86:835-842.
- Leprieur, F. Tedesco, P.A. Hugueny, B. Beauchard, O. Durr, H.H. Brosse, S. & Oberdorff T. 2011. Partitioning global patterns of freshwater fish betas diversity reveals contrasting signatures of past climate changes. *Ecol. Letters* 14:325-334.
- Li, K., B. Shieh, Yuh-wen C., Da-ji H., & Shih-hsiung L. 2016. Growth, diet composition and reproductive biology of the invasive freshwater fish chevron snakehead *Channa striata* (Bloch, 1793) on a subtropical island. *Zoological Studies*. 55(53): 1-11.
- Listiyanto, N. & Andriyanto, S., 2009. Ikan gabus (*Channa striata* Bloch, 1793) manfaat pengembangan dan alternatif teknik budidayanya. *J. Pascapanen dan Bioteknologi Kelautan dan Perikanan*. 4(1): 18–25.
- Liu LS, Yu XM & Tong JG. 2012. Molecular characterization of myostatin (MSTN) gene and association analysis with growth traits in the bighead carp (*Aristichthys nobilis*). *Mol. Biol. Rep*. 39: 9211-9221.
- Liu X., Liang H., Liang Y., Zhong L., Qin X., Zhang T., Zhang J., Zou G., & Hu G. 2017. Significant associations of polymorphisms in the growth hormone gene with growth traits in common carp (*Cyprinus carpio*). *Aquaculture*: 14 (38-41).
- Marcus L.F., & Corti M. 1996. Overview of the New, or Geometric Morphometrics. *Advances in Morphometrics*. pp: 1-13.
- Mahmud N.A., Rahman H.Md.H., Mostakim G.M., Khan M.G.Q., Shahjahan Md., Lucky N.S., and Islam M.S. 2016. Cyclic variations of gonad development of an air-breathing fish, *Channa striata* (Bloch, 1793) in the lentic and lotic environments. *Fisheries and Aquatic Sciences*. Vol. 19(5): 1-7.
- Maftuchah, & Winaya, A.Z., 2014. Teknik Dasar Analisis Biologi Molekuler.
- Makmur S, Rahardjo M.F., & Sutrisno Sukimin. 2003. Reproductive biology of snakehead fish, *Channa striata* (Bloch, 1793) in Flood Plain Area of Musi River, South Sumatera. *Jurnal Iktiologi Indonesia*. Vol.3(2):57-62
- Makmur S. 2006. Fecundities and eggs diameter of snakehead (*Channa striata* (Bloch, 1793) Bloch) in the flood plain area of Musi River, South Sumatera. *Jurnal Perikanan*. Vol. VIII(2): 254-259.
- Marimuthu K., Kumar D., & Haniffa A. 2007. Induced Spawning of Striped Snakehead, *Channa striatus*, Using Ovate. *J. Applied Aquaculture*. 19 (4): 95-103.
- Matthews W.J. & Marsh-Matthews E. 2017. Stream fish community dynamics: A critical synthesis. Johns Hopkins University Press, Maryland, USA. 345 p.
- Martinez-Gonzalez C.C., Gonzalez-Daza W., and Mojica J.I. 2018. Length-weight relationships of fishes in the Mira basin, Colombia. *Journal of Applied Ichthyology*: 1-4.

- Maruska K.P. & R.D. Fernald. 2010. Behavioral and physiological plasticity: Rapid changes during social ascent in an African cichlid fish. *Journal of Hormones and Behavior*. 58(2): 230-240.
- Matos L.S., Parisotto D.C., & Carvalho L.N., 2017. Length-weight relationship condition factor of the Characidae matrinsa, *Brycon falcatus* (Muller & Troschel, 1844) in the Teles Pires River, southern Amazon. *Journal of Applied Ichthyology*: 1-5.
- Mestanza-Ramon C., Henkanaththege S.M., Duchicela P.V., Tierras Y.V., Capa M.S., Mejia D.C., Gutierrez M.J., Guaman M.C., & Ramon P.M. 2020. In-Situ and Ex-Situ Biodiversity Conservation in Ecuador: A Review of Policies, Actions and Challenges. *Diversity*. 12: 1-18.
- Meilina M.F. 2019. Hubungan Panjang dengan berat serta faktor kondisi ikan gabus (*Channa striata* Bloch) asal dua perairan di wilayah Kabupaten Musi Banyuasin dan sumbangannya pada pembelajaran biologi. Skripsi, Universitas Sriwijaya.
- Michel M.J., H. Chien, C.E. Beachum, M.G Bennett, & J.H. Knouft. 2016. Climate change, hydrology, and fish morphology: predictions using phenotype-environment associations. *Climate Change*. 140: 563-576.
- Milton J., Bhat A.A., Haniffa M.A., Hussain S.A., Rather I.A., Al-Anazi K.M., Hailan W.A.Q., and Farah M.A. 2017. Ovarian development and histological observations of threatened dwarf snakehead fish, *Channa gachua* (Hamilton, 1822). *Saudi Journal of Biological Sciences*. Vol. 25: 149-153.
- Miranda, L.E. 2011. Depth as an organizer of fish assemblages in floodplain lakes. *Aquat. Sci*. 73:211–221.
- Mitu N.R., Alam M.M., Hussain M.A., Hasan M.R., & Singha C. 2019. Length-weight and length-length relationships, sex ratio and condition factors of the Asian striped dwarf catfish *Mytus tengara* (Hamilton, 1822) (Siluriformes: Bagridae) in the Ganges River, Northwestern Bangladesh. *Iran J. Ichtiol*. Vol. 6(1): 21-30.
- Miyan, K. Afzal, M. Kumar, D. & Khan, S. 2016. Truss morphometry and otolith microchemistry reveal stock discrimination in *Clarias batrachus* (Linnaeus, 1758) inhabiting the Gangetic river system, *Fisheries Research* 173: 294–302.
- Moder, K. Schlick-Steiner, B.C., Steiner, F.M. Cremer, S. Christian, E. & Seifert, B. 2006. Optimal species distinction by discriminant analysis: comparing established methods of character selection with a combination procedure using ant morphometrics as a case study. *J. of Zoological Syst. and Evolutionary Research* 45:82-87.
- Mohaddasi, M. Shabanipour, N. & Abdolmaleki, S. 2013. Morphometric variation among four populations of Shemaya (*Alburnus chalcoides*) in the South of Caspian sea using truss network. *The J. of Basic and Applied Zool*. 66:87–92.
- Morrongiello J.R., Bond N.R., Crook D.A., & Wong B.B.M. 2012. Spatial variation in egg size and egg number reflects trade-offs and bet-hedging in a freshwater fish. *Journal of Animal Ecology*. 81(4): 806-817.
- Mulyasari, Soelistyowati D.T., Kristanto A.H., & Kusmini I.I. 2010. Karakteristik geneik enam populasi ikan nilem (*Osteochilus hasselti*) di Jawa Barat. *Jurnal Riset Akuakultur*. 5(2): 175-182.
- Muntazania M.P.A., Amin S.M.N., Rahman M.A., Rahim A.A., & Marimuthu K. 2013. Asian *Journal of Animal and Veterinary Advances*. Vol. 8(2): 369-375
- Munir M.B., Hashim R., Chai Y.H., Marsh Y.H., Marsh T.L., & Mohd Nor S.A. 2016. Dietary prebiotics and probiotics influence growth performance, nutrient digestibility and the expression of immune regulatory genes in snakehead (*Channa striata* Bloch, 1793) fingerlings. *Aquaculture*. 460: 59–68.
- Muntaziana M.P.A., Amin S.M.N., Rahman M.A., Rahim A.A., & Marimuthu K. 2013. Present culture status of endangered snakehead. *Asian J. Animal and Veterinary Advances*. 8(2): 369–375.

- Muslim M., Sasanti A.D., & Apriana A. 2019. The Effect of Immersion Duration in Thyroxine Hormone on Growth of Snakehead Fish Larvae (*Channa striata*). *Journal of Aquaculture Science*. 4(1): 1-11.
- Mutmainah D. 2013. Hubungan panjang berat dan faktor kondisi ikan gabus (*Channa striata* (Bloch, 1793) Bloch, 1793) yang dibesarkan di rawa lebak, Provinsi Sumatera Selatan. *J. Depik*. 2(3): 184-190.
- Muthukumar S., M. Arunachalam, U. Ramesh, M. Umamaheswari, & A. Vanarajan. 2017. A new record of the dwarf snakehead, *Channa ornatipinnis* Britz 2007 (Perciformes: Channidae) from India. *International J. Aquatic Biology*. 5(1): 29-32.
- Mulyadi G., Sasanti A.D., & Yulisman. 2016. The Rearing of Snakehead (*Channa striata*) with Different Stocking Density in Biofloc Media. *Jurnal Akuakultur Rawa Indonesia*. 4(2): 154-174.
- Nagarajan, M. Haniffa, M.A. Gopalakrishnan, A. Basheer, V.S. & Muneer, A. 2006. Genetic variability of *Channa punctatus* population using Randomly Amplified Polymorphic DNA. *Aqualt. Research* 27:1151-11155.
- Nakkrasae L., K. Wisetdee, & N. Charoenphandhu. 2015. Osmoregulatory adaptations of freshwater air-breathing snakehead fish (*Channa striata*) after exposure to brackish water. *J. Comp. Physiol. B*. 185: 527-537.
- Ndobe S., Serdiati N., & Moore A. 2014. Domestication and length-weight relationship of striped snakehead *Channa striata* (Bloch, 1793) (Bloch). *Proceeding of International Conference of Aquaculture Indonesia (ICAI)*. 165-172.
- Neff B.D., Garner S.R., & Pitcher T.E., 2011. Conservation and enhancement of wild fish populations- preserving genetic quality versus genetic diversity. *Canada Journal of Fish Aquaculture Science*. 68: 1.139-1154.
- Nguyen, T. & Duong, L. 2016. Morphological and Genetic Differences Between Cultured and Wild Populations of *Channa striata* (Bloch, 1793) in Vietnam and its Phylogenetic Relationship with Other *Channa* Species. *Songklanakarin, J. Science Technology*. 38 (4): 427-434.
- Norainy. 2007. Morphological and genetic variability of Malaysian *Channa* spp. on morphometric and RAPD technique. Thesis. Universiti Sains Malaysia.
- Nontji A. 2016. Danau-danau alami Nusantara. *Oceanografi LIPI*. 294 p.
- Ouyang J.H., Xie L., Nie Q., Luo C., Liang Y., Zeng H., & Zhang X. 2014. Single nucleotide polymorphism (SNP) at the GHR gene and its associations with chicken growth and fat deposition traits. *British Poultry Science*. 49(2): 87-95.
- Paaby A.B., Blacket M.J., Hoffmann A.A., & Schmidt. 2010. Identification of a candidate adaptive polymorphism for *Drosophila* life history by parallel independent clines on two continents. *Molecular ecology*. 19:760-774.
- Pal A., Chakravarty A.K., Hattacharya, T.K., Joshi, B.K., Sharma A. 2004. Detection of Polymorphism of Growth hormone Gene for the Analysis of Relationship between Allele Type and Growth Traits in Karan Fries Cattle. *Asian-Australian Journal of Animal Sciences*. 17(10): 1334-1337.
- Pankova M.V., Kukhlevsky A.D., & Brykov V.A. 2017. Fish growth hormone genes: Divergence of coding sequence in salmonid fishes. *Russian Journal of Genetics*. 53(2): 221-232.
- Paez D.J. & J.J. Dodson. 2017. Environment-specific heritabilities and maternal effects for body size, morphology and survival in juvenile Atlantic salmon (*Salmo salar*): evidence from a field experiment. *Environmental Biology of Fishes*. 100: 209-221.
- Pasaribu A.F., Muslim M., & Syaifudin M. 2019. The Effect of Dipping Time in Thyroxine on Growth Rate and Survival Rate of Snakehead Larvae (*Channa striata*). *Jurnal Akuakultur Rawa Indonesia*. 7(1): 25-33.
- Pease, A.A. Gonzalez-diaz, A.A., Rodiles-hernandez, R. & Winemiller, K.O. 2012. *Funct. diversity and trait-enviro. relationships of stream fish assemblages in a large tropical*

- catchment. J. of Freshwater Biol. 64:367-379.
- Pervaiz K., Iqbal Z., Mirza M.R., Haved M.N., & Naeem M. 2012. Meristic and Morphometric Studies on Indus Mahseer *Tor macrolepis* (Teleostei: Cyprinidae) from District Attock, Pakistan. International Journal of Agriculture & Biology. 14(2): 169-175.
- Pham M.D., Nam S., Hien T.T.T., & Robert P. 2011. Sustainable snakehead aquaculture development in the Lower Mekong River Basin of Cambodia and Vietnam: Part 2: Striped Snakehead fish diseases and water quality analysis. Aqua fish CRSP Project USAID Grant. Inland Fisheries Research and Development Institute, Phnom Penh, Cambodia, Can Tho University, Vietnam and University of Connecticut, USA.
- Pinkert C.A. 2014. Transgenic animal technology. Elsevier. pp: 696.
- Pope K.L., Lochmann S.E., & Young M.K. 2010. Methods for assessing fish populations. inland fisheries management in North America, 3rd edition. America Fisheries Society, Bethesda, Maryland, 325-351 p.
- Prakoso V.A., Ath-thar F.M.H., Radona D., & Kusmini I.I. 2018. Respon pertumbuhan benih ikan gabus (*Channa striata*) dalam kondisi pemeliharaan bersalinitas. Limnotek. 25(1): 10-17.
- Prasad L., Dwivedi A.K., Dubey V.K., & Serrajudin M. 2011. Reproductive biology of freshwater murrel, *Channa punctatus* (Bloch, 1793) from river Varuna (a tributary of Ganga River) in India. J. Echophysiol. Occup. Hlth. Vol. 11: 69-80.
- Purnamawati, Djokosetiyanto D., Nirmala K., Harris E., Affandi R., 2017 Survival and growth of striped snakehead fish (*Channa striata* (Bloch, 1793) Bloch.) juvenile reared in acid sulfate water and rainwater medium. AACL Bioflux 10(2):265-273.
- Purnamawati, Nurmala, Shilman M.I., Dewantoro E., Utami A.S. 2018. Effects of ameliorant on the growth of snakehead fish (*Channa striata* Bloch, 1793) juvenile reared in acid sulfate water medium located in tidal land. AACL Bioflux. Vol. 11(6): 1919-1926.
- Purnamasari R. & D.R. Santi. 2017. Fisiologi Hewan. Penerbit Prodi Arsitektur UIN Sunan Ampel, Surabaya, Jawa Timur. 113 p.
- Rahim, M.H., Ismail P., Alias R., Muhammad N., Mat Jais A.M. 2012. PCR-RFLP analysis of mitochondrial DNA cytochrome b gene among Haruan (*Channa striatus*) in Malaysia. Gene. 494(1): 1-10.
- Rahman M.A., Arshad A., Amin S.M.N., & Shamsudin M.N. 2013. Growth and survival of fingerlings of a threatened snakehead of *Channa striates* (Bloch) in earthen nursery ponds. Asian Journal of animal and veterinary advances. Vol. 8(2): 216-226.
- Reid J.E. & Chaput G. 2012. Spawning history influence on fecundity, egg size, and egg survival of Atlantic salmon (*Salmo salar*) from the Miramichi River, New Brunswick, Canada. ICES Journal of Marine Science. 69(9): 1678-1685.
- Resfiza, Muslim dan Sasanti, A.D., 2014. Perbedaan jumlah kromosom ikan toman (*Channa micropeltes*) dengan Ikan Serandang (*Channa pleurophthalmus*). J. Akuakultur Rawa Indonesia. 2(2): 125-134.
- Requieron E.A., Torres R.M.A.J., & Demayo C.G. 2012. Applications of relative warp analysis in describing of scale shape morphology between sexes of the snakehead fish *Channa striata* (Bloch, 1793). International J. Biological, Ecology and Environmental Sciences. 1 (6): 205-209.
- Rinku G, Behera S, Bibha CB, Sonmoina B. 2013. Sexual dimorphism and gonadal development of a rare murrel species *Channa bleheri* (Bleher) in Assam. The Bioscan. 2013;8(4):1265-9.
- Rohlf F.J., & Marcus L.F. 1993. A revolution morphometrics. Trends in Ecology & Evolution. 8(4): 129-132.
- Rodriguez J.M., Angon E., Gonzalez M.A., 2017. Allometric relationship and growth models of juveniles of *Cichlasoma festae* (Perciforme: Cichlidae), a freshwater species native in Ecuador. Revista de Biología Tropical. 65(3): 1185-1193.



- Roy N. C., Chowdhury K., & Das S.K. 2016. Observation of hapa breeding technique of striped snakehead, *Channa striatus* (Bloch, 1793) under captive conditions. International J. Fisheries and Aquatic Studies. 4(5): 413-417.
- Samidjan I. & Rachmawati. 2016. Technology Engineering of Aquaculture Snakeheads *Channa striatus* (Bloch, 1793) using Cross Breeding from Different Waters for Determining the Genetic Variation of Superior Seeds. International Symposium on Aquatic Products Processing and Health. Aquatic Procedia. Vol. 7:136-145.
- Saputra F.M. 2009. Daerah Aliran Sungai Batanghari. Diakses dari https://staff.blog.ui.ac.id/tarsoen.waryono/files/2009/12/punya_tile.pdf (29 Desember 2009).
- Saputra A., Ath-thar M.H.F., Samsudin R., Putri F.P., & Prakoso V.A. 2017. Reproductive biology of striped snakehead (*Channa striata* (Bloch, 1793) Bloch, 1973) in Bogor and Bekasi, West Java. 16(3): 309-314
- Saputra A., Budiardi T., Samsudin R., & Rahmadya N.D. 2018. Growth performance and survival of snakehead *Channa striata* juvenile with different stocking density reared in recirculation system. Jurnal Akuakultur Indonesia. 17(2): 104-112.
- Sarkar UK, Khan GE, Dabas A, Pathak AK, Mir JI, Rebello SC *et al.* 2013. Length weight relationship and condition factor of selected freshwater fish species found in River Ganga, Gomti and Rapti, India. Journal of Environmental Biology. 34(5):951.
- Schunter C., Carreras-Carbonell J., Macpherson E., Tintore J., Vidal-Vijande E., Pascual A., Guidetti P., & Pascual M. 2011. Matching genetics with oceanography: Directional gene flow in a Mediterranean fish species. Molecular Ecology. 20(24): 5167–5181.
- Sebrina, N. & Rahajoe, P.S. 2012. Perbedaan Antara Pemberian Asupan Putih Telur Ayam Kampung dan Ekstrak Ikan Gabus Terhadap Pembentukan Kolagen pada Penyembuhan Luka Gingiva Labial (Kajian in Vivo pada Tikus Wistar). Skripsi. Universitas Gadjah Mada.
- Setyawati A. 2009. DAS Sungai Siak Provinsi Riau. Diakses dari https://staff.blog.ui.ac.id/tarsoen.waryono/files/2009/12/das_siak_nuranitasetyawati_0706265705.pdf (29 Desember 2020).
- Senguttuvan M. & A.A. Sivakumar. 2016. Studies on the maturation and spawning of *Channa striata* (Bloch, 1793) in Ukkadam Lake, Coimbatore, Talimandu, India. Journal of Aquaculture in the Tropics. 31(1): 13-23.
- Sharpe D.M.T., Rasanen K., Berner D., & Hendry A.P. 2008. Genetic and environmental contributions to the morphology of lake and stream stickleback: implications for gene flow and reproductive isolation. J. Evolutionary Ecology Research. 10 (6): 849-866.
- Shen H.M., Chen X.R., Chen W.Y., Lin S.M., Chen Y.J., Zhang L., and Luo L. 2016. Influence of dietary phosphorus levels on growth, body composition, metabolic response and antioxidant capacity of juvenile snakehead (*Channa argus* x *Channa maculata*). Aquaculture nutrition. 1-9.
- Shrestha T.R. 2009. Growth hormone gene manipulation in *Channa striatus* Bloch. Disertation. Center for Cellular and Molecular Biologi, India. 122 p.
- Shuai F., S.Yu, S. Lek, & X. Li. 2018. Habitat effects on intra-species variation in functional morphology: Evidence from freshwater fish. Ecology and Evolution. 8(22): 10902-10913.
- Simon K.D., Y.Bakar, S.E. Temple & A.G. Mazlan. 2010. Morphometric and meristic variation in two congeneric archer fishes *Toxotes chatareus* (Hamilton 1822) and *Toxotes jaculatrix* (Pallas 1767) inhabiting Malaysian coastal waters. Journal of Zhejiang University Science B. 11:871-879.
- Singh C.P., Ram R.N., & Singh R.N. 2012. Food and feeding pattern of *Channa punctatus* in two different habitats at Tarai region of Uttarakhand. Journal of Environmental Biology. Vol. 34: 789-792.
- Singh M. & Serajuddin M. 2017. Length-weight, length-length relationship and condition factor of *Channa punctatus* collected from three different rivers of India. Journal of Entomology and Zoology Studies. Vol. 5(1):191-197.

- Song L.M., Munian K., Rashid Z.A., & Bhassu S. 2013. Characterisation of Asian snakehead murrel *Channa striata* (Bloch, 1793) (*Channidae*) in Malaysia: An insight into molecular data and morphological approach. *The Scientific World J.* 1–16.
- Sfakianakis, D.G. & Somarakis, S. 2012. Morphological differences between wild and farmed Mediterranean fish, *J. Hydrobiologia* 679: 217–231.
- Strauss R.E & F.L. Bookstein. 1982. The truss: Body from reconstruction in morphometrics. *System Zoology*. 31(2): 113-135.
- Strychalski J., Peierzchala M., Pareek C.S. 2011. Relationship between the insertion/deletion polymorphism within the promoter and the intron 1 sequence of the PRNP gene and milk performance traits in cattle. *Czech Journal of Animal Science*. 56(4): 151-156.
- Strüssmann, C.A. & Nakamura, M. 2002. Morph., endocrinology and enviro. modulation of gonadal sex differentiation in teleost fishes. *Fish Physiology and Biochemistry* 26:13–29.
- Suardika I.P., Edi D.G.S., Arya I.W., & Darmadi N.M. 2019. Growth pattern and maturity level of Chevron Snakehead's gonad (*Channa striata* Bloch, 1793). Series. 4th Annual Applied Science and Engineering Conference. *Journal of Physiscs: Conference*. IOP Publishing. 1402(2019)033061: 1-7
- Subagdja, Mutmainah D., Sawestri S., Atminarso D., Makri & Sudrajat A. 2013. Laporan teknis: Ekologi, biologi dan kapasitas penangkapan sumberdaya ikan di Danau Ranau, Provinsi Sumatera Selatan. Balai Penelitian Perikanan Perairan Umum Palembang, Kementerian Kelautan dan Perikanan. 77 p.
- Sugiyono. 2012. Metode penelitian kuantitatif, kualitatif dan R&D. Alfabeta, Bandung. pp. 380.
- Supiwong, W., Jearanaiprepame, P. & Tanomtong, A., 2009. A new report of karyotype in the chevron snakehead. 74(3): 317–322.
- Syah F., Yustina, & Suwondo. 2020. Keanekaragaman ikan Kabupaten Kampar. Penerbit Lakeisha, Klaten, Jawa Tengah. 150 p.
- Syafei L.S. 2017. Keanekaragaman hayati dan konservasi ikan air tawar. *Jurnal Penyuluhan Kelautan dan Perikanan Indonesia*. 11(1): 48-62.
- Tan, M.P. Jamsari A.F.J., Muchisin Z.A., & Siti Azizah M.N. 2015. Mitochondrial genetic variation and population structure of the striped snakehead, *Channa striata* (Bloch, 1793) in Malaysia and Sumatra, Indonesia. *Biochemical Systematics and Ecology*. 60: 99–105.
- Tan, M.P., Jamsari, A.F.J. & Siti Azizah, M.N., 2012. Phylogeographic pattern of the striped snakehead, *Channa striata* (Bloch, 1793) in Sundaland: Ancient river connectivity, geographical and anthropogenic singnatures. *PLoS ONE*. 7(12): 1–11.
- Tanamati F., Silva S.C.C., Rodriguez M.D.P.R., Schuroff G.P., Nascimento C.S., Vesco A.P.D., & Gasparino E. 2015. GHR and IGF-I gene expression and production characteristics associated with GH gene polymorphism in Nile tilapia. *Aquaculture*. 435: 195-199.
- Tatangindatu F., Kalesaran O., dan Rompas R. 2013. Studi parameter fisika dan kimia air pada areal budidaya ikan di Danau Tondano, Desa Paleloan, Kabupaten Minahasa. *Jurnal Budidaya Perairan* Vol. 1(2): 8-19.
- Thanh N.M., Barnes A.C., Mather P.B., Li Y., and Lyons R.E. 2010. Single nucleotide polymorphisms in the actin and crustacean hyperglycemic hormone genes and their correlation with individual growth performance freshwater prawn *Macrobrachium rosenbergii*. *Aquaculture*. 301:7-15.
- Thermo Scientific. 2012. Product Information Thermo Scientific Verso 1-Setp RT-PCR Hot-Start Kit.
- Topal, M. Yaganogle, A.M. Sonmez, A.Y. Arslan, G. & Hisar, O. 2010. Using discriminant and CHAID analysis methods to identify sex in brown trout (*Salmo trutta fario*) by morphometric features. *The Israeli J. of Aquaculture-Bamidgeh* 62:251-259.
- Turan C. 1999. A note on the examination of morphometric differentiation among fish populations: The truss system. *Turki J. Zoology*. 23: 259-263.

- Tzeng T-D. 2004. Morphological variation between populaitons of spotted mackerel (*Scomber australasicus*) off Taiwan. J. Fisheries Research. 68: 45-55.
- Utomo A.D., & Krismono. 2006. Aspek biologi beberapa jenis ikan langka di Sungai Musi, Sumatera Selatan. Prosiding Seminar Nasional Ikan IV. Jatiluhur, 29-30 Agustus 2006. Pp. 309-330.
- Utomo A.D., Wibowo A., & Mohtar H. 2017. Pengelolaan sumberdaya perikanan di Kabupaten Ogan Komering Ulu (OKU) Selatan, Provinsi Sumatera Selatan. Laporan Badan Perencanaan Pembangunan Daerah Kabupaten Ogan n Komering Ulu Selatan dengan Balai Riset Perikanan Perairan Umum dan Penyulihan Perikanan Palembang. Pp. 139.
- Villeger, S. Miranda, J.R. Hernandez, D.F. & Mouillot, D. 2010. Contrasting changes in taxonomic vs. funct. diversity of tropical fish communities after habitat degradation. Ecol. of Freshwater Fish 28:41-52.
- Voris H.K.2000. Maps of *Pleisocene* sea levels in Southeast Asia: shorelines, river system and time durations. J. Biogeography. 27 :1153-1167.
- Wahyuni S., Windarti, & Putra R.W. Comparative study on histological structure of gill and kidney of snakehead fish (*Channa striata*, Bloch 1793) from the Kulim and Sibam Rivers, Riau Province. Jurnal Online Mahasiswa Fakultas Perikanan dan Ilmu Kelautan. 4(2): 1-14.
- Wang XL, Meng XY, Song B, Qiu XM, *et al.* (2010). SNPs in the myostatin gene of the mollusk *Chlamys farreri*: Association with growth traits. Comp. Biochem. Physiol. B 155: 327-330.
- Wang Y., Yang L., Zhou K., Zhang Y., Song Z., & He S. 2015. Evidence for adaptation to the Tibetan Plateau inferred from Tibetan Loach transcriptomes. Genom Biology Evolution. 7(11): 2970-2982.
- Wang X., Fu B., Yu X., Qu C., Zhang Q., and Tong J. 2018. Fine mapping of growth-related quantitative trait loci in Yellow river carp (*Cyprinus carpio* Haematoperus). Aquaculture. 484: 277-285.
- Wardoyo S.A. 2006. Pengaruh kerusakan penutupan vegetasi rawa gambut terhadap komunitas ikan di Sungai Merang Kabupaten Musi Banyuasin, Sumatera Selatan. Tesis. Prodi Pengelolaan Sumberdaya Alam dan Lingkungan. Institut Pertanian Bogor.
- Wikiandy N. 2013. Dampak penemaran limbah industri tekstil terhadap kerusakan struktur organ ikan yang hidup di DAS (Daerah Aliran Sungai) Citarum bagian hulu. Skripsi. Fakultas Perikanan dan Ilmu Kelautan, Universitas Padjajaran, Bandung.
- Yousefian, Mehdi, Shirzad, & Elham. 2011. The review of effect of growth hormone on immune system, metabolism, and osmoregulation of fish. Australian Journal of Basic and Applied Sciences. 5(5): 467-475.
- Zhao H., Ma H.J., Gao S.N., Chen X.R., Chen Y.J., & Lin S.M. 2016. Evaluation of dietary vitamin E supplementation on growth performance and antioxidant status in hybrid snakehead (*Channa argus* x *Channa maculate*). Aquaculture Nutrition. 2017:1-8.
- Zhang J., Li L., Gao N., Wang D., Gao Q., & Jiang S., 2010. Feature extraction and selection from volatile compounds for analytical classification of Chinese red wines from different varieties. Analytica Chimica Acta. 662(2): 137-142.