

DAFTAR PUSTAKA

- Arunangshu, G., Pradip, T., Nabarun, B., Bipan, T., Nagen, G., Rajib, B. 2012. Estimation of Theaflavin Content in Black Tea Using Eelectronic Tongue. *Journal of Food Engineering* 110 : 71–79
- Astika, W., D. Muchtar, dan Sutrisno. 1996. Klon-klon teh baru yang telah dilepas oleh Balai Penelitian Teh dan Kina, Gambung. *Warta Teh dan Kina*. 7: 6-15.
- Baldermann, S., Fleischmann, P., Bolten, M., Watanabe, N., Winterhalter, P., & Ito, Y. (2009). Centrifugal precipitation chromatography , a powerful technique for the isolation of active enzymes from tea leaves (*Camellia sinensis*), 1216, 4263–4267.
- Bhattacharyya, N.; Bandyopadhyay, R.; Bhuyan, M.; Tudu, B.; Ghosh, D.; Jana, A. Electronic nose for black tea classification and correlation of measurements with “tea taster” marks. *IEEE Trans. Instrum. Meas.* 2008, 57, 1313–1321.
- Biswas, A.K., and Sarkar, A.R. 1971. Biological and chemical factors affecting the valuation of North East Indian plains teas. II. Statistical evaluation of biochemical constituents and their effects on briskness, quality and cash valuations of black teas. *J. Sci. Food Agric.* 22, 196–204.
- Bonnely, S., Davis, A. L., Lewis, J. R., & Astill, C. 2003. A model oxidation system to study oxidised phenolic compounds present in black tea. *Food Chemistry* 83, 483e492.
- Borse, B.B., and Rao, L.J.M. 2012. Novel Bio-Chemical of Indian Black Teas with Reference to Quality Parameter. *Journal of Bioequivalence and Bioavailability* S14
- Buzun, G.A., Dzhemukhadze, K.M. and Milesenko, L.F. 1984. O-Diphenol oxidase of tea and quality of the raw material. *Subtrop Ku’lt* 1, 35–38.
- Cabrera, C., Gimenez, R., Lopez, C., 2003. Determination of tea components with antioxidant activity. *J. Agric Food Chem.* 51, 4427e4435
- Cairns, J. R. K. & Esen, A. 2010. Review:β-glucosidases. *Cellular anda Molecular Life Science.* 67:3389–3405
- Carlioni, P., Tiano, C., Padella, C., Bacchetti Z., Customu, C., Alexander, K., Damian, I., 2018. Antioxidant activity of white, green and black tea obtained from the sametea cultivar. *Food Research International* 53: 900–908
- Carlin, S., Masuero, D., Guella, G., Vrhovsek, U., Mattivi, F. 2019. Methyl Salicylate Glycosides in Some Italian Varietal Wines. *Molecules* 24, 3260
- Celik, A., Dincer, A., & Aydemir, T. 2016. Characterization of β-glucosidase immobilized on chitosan-multiwalled carbon nanotubes (MWCNTS) and their application on tea extracts for aroma enhancement. *International Journal of Biological Macromolecules* 89: 406-414

- Chaturvedula, V. S. P., & Prakash, I. 2011. The aroma, taste, color and bioactive constituents of tea. *Journal of Medicinal Plants Research*, 5, 2110–2124.
- Cleverdon, R., Elhalaby, Y., McAlpine, M., Gittings, W., and Ward, W. 2018. Total Polyphenol Content and Antioxidant Capacity of Tea Bags: Comparison of Black, Green, Red Rooibos, Chamomile and Peppermint over Different Steep Times. *Beverages* 4(15): 1–13.
- Cooper, R. 2011. Green tea and theanine: health benefits. *Int. J. Food Sci. Nutr.* 63 : 90–97
- Cui, J. L., Katsuno, T., Totsuka, K., Ohnishi, T., Takemoto, H., Mase, N., ... Tong, H. R. 2016. Characteristic fluctuations in glycosidically bound volatiles during tea processing and identification of their unstable derivatives. *Journal of Agricultural and Food Chemistry*, 64(5), 1151–1157.
- Daenen, L., Sterckx, F., Delvaux, F. R., Verachtert, H., Derdelinckx, G. 2008. Evaluation of the glycoside hydrolase activity of a *Brettanomyces* strain on glycosides from sour cherry (*Prunus cerasus* L.) used in the production of special fruit beers. *Federation of European Microbiological Societies*, 1103–1114.
- Deb, S., Jolvis, K.R. 2016. A review of withering in the processing of black tea. *Journal of Biosystems*. 41(4): 365-372
- DeGarmo, E.P., W.G. Sullivan, dan C.R Candra. 1984. *Engineering Economy*. 7th edition Mc Millan Publ. Co. New York.
- Du, L., Jianxun L., Wei L., Yunfei L., Tao L., Dongguang, X. 2014. Characterization of volatile compounds of pu-erh tea using solid-phase microextraction and simultaneous distillation–extraction coupled with gas chromatography–mass spectrometry : *Food Research International* 57;61–70
- FAO. 2017. Intergovernmental Group On Tea Intersessional Meeting : Report of the Working Group on Global Tea Market Analysis and Promotion. IGG:TE ISM 17/3
- Finger, A.; Kuhr, S.; Engelhardt, U. H. Chromatography of tea constituents. *J. Chromatogr.* **1992**, 624, 293-315.
- Gebely, T., 2015. Chemical Compounds in Tea. Retrieved 2nd December, 2016, from: <http://www.worldoftea.org/tea-chemistry/>.
- Gui, J., Fu, X., Zhou, Y., Katsuno, T., Mei, X., Deng, R., Yang, Z. 2015. Does Enzymatic Hydrolysis of Glycosidically Bound Volatile Compounds Really Contribute to the Formation of Volatile Compounds during the Oolong Tea Manufacturing Process? *Journal of Agricultural and Food Chemistry*, 63(31), 6905–6914.
- Haibo Pan, Ying Gao, Youying Tu, 2016. Mechanisms of body weight reduction by black tea polyphenols. *Molecules* 21, 1659.
- Han, Z. X., Rana, M. M., Liu, G. F., Gao, M. J., Li, D. X., Wu, F. G., ... Wei, S. 2016. *Green Tea Flavour Determinants And Their Changes Over Manufacturing Processes*. *Food Chemistry*, 212 : 739–748
- Haslam, E. *Practical Polyphenolics, From Structure to Molecular Recognition and Physiological Action*; Cambridge University Press: New York, 1998; pp 335-373.

- Ho, C.-T., Zheng, X., & Li, S. 2015. Tea aroma formation. *Food Science and Human Wellness*, 4(1), 9–27.
- Ichsan, O. A. N. 2019. Seleksi Klon Teh Unggulan PT Pagilaran sebagai Bahan Baku Teh Hitam Kualitas Premium. [Tesis]. Yogyakarta: Fakultas Teknologi Pertanian Universitas Gadjah Mada.
- Kalidass, S., Vijaya, K.U., and Kumar, R.R. 2019. Biochemical Changes During Withering Process of CTC black Tea Manufacture. *Oriental Journal of Chemistry*, 35, (4): 1313-1319
- Kang, S., Yan, H., Zhu, Y., Liu, X., Lv, H. P., Zhang, Y., ... Lin, Z. 2019. Identification and quantification of key odorants in the world's four most famous black teas. *Food Research International*, 121(January), 73–83.
- Kato, M., Shibamoto, T. 2001 Variation of major volatile constituents in various green teas from southeast Asia, *J. Agric. Food Chem.* 49 : 1394–1396.
- Kawakami, M., Kobayashi, A., 2000. Carotenoid-derived aroma compound in tea. *Abstr. Pap. Amer. Chem. Soc.* 219 U32.
- Kinoshita, T., Hirata, S., Yang, Z. Y., Baldermann, S., Kitayama, E., Matsumoto, S., et al. 2010. Formation of damascenone derived from glycosidically bound precursors in green tea infusions. *Food Chemistry*, 123(3), 601–606.
- Kraujalyte, V., Pelvan, V., Alasalvar C., 2016. Volatile compounds and sensory characteristics of various instant teas produced from black tea. *Food Chemistry* 194, 864–872
- Kumazawa, K., & Masuda, H. 2001. Change in the flavor of black tea drink during heat processing. *Journal of Agricultural and Food Chemistry*, 49(7), 3304–3309.
- Li, Y., Shibahara, A., Matsuo, Y., Tanaka, T., dan Kuono, I. 2010. Reaction of The Black Tea Pigment Theaflavin during Enzymatic Oxidation of Tea Catechins. *Journal of Natural Product* 73(1):33-39.
- McDowell, I.; Taylor, S.; Gay, C. The phenolic pigment composition of black tea liquors. Part I. Predicting quality. *J. Sci. Food Agric.* 1995, 69, 467–474.
- Nanninga, A. W. 1916. *De tehecultuur in Ned*, Indie, Amsterdam.
- Ni, H., Jiang, Q., Lin, Q., Ma, Q., Wang, L., Weng, S., Huang, G., Li, L., Chen, F. 2020. Enzymatic hydrolysis and auto-isomerization during β -glucosidase treatment improve the aroma of instant white tea infusion. *Food Chemistry*.
- Mitrowihardjo, S., Mangoendidjojo, W., Hartiko, H., dan Yudono, P. 2012. *Kandungan Katekin dan Kualitas (Warna Air Seduhan, Flavor, Kenampakan) Enam Klon Teh (Camellia sinensis (L.) O. Kuntze) Di Ketinggian Yang Berbeda*. *Agritech*, 32 (2) : 199–206.
- Mizukami, Y., Sawai, Y., & Yamaguchi, Y. (2008). Changes in the concentrations of acrylamide, selected odorants, and catechins caused by roasting of green tea. *Journal of Agricultural and Food Chemistry*, 56(6), 2154–2159.
- Ogawa, K., Moon, J. H., Guo, W., Yagi, A., Watanabe, N., & Sakata, K. 1995. A Study on Tea Aroma Formation Mechanism: Alcoholic Aroma Precursor Amounts and Glycosidase Activity in Parts of the Tea Plant. *Zeitschrift*

- Fur Naturforschung - Section C Journal of Biosciences*, 50(7–8), 493–498.
- Owuor, P. O., & Obanda, M. (2007). The use of green tea (*Camellia sinensis*) leaf flavan-3-ol composition in predicting plain black tea quality. *Food Chemistry*, 100 (3):873 -884
- Price, W. E., and Spitzer, J. C.. 1993. Variations In The Amounts of Individual Flavanols In A Range of Green Teas. *Food Chem.*, 47 (3) : 271–276.
- Puangpronpitag, D., Niamsa, N., & Sittiwet, C. 2009. Anti-Microbial Properties of Clove (*Eugenia caryophyllum* Bullock and Harrison) Aqueous Extract Against Food-Borne Pathogen Bacteria. *International Journal of Pharmacology*
- Qi, D., Miao, A., Cao, J., Wang, W., Chen, W., Pang, S., ... Ma, C. (2018). Study on the effects of rapid aging technology on the aroma quality of white tea using GC-MS combined with chemometrics: In comparison with natural aged and fresh white tea. *Food Chemistry*, 265, 189–199.
- Ravichandran, R. and Partibhan, R. 1998. Changes in enzyme activities (polyphenol oxidase and phenylalanine ammonia lyase) with type of tea leaf and during black tea manufacture and the effect of enzyme supplementation of dhoool on black tea quality. *Food Chem.* 62, 277–281.
- Robertson, A., *Phytochemistry* 1983, 22, 889–896.
- Robertson, A., Wilson, K. C., Clifford, M. N., Eds.; Chapman and Hall *The chemistry and biochemistry of black tea production: the non-volatiles. In Tea: Cultivation to Consumption: London, U.K., 1992; pp 553–601.*
- Schuh, C., & Schieberle, P. (2006). Characterization of the key aroma compounds in beverage prepared from Darjeeling black tea: Quantitative differences between tea leaves and infusion. *Journal of Agricultural and Food Chemistry*, 54(3), 916–924.
- Shahidi, F.; Naczki, M. *Phenolics in Foods and Nutraceuticals*; CRC Press: Boca Raton, FL, 2004
- Santoso, U. 2016. *Antioksidan Pangan*. Gadjah Mada University Press. Yogyakarta
- Senthil Kumar, R. S., Murugesan, S., Kottur, G., & Gyamfi, D. 2013. Black Tea: The Plants, Processing/Manufacturing and Production. *Tea in Health and Disease Prevention*, 41–57.
- Shukla, Y. 2007. Tea and Cancer Chemoprevention : A comprehensive Riview. *Asian Pasific Journal of Cancer Preventiom*, Vol. 8, Hal: 155-166
- Su, E., Xia, T., Gao, L., Dai, Q., & Zhang, Z. 2010. Immobilization of β -glucosidase and its aroma-increasing effect on tea beverage. *Food and Bioproducts Processing*, 88(2–3), 83–89.
- Sudaryat, Y., Kusmiyati, M., Pelangi, C.R., Rustamsyah, A., and Rohdiana, D. 2015. Aktivitas Antioksidan Seduhan Sepuluh Jenis Mutu Teh Hitam (*Camelia sinensis* (L.) O. Kuntze) Indonesia. *Jurnal Penelitian Teh dan Kina* 18 (2),95-100.
- Takeo, T. and Baker, J.E. 1973. Changes in multiple forms of polyphenol oxidase during maturation of tea leaves. *Physicochemistry* 12, 21–24.

- Takeo, T. 1992. *Green and semi-fermented teas*, In: *Tea: Cultivation to Consumption*, Willson, K. C. and Clifford, M. N. (eds.), London: Chapman and Hall, pp. 413–457
- Tampubolon, TW., 2011, Pengaruh Temperatur Terhadap Kadar Tanin Pada Pembuatan Teh Botol Sosro di PT Sinar Sosro KPB Deli Serdang, Skripsi. Progam Studi Analis Farmasi dan Makanan USU.
- Teng, J., Gong, Z., Deng, Y., Chen, L., Li, Q., Shao, Y., Lin, L., Xiao, W. Purification, Characterization and Enzymatic Synthesis of Theaflavins of Polyphenol Oxidase Isozyme from Tea Leaf (*Camellia sinensis*). *LWT-Food Science and Technology*, 84 (2017) 263-270.
- Thanaraj, S.N.S. and Seshadri, R. 1990. Influence of polyphenol oxidase activity and polyphenol content of tea shoot on quality of black tea. *J. Sci. Food Agric.* 51, 57–69.
- Toda, M., Mase, N., Matsuo, T., Tong, H., Yang, Z., Takemoto, H., ... Cui, J. 2016. Characteristic Fluctuations in Glycosidically Bound Volatiles during Tea Processing and Identification of Their Unstable Derivatives. *Journal of Agricultural and Food Chemistry*, 64(5), 1151–1157.
- Ullah, M. R. 1986. A rapid procedure for estimating theaflavins and thearubigins of black tea. *Two and a Bud*, 33, 46–48
- Ullah, M.R. and Roy, P.C. 1982. Effect of withering on the polyphenol oxidase level in the tea leaf. *J. Sci. Food Agric.* 33, 492–495.
- Venkatesan, S. and Sujitha, S. D. A. 2007. Influence of Genotype and Manufacturing Process on the Activity of β -D-Glucosidase and β -Galactosidase in Tea. *American Journal of Plant Physiology*, 2 (2): 175-182
- Wang, D., Yoshimura, T., Kubota, K., & Kobayashi, A. 2000. Analysis of glycosidically bound aroma precursors in tea leaves. 1. Qualitative and quantitative analyses of glycosides with aglycons as aroma compounds. *Journal of Agricultural and Food Chemistry*, 48(11), 5411–5418.
- Wang, D., Kubota, K., Kobayashi, A., & Juan, I. M. 2001. Analysis of glycosidically bound aroma precursors in tea leaves. 3. Change in the glycoside content of tea leaves during the oolong tea manufacturing process. *Journal of Agricultural and Food Chemistry*, 49(11), 5391–5396.
- Wanga, W., Zhanga, S., Lishuang L, Sanga S. (2018). A new method to prepare and redefine black tea thearubigins. *Journal of Chromatography A*, 1563 : 82–88
- Widjoseno, G. 2012. Potensi hasil dan Toleransi Curah Hujan Beberapa Klon Teh (*Camellia sinensis* (L.) O. Kuntze) PGL di Bagian Kebun Kayulandak PT Pagilaran. Fakultas Pertanian. Universitas Gadjah Mada. Skripsi.
- Winterhalter, P., and G. K. Skouroumounis. 1997. Glycoconjugated aroma compounds: occurrence, role and biotechnological transformation. In *Biotechnology of aroma compounds*, ed. R. G. Berger, 73–105. New York: Springer.
- Yamanishi, T., Kobayashi, A. 1999. Progress of tea aroma chemistry, in: R. Teranishi, E.L. Wick, I. Hornstein (Eds.), *Flavor Chemistry: 30 Years of Progress*, Springer, New York, NY, pp. 135–145.

- Yang, Z., Baldermann, S., & Watanabe, N. 2013. Recent studies of the volatile compounds in tea. *Food Research International*, 53(2), 585–599
- Yang, X., Wan, X., Li, Y., Wang, Y., Feng, Z., Zhang, L., & Li, M. 2019. Tea aroma formation from six model manufacturing processes. *Food Chemistry*, 285(October 2018), 347–354.
- Yuliana, R.A; Indradewa, Didik; Ambarwati, Erlina. 2013. *Potensi Hasil dan Tanggapan Sembilan Klon Teh (Camellia sinensis (L.) O. Kuntze) PGL Terhadap Variasi Curah Hujan di Kebun Bagian Pagilaran*. *Vegetatika* Vol.2 No.3, 2013 : 54 – 67.
- Zamora, A.J., Andrade, C.D., and Henares, J.A.R. 2016. Antioxidant capacity, total phenols and color profile during the storage of selected plants used for infusion. *Food Chemistry* 199 : 339–346
- Zhang, L. Z., Ni, H., Zhu, Y. B., Yang, Y. F., Li, L. J., Jiang, Z. D., ... Chen, F. 2018. Characterization of aromas of instant oolong tea and its counterparts treated with two crude enzymes from *Aspergillus niger*. *Journal of Food Processing & Preservation*, 42(2), e13500.
- Zhang, T., Fang, K., Ni, H., Li, T., Li, L. J., Li, Q. B., Chen, F. 2020. Aroma enhancement of instant green tea infusion using β -glucosidase and β -xylosidase. *Food Chemistry*, 315 (2020) 126287.
- Zhen, Y.-S., 2002. *Tea: Bioactivity and Therapeutic Potential*. CRC Press, USA.
- Zheng, X. Q., Li, Q. S., Xiang, L. P., Liang, Y. R. 2016. Review: Recent Advances in Volatile of Teas. *Molecules*, 338.
- Zhu, J. C., Chen, F., Wang, L. Y., Niu, Y. W., Yu, D., Shu, C., ... Xiao, Z. B. 2015. Comparison of aroma-active volatiles in oolong tea infusions using GC-olfactometry, GC-FPD, and GC-MS. *Journal of Agricultural and Food Chemistry*, 63(34), 7499–7510.
- Zeng, L., Watanabe, N., & Yang, Z. 2018. Understanding the biosyntheses and stress response mechanisms of aroma compounds in tea (*Camellia sinensis*) to safely and effectively improve tea aroma . *Critical Reviews in Food Science and Nutrition*, 8398, 1–14.