

DAFTAR PUSTAKA

- Abasi, S., Minaei, S., Jamshidi, B., Fathi, D., & Khoshtaghaza, M. H. (2019). Rapid measurement of apple quality parameters using wavelet de-noising transform with Vis/NIR analysis. *Scientia Horticulturae*, 252(November 2018), 7–13. <https://doi.org/10.1016/j.scienta.2019.02.085>
- Acharya, U. K., Subedi, P. P., & Walsh, K. B. (2017). Robustness of Tomato Quality Evaluation Using a Portable Vis-SWNIRS for Dry Matter and Colour. *International Journal of Analytical Chemistry*, 2017. <https://doi.org/10.1155/2017/2863454>
- Ali, M. M., Janius, R. B., Nawi, N. M., & Hashim, N. (2018). Prediction of total soluble solids and pH in banana using near infrared spectroscopy. *Journal of Engineering Science and Technology*, 13(1), 254–264.
- Anonim. (2018). *How Does a Spectrometer Work*. https://www.youtube.com/results?search_query=how+does+a+spectrometer+work
- Anthon, G. E., & Barrett, D. M. (2006). Potential and limitations for determining lycopene in tomatoes by optical methods. *Acta Horticulturae*, 724, 243–248. <https://doi.org/10.17660/ActaHortic.2006.724.30>
- Anthon, G. E., Lestrangle, M., & Barrett, D. M. (2011). Changes in pH, acids, sugars and other quality parameters during extended vine holding of ripe processing tomatoes. *Journal of the Science of Food and Agriculture*, 91(7), 1175–1181. <https://doi.org/10.1002/jsfa.4312>
- Aquino, C. F., Carlos, L., Salomão, C., & Cecon, P. R. (2017). *Physical, Chemical And Morphological Characteristics Of Banana Cultivars Depending On Maturation Stages*. 2125, 87–96.
- Arendse, E., Fawole, O. A., Magwaza, L. S., Nieuwoudt, H., & Opara, U. L. (2018). Fourier transform near infrared diffuse reflectance spectroscopy and two spectral acquisition modes for evaluation of external and internal quality of intact pomegranate fruit. *Postharvest Biology and Technology*, 138(October 2017), 91–98. <https://doi.org/10.1016/j.postharvbio.2018.01.001>
- Arendse, E., Fawole, O. A., Magwaza, L. S., & Opara, U. L. (2018). Non-destructive prediction of internal and external quality attributes of fruit with thick rind: A review. *Journal of Food Engineering*, 217, 11–23. <https://doi.org/10.1016/j.jfoodeng.2017.08.009>
- Arias, R., Lee, T. C., Logendra, L., & Janes, H. (2000). Correlation of lycopene measured by HPLC with the L*, a*, b* color readings of a hydroponic tomato and the relationship of maturity with color and lycopene content. *Journal of Agricultural and Food Chemistry*, 48(5), 1697–1702. <https://doi.org/10.1021/jf990974e>
- Ashebir, D., Jezik, K., Weingartemann, H., & Gretzmacher, R. (2009). Change in color and other fruit quality characteristics of tomato cultivars after hot-air drying at low final-moisture content. *International Journal of Food Sciences and Nutrition*, 60(SUPPL. 7), 308–315. <https://doi.org/10.1080/09637480903114128>
- Ashokkumar, K., Sivakumar, P., Elayabalan, S., Shobana, V. G., & Pandiyan, M.

- (2018). Nutritional value of cultivars of Banana (*Musa spp.*) and its future prospects. ~ 2972 ~ *Journal of Pharmacognosy and Phytochemistry*, 7(3), 2972–2977.
- Ayvaz, H., Santos, A. M., & Rodriguez-Saona, L. E. (2016). Understanding Tomato Peelability. *Comprehensive Reviews in Food Science and Food Safety*, 15(3), 619–632. <https://doi.org/10.1111/1541-4337.12195>
- Campuzano, A., Rosell, C. M., & Cornejo, F. (2018). Physicochemical and nutritional characteristics of banana flour during ripening. *Food Chemistry*, 256(November 2017), 11–17. <https://doi.org/10.1016/j.foodchem.2018.02.113>
- Cao, Fang, Wu, D., & He, Y. (2010). Soluble solids content and pH prediction and varieties discrimination of grapes based on visible-near infrared spectroscopy. *Computers and Electronics in Agriculture*, 71(SUPPL. 1). <https://doi.org/10.1016/j.compag.2009.05.011>
- Cao, Fei, Guan, C., Dai, H., Li, X., & Zhang, Z. (2015). Soluble solids content is positively correlated with phosphorus content in ripening strawberry fruits. *Scientia Horticulturae*, 195, 183–187. <https://doi.org/10.1016/j.scienta.2015.09.018>
- Cao, N. (2013). Calibration optimization and efficiency in near infrared spectroscopy. *Iowa State University Capstones, Theses and Dissertations*, 183. <http://lib.dr.iastate.edu/etd/13199/>
- Carlini, P., Massantini, R., Mencarelli, F., Carlini, P., Massantini, R., & Mencarelli, F. (2000). Vis-NIR Measurement of Soluble Solids in Cherry and Apricot by PLS Regression and Wavelength Selection Vis-NIR Measurement of Soluble Solids in Cherry and Apricot by PLS Regression and Wavelength Selection. *Journal of Agricultural and Food Chemistry*, 48(11), 5236–5242. <https://doi.org/10.1021/jf000408f>
- Casierra-Posada, F., & Aguilar-Avenidaño, O. E. (2009). Incidence of maturity stage on tomato (*solanum lycopersicum L.*) fruit quality. *Acta Horticulturae*, 821, 229–234. <https://doi.org/10.17660/ActaHortic.2009.821.26>
- Chang, P. T., Hsieh, C. C., & Jiang, Y. L. (2016). Responses of “Shih Huo Chuan” pitaya (*Hylocereus polyrhizus* (Weber) Britt. & Rose) to different degrees of shading nets. *Scientia Horticulturae*, 198, 154–162. <https://doi.org/10.1016/j.scienta.2015.11.024>
- Chauhan, N., & Jethva, K. R. (2008). *Drying Characteristics of Banana Powder*. 7(1), 119–134.
- Chen, C. R., & Ramaswamy, H. S. (2002). Color and texture change kinetics in ripening bananas. *LWT - Food Science and Technology*, 35(5), 415–419. <https://doi.org/10.1006/fstl.2001.0875>
- Chia, K. S., Abdul Rahim, H., & Abdul Rahim, R. (2012). Prediction of soluble solids content of pineapple via non-invasive low cost visible and shortwave near infrared spectroscopy and artificial neural network. *Biosystems Engineering*, 113(2), 158–165. <https://doi.org/10.1016/j.biosystemseng.2012.07.003>
- Chia, K. S., Abdul Rahim, H., & Abdul Rahim, R. (2013). Evaluation of common pre-processing approaches for visible (VIS) and shortwave near infrared

- (SWNIR) spectroscopy in soluble solids content (SSC) assessment. *Biosystems Engineering*, 115(1), 82–88. <https://doi.org/10.1016/j.biosystemseng.2013.02.008>
- Clément, A., Dorais, M., & Vernon, M. (2008). Nondestructive measurement of fresh tomato lycopene content and other physicochemical characteristics using visible NIR spectroscopy. *Journal of Agricultural and Food Chemistry*, 56(21), 9813–9818. <https://doi.org/10.1021/jf801299r>
- Cordeiro, M. H. M., Da Silva, J. M., Mizobutsi, G. P., Mizobutsi, E. H., & Da Mota, W. F. (2015). Caracterização física, química e nutricional da pitaiia-rosa de polpa vermelha. *Revista Brasileira de Fruticultura*, 37(1), 20–26. <https://doi.org/10.1590/0100-2945-046/14>
- Cortés, V., Ortiz, C., Aleixos, N., Blasco, J., Cubero, S., & Talens, P. (2016). A new internal quality index for mango and its prediction by external visible and near-infrared reflection spectroscopy. *Postharvest Biology and Technology*, 118, 148–158. <https://doi.org/10.1016/j.postharvbio.2016.04.011>
- de Freitas, S. T., & Mitcham, E. J. (2013). Quality of pitaya fruit (*Hylocereus undatus*) as influenced by storage temperature and packaging. *Scientia Agricola*, 70(4), 257–262. <https://doi.org/10.1590/S0103-90162013000400006>
- Ding, P., Koi Chew, M., Abdul Aziz, S., Ming Lai, O., & Ong Abdullah, J. (2009). Red-fleshed pitaya (*Hylocereus polyrhizus*) fruit colour and betacyanin content depend on maturity. *International Food Research Journal*, 16(2), 233–242.
- dos Santos Costa, D., Oliveros Mesa, N. F., Santos Freire, M., Pereira Ramos, R., & Teruel Mederos, B. J. (2019). Development of predictive models for quality and maturation stage attributes of wine grapes using vis-nir reflectance spectroscopy. *Postharvest Biology and Technology*, 150(May 2018), 166–178. <https://doi.org/10.1016/j.postharvbio.2018.12.010>
- dos Santos Neto, J. P., de Assis, M. W. D., Casagrande, I. P., Cunha Júnior, L. C., & de Almeida Teixeira, G. H. (2017). Determination of ‘Palmer’ mango maturity indices using portable near infrared (VIS-NIR) spectrometer. *Postharvest Biology and Technology*, 130(May), 75–80. <https://doi.org/10.1016/j.postharvbio.2017.03.009>
- Drissner, D., & Zuercher, U. (2014). Safety of Food and Beverages: Fruits and Vegetables. In *Encyclopedia of Food Safety*. <https://doi.org/10.1016/B978-0-12-378612-8.00280-8>
- Dziezak, J. D. (2003). ACIDS | Natural Acids and Acidulants. In *Encyclopedia of Food Sciences and Nutrition* (pp. 12–17).
- Elfadl, E., Reinbrecht, C., & Claupein, W. (2010). Development of near infrared reflectance spectroscopy (NIRS) calibration model for estimation of oil content in a worldwide safflower germplasm collection. *International Journal of Plant Production*, 4(4), 259–270. <https://doi.org/10.22069/ijpp.2012.709>
- Eyarkai Nambi, V., Thangavel, K., Shahir, S., & Geetha, V. (2015). Evaluation of colour behavior during ripening of Banganapalli mango using CIE-Lab and RGB colour coordinates. *Journal of Applied Horticulture*, 17(3), 205–209. <https://doi.org/10.37855/jah.2015.v17i03.38>

- Farcuh, M., Toubiana, D., Sade, N., Rivero, R. M., Doron-Faigenboim, A., Nambara, E., Sadka, A., & Blumwald, E. (2019). Hormone balance in a climacteric plum fruit and its non-climacteric bud mutant during ripening. *Plant Science*, 280(November 2018), 51–65. <https://doi.org/10.1016/j.plantsci.2018.11.001>
- Fernández-Navales, J., Garde-Cerdán, T., Tardáguila, J., Gutiérrez-Gamboa, G., Pérez-Álvarez, E. P., & Diago, M. P. (2019). Assessment of amino acids and total soluble solids in intact grape berries using contactless Vis and NIR spectroscopy during ripening. *Talanta*, 199(February), 244–253. <https://doi.org/10.1016/j.talanta.2019.02.037>
- Fernández-Navales, J., López, M. I., Sánchez, M. T., Morales, J., & González-Caballero, V. (2009). Shortwave-near infrared spectroscopy for determination of reducing sugar content during grape ripening, winemaking, and aging of white and red wines. *Food Research International*, 42(2), 285–291. <https://doi.org/10.1016/j.foodres.2008.11.008>
- García-Salinas, C., Ramos-Parra, P. A., & Díaz de la Garza, R. I. (2016). Ethylene treatment induces changes in folate profiles in climacteric fruit during postharvest ripening. *Postharvest Biology and Technology*, 118, 43–50. <https://doi.org/10.1016/j.postharvbio.2016.03.011>
- Garner, D., Crisosto, C. H., Wiley, P., & Crisosto, G. M. (2005). Measurement of Soluble Solids Content. *Quality Evaluation Methodology*.
- Gatius, F., Miralbés, C., David, C., & Puy, J. (2017). Chemometrics and Intelligent Laboratory Systems Comparison of CCA and PLS to explore and model NIR data. *Chemometrics and Intelligent Laboratory Systems*, 164(December 2016), 76–82. <https://doi.org/10.1016/j.chemolab.2017.03.011>
- Gayo, J., & Hale, S. A. (2007). Detection and quantification of species authenticity and adulteration in crabmeat using visible and near-infrared spectroscopy. *Journal of Agricultural and Food Chemistry*, 55(3), 585–592. <https://doi.org/10.1021/jf061801+>
- Golic, M., Walsh, K., & Lawson, P. (2003). Short-wavelength near-infrared spectra of sucrose, glucose, and fructose with respect to sugar concentration and temperature. *Applied Spectroscopy*, 57(2), 139–145. <https://doi.org/10.1366/000370203321535033>
- Gómez, A. H., He, Y., & Pereira, A. G. (2006). Non-destructive measurement of acidity, soluble solids and firmness of Satsuma mandarin using Vis/NIR-spectroscopy techniques. *Journal of Food Engineering*, 77(2), 313–319. <https://doi.org/10.1016/j.jfoodeng.2005.06.036>
- Grémy-Gros, C., & Mehinagic, E. (2011). Non-destructive measurements for evaluation of fruit quality. *Qualita 2011, March 2011*. https://www.researchgate.net/publication/235652279_Non-destructive_measurements_for_evaluation_of_fruit_quality
- Guo, Z., Huang, W., Peng, Y., Chen, Q., Ouyang, Q., & Zhao, J. (2016). Color compensation and comparison of shortwave near infrared and long wave near infrared spectroscopy for determination of soluble solids content of “Fuji” apple. *Postharvest Biology and Technology*, 115, 81–90. <https://doi.org/10.1016/j.postharvbio.2015.12.027>

- Hardesty, J. H., & Attili, B. (2010). Spectrophotometry and the Beer-Lambert Law: An Important Analytical Technique in Chemistry. *Collin College*.
Harvard University. (2015). Leaf Pigments | Harvard Forest. In *The President and Fellows of Harvard College*.
<http://harvardforest.fas.harvard.edu/leaves/pigment>
- Hastürk Şahin, F., Ülger, P., Aktas, T., & Orak, H. (2010). Effects of Different Drying Techniques on Some Nutritional Components of Tomato (*Lycopersicon esculentum*). *Tarım Makinaları Bilimi Dergisi*, 6(1), 71–78.
- He, Y., Zhang, Y., Pereira, A. G., Gómez, A. H., & Wang, J. (2005). Nondestructive Determination of Tomato Fruit Quality Characteristics Using Vis / NIR Spectroscopy Technique. *International Journal of Information Technology*, 11(11), 97–108.
- Hu, H., Liu, H. Q., Zhang, H., Zhu, J. H., Yao, X. G., Zhang, X. Bin, & Zheng, K. F. (2010). Assessment of chlorophyll content based on image color analysis, comparison with SPAD-502. *2nd International Conference on Information Engineering and Computer Science - Proceedings, ICIECS 2010*, 476–478.
<https://doi.org/10.1109/ICIECS.2010.5678413>
- Hu, R., Zhang, L., Yu, Z., Zhai, Z., & Zhang, R. (2019). Optimization of soluble solids content prediction models in ‘Hami’ melons by means of Vis-NIR spectroscopy and chemometric tools. *Infrared Physics and Technology*, 102(May), 102999. <https://doi.org/10.1016/j.infrared.2019.102999>
- Huang, Y., Lu, R., & Chen, K. (2018). Assessment of tomato soluble solids content and pH by spatially-resolved and conventional Vis/NIR spectroscopy. *Journal of Food Engineering*, 236(May), 19–28.
<https://doi.org/10.1016/j.jfoodeng.2018.05.008>
- Islam, M., Khan, M., Hoque, M., & Rahman, M. (2012). Studies on the Processing and Preservation of Dragon Fruit (*Hylocereus undatus*) Jelly. *The Agriculturists*, 10(2), 29–35. <https://doi.org/10.3329/agric.v10i2.13139>
- Jaiswal, P., Jha, S. N., & Bharadwaj, R. (2012). Non-destructive prediction of quality of intact banana using spectroscopy. *Scientia Horticulturae*, 135, 14–22. <https://doi.org/10.1016/j.scienta.2011.11.021>
- Jaiswal, P., Jha, S. N., Kaur, P. P., Bhardwaj, R., Singh, A. K., & Wadhawan, V. (2014). Prediction of textural attributes using color values of banana (*Musa sapientum*) during ripening. *Journal of Food Science and Technology*, 51(6), 1179–1184. <https://doi.org/10.1007/s13197-012-0614-2>
- Jamilah, B., Shu, C. E., Kharidah, M., Dzulkifly, M. A., & Noranizan, A. (2011). Physico-chemical characteristics of red pitaya (*Hylocereus polyrhizus*) peel. *International Food Research Journal*, 18(1), 279–286.
- Jamshidi, B., Minaei, S., Mohajerani, E., & Ghassemian, H. (2012). Reflectance Vis/NIR spectroscopy for nondestructive taste characterization of Valencia oranges. *Computers and Electronics in Agriculture*, 85, 64–69.
<https://doi.org/10.1016/j.compag.2012.03.008>
- Janik, L. J., Cozzolino, D., Damberg, R., Cynkar, W., & Gishen, M. (2007). The prediction of total anthocyanin concentration in red-grape homogenates using visible-near-infrared spectroscopy and artificial neural networks. *Analytica Chimica Acta*, 594(1), 107–118. <https://doi.org/10.1016/j.aca.2007.05.019>

- Janse Van Vuuren, J. A., & Groenewald, C. A. (2013). Use of Scanning Near-Infrared Spectroscopy as a Quality Control Indicator for Bulk Blended Inorganic Fertilizers. *Communications in Soil Science and Plant Analysis*, 44(1–4), 120–135. <https://doi.org/10.1080/00103624.2013.736141>
- Jiang, H., Yoon, S. C., Zhuang, H., Wang, W., Li, Y., Lu, C., & Li, N. (2018). Non-destructive assessment of final color and pH attributes of broiler breast fillets using visible and near-infrared hyperspectral imaging: A preliminary study. *Infrared Physics and Technology*, 92(April), 309–317. <https://doi.org/10.1016/j.infrared.2018.06.025>
- Jiang, Y. L., Chen, L. Y., Lee, T. C., & Chang, P. T. (2020). Improving postharvest storage of fresh red-fleshed pitaya (*Hylocereus polyrhizus* sp.) fruit by pre-harvest application of CPPU. *Scientia Horticulturae*, 273(August), 109646. <https://doi.org/10.1016/j.scienta.2020.109646>
- Jie, D., Xie, L., Rao, X., & Ying, Y. (2014). Using visible and near infrared diffuse transmittance technique to predict soluble solids content of watermelon in an on-line detection system. *Postharvest Biology and Technology*, 90, 1–6. <https://doi.org/10.1016/j.postharvbio.2013.11.009>
- Jones, & Bartlett. (1964). Light and the electromagnetic spectrum. *Physics*, 95–122.
- Khodabakhshian, R., Emadi, B., Khojastehpour, M., Golzarian, M. R., & Sazgarnia, A. (2017). Non-destructive evaluation of maturity and quality parameters of pomegranate fruit by visible/near infrared spectroscopy. *International Journal of Food Properties*, 20(1), 41–52. <https://doi.org/10.1080/10942912.2015.1126725>
- Khoo, H. E., Prasad, K. N., Kong, K. W., Jiang, Y., & Ismail, A. (2011). Carotenoids and their isomers: Color pigments in fruits and vegetables. *Molecules*, 16(2), 1710–1738. <https://doi.org/10.3390/molecules16021710>
- Khuriyati, N., & Matsuoka, T. (2004). Near Infrared Transmittance Method for Nondestructive Determination of Soluble Solids Content in Growing Tomato Fruits. *Environment Control in Biology*, 42(3), 217–223. <https://doi.org/10.2525/ecb1963.42.217>
- Kookal, S. K., & Thimmaiah, A. (2018). Nutritional Composition of Staple Food Bananas of Three Cultivars in India. *American Journal of Plant Sciences*, 09(12), 2480–2493. <https://doi.org/10.4236/ajps.2018.912179>
- Kuang, B., & Mouazen, A. M. (2011). Calibration of visible and near infrared spectroscopy for soil analysis at the field scale on three European farms. *European Journal of Soil Science*, 62(4), 629–636. <https://doi.org/10.1111/j.1365-2389.2011.01358.x>
- Kusumiyati, K., Kurniasari, I., & Oktavia, A. R. (2017). The color detection of two cucumber cultivars by NIR Spectroscopy. *Asian Journal of Agriculture*, 1(02), 59–65. <https://doi.org/10.13057/asianjagric/g010202>
- Kusumiyati, Mubarak, S., Hamdani, J. S., Farida, Sutari, W., Hadiwijaya, Y., Putri, I. E., & Mutiarawati, T. (2018). *Evaluation of sapodilla fruit quality using near-infrared spectroscopy*. 49–53. <https://doi.org/10.31227/osf.io/mghf2>
- Kusumiyati, Sutari, W., Farida, Mubarak, S., & Hamdani, J. S. (2019). Prediction of surface color of “crystal” guava using UV-Vis-NIR spectroscopy and multivariate analysis. *IOP Conference Series: Earth and Environmental*

- Science*, 365(1). <https://doi.org/10.1088/1755-1315/365/1/012026>
- Lammertyn, J., Peirs, A., De Baerdemaeker, J., & Nicolaï, B. (2000). Light penetration properties of NIR radiation in fruit with respect to non-destructive quality assessment. *Postharvest Biology and Technology*, 18(2), 121–132. [https://doi.org/10.1016/S0925-5214\(99\)00071-X](https://doi.org/10.1016/S0925-5214(99)00071-X)
- Li, J., Tian, X., Huang, W., Zhang, B., & Fan, S. (2016). Application of Long-Wave Near Infrared Hyperspectral Imaging for Measurement of Soluble Solid Content (SSC) in Pear. *Food Analytical Methods*, 9(11), 3087–3098. <https://doi.org/10.1007/s12161-016-0498-2>
- Li, M., Han, D., & Liu, W. (2019). Non-destructive measurement of soluble solids content of three melon cultivars using portable visible/near infrared spectroscopy. *Biosystems Engineering*, 188, 31–39. <https://doi.org/10.1016/j.biosystemseng.2019.10.003>
- Li, M., Lv, W., Zhao, R., Guo, H., Liu, J., & Han, D. (2017). Non-destructive assessment of quality parameters in ‘Friar’ plums during low temperature storage using visible/near infrared spectroscopy. *Food Control*, 73, 1334–1341. <https://doi.org/10.1016/j.foodcont.2016.10.054>
- Liew, C. Y., & Lau, C. Y. (2012). Determination of quality parameters in Cavendish banana during ripening by NIR spectroscopy. *International Food Research Journal*, 19(2), 751–758.
- Lira, S. M., Dionísio, A. P., Holanda, M. O., Marques, C. G., Silva, G. S. da, Correa, L. C., Santos, G. B. M., de Abreu, F. A. P., Magalhães, F. E. A., Rebouças, E. de L., Guedes, J. A. C., Oliveira, D. F. de, Guedes, M. I. F., & Zocolo, G. J. (2020). Metabolic profile of pitaya (*Hylocereus polyrhizus* (F.A.C. Weber) Britton & Rose) by UPLC-QTOF-MSE and assessment of its toxicity and anxiolytic-like effect in adult zebrafish. *Food Research International*, 127(May 2019). <https://doi.org/10.1016/j.foodres.2019.108701>
- Lismeri, L., Herdiana, N., & Darni, Y. (2019). Diversifikasi Produk Olahan Tomat Sebagai Alternatif “ Camilan Sehat Dan Lezat ” Guna Meningkatkan Nilai Gizi Langkapura Bandar Lampung. *SAKAI SAMBAYAN-Pengabdian Kepada Masyarakat*, 3, 75–82.
- Liu, F., He, Y., & Wang, L. (2008). Comparison of calibrations for the determination of soluble solids content and pH of rice vinegars using visible and short-wave near infrared spectroscopy. *Analytica Chimica Acta*, 610(2), 196–204. <https://doi.org/10.1016/j.aca.2008.01.039>
- Lo’ay, A. A., & EL-Khateeb, A. Y. (2019). Impact of exogenous pyridoxin treatments of ‘Williams’ banana on the activities of starch degrading enzymes. *Scientia Horticulturae*, 243(May 2018), 48–54. <https://doi.org/10.1016/j.scienta.2018.08.007>
- Londoño-Giraldo, L. M., Baena-Pedroza, A. M., Martinez-Seidel, F., Corpas-Iguarán, E., & Taborda-Ocampo, G. (2021). Gone wild: Integration of antioxidative, physicochemical, volatilomic and sensorial profiles ratify rustic relatives of cherry tomato as ideal mating partners. *Scientia Horticulturae*, 277(October 2020). <https://doi.org/10.1016/j.scienta.2020.109814>
- López Camelo, A. F., & Gómez, P. A. (2004). Comparison of color indexes for tomato ripening. *Horticultura Brasileira*, 22(3), 534–537.

- <https://doi.org/10.1590/s0102-05362004000300006>
- Lu, R., & Ariana, D. (2002). A near-infrared sensing technique for measuring internal quality of apple fruit. *Applied Engineering in Agriculture*, *18*(5), 585–590.
- Ma, T., Xia, Y., Inagaki, T., & Tsuchikawa, S. (2021). Non-destructive and fast method of mapping the distribution of the soluble solids content and pH in kiwifruit using object rotation near-infrared hyperspectral imaging approach. *Postharvest Biology and Technology*, *174*(September 2020). <https://doi.org/10.1016/j.postharvbio.2020.111440>
- MacDougall, D. B. (2002). Colour in food. In *Colour in food*. <https://doi.org/10.1533/9781855736672>
- Magalhães, D. S., da Silva, D. M., Ramos, J. D., Salles Pio, L. A., Pasqual, M., Vilas Boas, E. V. B., Galvão, E. C., & de Melo, E. T. (2019). Changes in the physical and physico-chemical characteristics of red-pulp dragon fruit during its development. *Scientia Horticulturae*, *253*(April), 180–186. <https://doi.org/10.1016/j.scienta.2019.04.050>
- Magwaza, L. S., Opara, U. L., Cronje, P. J. R., Landahl, S., Nieuwoudt, H. H., Mouazen, A. M., Nicolai, B. M., & Terry, L. A. (2014). Assessment of rind quality of “Nules Clementine” mandarin fruit during postharvest storage: 2. Robust Vis/NIRS PLS models for prediction of physico-chemical attributes. *Scientia Horticulturae*, *165*, 421–432. <https://doi.org/10.1016/j.scienta.2013.09.050>
- Magwaza, L. S., Opara, U. L., Nieuwoudt, H., Cronje, P. J. R., Saeys, W., & Nicolai, B. (2012). NIR Spectroscopy Applications for Internal and External Quality Analysis of Citrus Fruit-A Review. *Food and Bioprocess Technology*, *5*(2), 425–444. <https://doi.org/10.1007/s11947-011-0697-1>
- Masithoh, R. E., Haff, R., & Kawano, S. (2016). Determination of soluble solids content and titratable acidity of intact fruit and juice of satsuma Mandarin using a hand-held near infrared instrument in transmittance mode. *Journal of Near Infrared Spectroscopy*, *24*(1), 83–88. <https://doi.org/10.1255/jnirs.1196>
- Masithoh, R. E., Lohumi, S., Yoon, W. S., Amanah, H. Z., & Cho, B. K. (2020). Development of multi-product calibration models of various root and tuber powders by fourier transform near infra-red (FT-NIR) spectroscopy for the quantification of polysaccharide contents. *Heliyon*, *6*(10), e05099. <https://doi.org/10.1016/j.heliyon.2020.e05099>
- Masithoh, R. E., Rahardjo, B., Sutiarso, L., & Hadjoko, A. (2012). Development of a simple Computer Vision System to determine tomato quality. *Agritech*, *31*(02). <https://doi.org/10.22146/agritech.9734>
- Merk, H. L., Yarnes, S. C., Van Deynze, A., Tong, N., Menda, N., Mueller, L. A., Mutschler, M. A., Loewen, S. A., Myers, J. R., & Francis, D. M. (2012). Trait diversity and potential for selection indices based on variation among regionally adapted processing Tomato Germplasm. *Journal of the American Society for Horticultural Science*, *137*(6), 427–437. <https://doi.org/10.21273/jashs.137.6.427>
- Merzlyak, M. N., Solovchenko, A. E., & Gitelson, A. A. (2003). Reflectance spectral features and non-destructive estimation of chlorophyll, carotenoid and

- anthocyanin content in apple fruit. *Postharvest Biology and Technology*, 27(2), 197–211. [https://doi.org/10.1016/S0925-5214\(02\)00066-2](https://doi.org/10.1016/S0925-5214(02)00066-2)
- Moradi, M., Rahmatian, M., & Raoufat, M. H. (2017). *Physical Properties of Three Varieties of Tomato , A Comparative Study Physical Properties of Three Varieties of Tomato , a Comparative Study*. June.
- Mouen, J., Chillet, M., Jullien, A., Lapeyre, L. De, & Bellaire, D. (2003). Article original Le gainage précoce des régimes de bananes améliore la croissance des fruits et leur état sanitaire vis-à-vis de l' anthracnose (Colletotrichum musae). *Fruits*, 58(2), 71–81. <https://doi.org/10.1051/fruits>
- Mwende, R., Owino, W., & Imathiu, S. (2018). Effects of pretreatment during drying on the antioxidant properties and color of selected tomato varieties. *Food Science and Nutrition*, 6(2), 503–511. <https://doi.org/10.1002/fsn3.581>
- Newilah, G. N., & Tomekpe, K. (2010). *Physicochemical Changes during Early Fruit Development and Ripening of Three Musa Cultivars in Cameroon*. 401–406.
- Nicola, B. M., Beullens, K., Bobelyn, E., Peirs, A., Saeys, W., Theron, K. I., & Lammertyn, J. (2007). *Nondestructive measurement of fruit and vegetable quality by means of NIR spectroscopy: A review*. 46, 99–118. <https://doi.org/10.1016/j.postharvbio.2007.06.024>
- Nielsen, S. S. (2017). Food Analysis Fifth Edition. In *Food Analysis*. <https://doi.org/10.1038/1841347a0>
- Nordey, T., Davrieux, F., & Léchaudel, M. (2019). Predictions of fruit shelf life and quality after ripening: Are quality traits measured at harvest reliable indicators? *Postharvest Biology and Technology*, 153(June 2018), 52–60. <https://doi.org/10.1016/j.postharvbio.2019.03.011>
- Onsawai, P., Phetpan, K., Khurnpoon, L., & Sirisomboon, P. (2020). Evaluation of physiological properties and texture trait of durian pulp using pulp and intact fruit spectra by near infrared spectroscopic technique. In *Measurement*. Elsevier Ltd. <https://doi.org/10.1016/j.measurement.2020.108684>
- Opara, U. L., Al-Ani, M. R., & Al-Rahbi, N. M. (2012). Effect of Fruit Ripening Stage on Physico-Chemical Properties, Nutritional Composition and Antioxidant Components of Tomato (*Lycopersicum esculentum*) Cultivars. *Food and Bioprocess Technology*, 5(8), 3236–3243. <https://doi.org/10.1007/s11947-011-0693-5>
- Ortiz, T. A., & Takahashi, L. S. A. (2015). Physical and chemical characteristics of pitaya fruits at physiological maturity. *Genetics and Molecular Research*, 14(4), 14422–14439. <https://doi.org/10.4238/2015.November.18.5>
- Phetpan, K., Udompetaikul, V., & Sirisomboon, P. (2018). An online visible and near-infrared spectroscopic technique for the real-time evaluation of the soluble solids content of sugarcane billets on an elevator conveyor. *Computers and Electronics in Agriculture*, 154(September), 460–466. <https://doi.org/10.1016/j.compag.2018.09.033>
- Phuangsoambut, K., Phuangsoambut, A., Talabnark, A., & Terdwongworakul, A. (2018). Empirical reduction of rind effect on rind and flesh absorbance for evaluation of durian maturity using near infrared spectroscopy. *Postharvest Biology and Technology*, 142(January), 55–59.

- <https://doi.org/10.1016/j.postharvbio.2018.04.004>
- Priyankara, G. G. D. S., Karunarathne, C. L. S. M., Sarananda, K. H., & Ariyaratne, M. (2017). Effect of maturity stage on ripening and quality characters of four tomato (*Solanum lycopersicum* L.) varieties of Sri Lanka. *Tropical Agricultural Research*, 28(4), 496. <https://doi.org/10.4038/tar.v28i4.8249>
- Qin, J., Chao, K., & Kim, M. S. (2012). Nondestructive evaluation of internal maturity of tomatoes using spatially offset Raman spectroscopy. *Postharvest Biology and Technology*, 71, 21–31. <https://doi.org/10.1016/j.postharvbio.2012.04.008>
- Rady, A., Fischer, J., Reeves, S., Logan, B., & Watson, N. J. (2020). The Effect of Light Intensity, Sensor Height, and Spectral Pre - Processing Methods when using NIR Spectroscopy to Identify Different Allergen - Containing Powdered Foods. *Sensors*, 20(230). <https://doi.org/10.3390/s20010230>
- Rahman, A., Kandpal, L. M., Lohumi, S., Kim, M. S., Lee, H., Mo, C., & Cho, B. K. (2017). Nondestructive estimation of moisture content, pH and soluble solid contents in intact tomatoes using hyperspectral imaging. *Applied Sciences (Switzerland)*, 7(1). <https://doi.org/10.3390/app7010109>
- Randolph, T. W. (2006). Scale-based normalization of spectral data. *Cancer Biomarkers*, 2(3–4), 135–144. <https://doi.org/10.3233/CBM-2006-23-405>
- Rinnan, Å., Berg, F. van den, & Engelsen, S. B. (2009). Review of the most common pre-processing techniques for near-infrared spectra. In *TrAC - Trends in Analytical Chemistry*. <https://doi.org/10.1016/j.trac.2009.07.007>
- Roy Choudhury, A. K. (2014). Principles of Colour and Appearance Measurement. In *Principles of Colour and Appearance Measurement*. <https://doi.org/10.1016/C2014-0-01832-1>
- Rupawala, M., Dehghani, H., Lucas, S. J. E., Tino, P., & Cruse, D. (2018). Shining a light on awareness: A review of functional near-infrared spectroscopy for prolonged disorders of consciousness. *Frontiers in Neurology*, 9(MAY), 1–17. <https://doi.org/10.3389/fneur.2018.00350>
- Sakuta, M. (2014). Diversity in plant red pigments: Anthocyanins and betacyanins. *Plant Biotechnology Reports*, 8(1), 37–48. <https://doi.org/10.1007/s11816-013-0294-z>
- Sanaeifar, A., Bakhshipour, A., & De La Guardia, M. (2016). Prediction of banana quality indices from color features using support vector regression. *Talanta*, 148, 54–61. <https://doi.org/10.1016/j.talanta.2015.10.073>
- Sánchez, F. D., López, E. M. S., Kerstupp, S. F., Ibarra, R. V., & Scheinvar, L. (2006). Colorant Extraction from Red Prickly Pear (*Opuntia lasiacantha*) for Food Application. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 5(2), 1330–1337.
- Sánchez, M. T., De La Haba, M. J., Benítez-López, M., Fernández-Novales, J., Garrido-Varo, A., & Pérez-Marín, D. (2012). Non-destructive characterization and quality control of intact strawberries based on NIR spectral data. *Journal of Food Engineering*, 110(1), 102–108. <https://doi.org/10.1016/j.jfoodeng.2011.12.003>
- Sandarani, M., Dasanayaka, D., & Jayasinghe, C. (2018). Strategies Used to Prolong the Shelf Life of Fresh Commodities. *Journal of Agricultural Science*

- and *Food Research*, 9(1), 117–118.
<https://doi.org/10.1017/S0021859609990268>
- Santos, G. B. M., Dionísio, A. P., Magalhães, H. C. R., Abreu, F. A. P. de, Lira, S. M., Lima, A. C. V. de, Silva, G. S. da, Guedes, J. A. C., da Silva Araujo, I. M., Artur, A. G., Pontes, D. F., & Zocolo, G. J. (2020). Effects of processing on the chemical, physicochemical, enzymatic, and volatile metabolic composition of pitaya (*Hylocereus polyrhizus* (F.A.C. Weber) Britton & Rose). *Food Research International*, 127(August 2019), 108710.
<https://doi.org/10.1016/j.foodres.2019.108710>
- Sarfraz, M. (2014). *Computer Vision and Image Processing in Intelligent Systems and Multimedia Technologies A volume in the Advances in Computational Intelligence and Robotics (ACIR) Book Series: Vol. i*. <http://www.igi-global.com>
- Sass-Kiss, A., Kiss, J., Milotay, P., Kerek, M. M., & Toth-Markus, M. (2005). Differences in anthocyanin and carotenoid content of fruits and vegetables. *Food Research International*, 38(8–9), 1023–1029.
<https://doi.org/10.1016/j.foodres.2005.03.014>
- Schaare, P. N., & Fraser, D. G. (2000). Comparison of reflectance, interactance and transmission modes of visible-near infrared spectroscopy for measuring internal properties of kiwifruit (*Actinidia chinensis*). *Postharvest Biology and Technology*, 20(2), 175–184. [https://doi.org/10.1016/S0925-5214\(00\)00130-7](https://doi.org/10.1016/S0925-5214(00)00130-7)
- Schneider, A., & Feussner, H. (2007). Chapter 7 - Operative (Surgical) Laparoscopy. In *Biomedical Engineering in Gastrointestinal Surgery* (pp. 269–327). Academic Press.
- Ścibisz, I., Reich, M., Bureau, S., Gouble, B., Causse, M., Bertrand, D., & Renard, C. M. G. C. (2011). Mid-infrared spectroscopy as a tool for rapid determination of internal quality parameters in tomato. *Food Chemistry*, 125(4), 1390–1397. <https://doi.org/10.1016/j.foodchem.2010.10.012>
- Shahir, S., & Visvanathan, R. (2014). Changes in Colour Value of Banana var. Grand Naine during Ripening. *Trends in Biosciences*, 7, 726–728.
- Shao, Y., He, Y., Gómez, A. H., Pereir, A. G., Qiu, Z., & Zhang, Y. (2007). Visible/near infrared spectrometric technique for nondestructive assessment of tomato “Heatwave” (*Lycopersicon esculentum*) quality characteristics. *Journal of Food Engineering*, 81(4), 672–678.
<https://doi.org/10.1016/j.jfoodeng.2006.12.026>
- Sheng, R., Cheng, W., Li, H., Ali, S., Akomeah Agyekum, A., & Chen, Q. (2019). Model development for soluble solids and lycopene contents of cherry tomato at different temperatures using near-infrared spectroscopy. *Postharvest Biology and Technology*, 156(July), 110952.
<https://doi.org/10.1016/j.postharvbio.2019.110952>
- Skalicky, M., Kubes, J., & Shokoofeh, H. (2020). *Betacyanins and Betaxanthins in Cultivated Varieties of Beta vulgaris L. Compared to Weed Beets*. 1–15.
- Someya, S., Yoshiki, Y., & Okubo, K. (2002). Antioxidant compounds from bananas (*Musa Cavendish*). *Food Chemistry*, 79(3), 351–354.
[https://doi.org/10.1016/S0308-8146\(02\)00186-3](https://doi.org/10.1016/S0308-8146(02)00186-3)
- Sugiyama, J. (1999). Visualization of sugar content in the flesh of a melon by near-

- infrared imaging. *Journal of Agricultural and Food Chemistry*, 47(7), 2715–2718. <https://doi.org/10.1021/jf981079i>
- Suhandy, D. (2009). Nondestructive measurement of soluble solids content in pineapple fruit using short wavelength near infrared (SW-NIR) spectroscopy. *International Journal of Applied Engineering Research*, 4(1), 107–114. <http://www.ripublication.com/ijaer.html>
- Taga, G., Homae, F., & Watanabe, H. (2007). Effects of source-detector distance of near infrared spectroscopy on the measurement of the cortical hemodynamic response in infants. *NeuroImage*, 38(3), 452–460. <https://doi.org/10.1016/j.neuroimage.2007.07.050>
- Tee, Y. K., Ding, P., & Rahman, N. A. A. (2011). Physical and cellular structure changes of Rastali banana (*Musa AAB*) during growth and development. *Scientia Horticulturae*, 129(3), 382–389. <https://doi.org/10.1016/j.scienta.2011.03.050>
- Thaiphanit, S., & Anprung, P. (2010). PHysicochemical and flavor changes of fragrant banana (*musa acuminata* aaa group “gross michel”) during ripening. *Journal of Food Processing and Preservation*, 34(3), 366–382. <https://doi.org/10.1111/j.1745-4549.2008.00314.x>
- Tian, S., Zhang, J., Zhang, Z., Zhao, J., Zhang, Z., & Zhang, H. (2019). Effective modification through transmission Vis/NIR spectra affected by fruit size to improve the prediction of moldy apple core. *Infrared Physics and Technology*, 100(March), 117–124. <https://doi.org/10.1016/j.infrared.2019.05.015>
- Travers, S., Bertelsen, M. G., Petersen, K. K., & Kucheryavskiy, S. V. (2014). Predicting pear (cv. Clara Frijs) dry matter and soluble solids content with near infrared spectroscopy. *LWT - Food Science and Technology*, 59(2P1), 1107–1113. <https://doi.org/10.1016/j.lwt.2014.04.048>
- Verboven, S., Hubert, M., & Goos, P. (2012). Robust preprocessing and model selection for spectral data. *Journal of Chemometrics*, 26(6), 282–289. <https://doi.org/10.1002/cem.2446>
- Wang, A., Hu, D., & Xie, L. (2014). Comparison of detection modes in terms of the necessity of visible region (VIS) and influence of the peel on soluble solids content (SSC) determination of navel orange using VIS-SWNIR spectroscopy. *Journal of Food Engineering*, 126, 126–132. <https://doi.org/10.1016/j.jfoodeng.2013.11.011>
- Wang, H., Peng, J., Xie, C., Bao, Y., & He, Y. (2015). Fruit quality evaluation using spectroscopy technology: A review. *Sensors (Switzerland)*, 15(5), 11889–11927. <https://doi.org/10.3390/s150511889>
- Wanitchang, J., Terdwongworakul, A., Wanitchang, P., & Noypitak, S. (2010). Maturity sorting index of dragon fruit: *Hylocereus polyrhizus*. *Journal of Food Engineering*, 100(3), 409–416. <https://doi.org/10.1016/j.jfoodeng.2010.04.025>
- Wasala, W. M. C. B., Dhramasena, D. A. ., Dissanayake, T. M. ., & Thilakarathne, B. M. K. S. (2012). *Physical and Mechanical Properties of Three Commercially Grown Banana (Musa acuminata Colla) Cultivars in Sri Lanka*. 24(1), 42–53.
- Wati, R. K., Fahri, M., Pahlawan, R., & Masithoh, R. E. (2020). Development of

- calibration model for pH content of intact tomatoes using a low-cost Vis / NIR spectroscopy. *IOP Conference Proceedings*.
- Williams, D., Kenyon, A., & Adamson, D. (2010). Chapter Ten - Physiology. In *Basic Science in Obstetrics and Gynaecology* (pp. 173–230). <https://doi.org/10.1016/b978-0-443-10281-3.x0001-2>
- Williams, P., & Norris, K. (2001). Near-infrared technology in the agricultural and food industries. *American Association of Cereal Chemist*.
- Woo, K. K., Ngou, F. H., Ngo, L. S., Soong, W. K., & Tang, P. Y. (2011). Stability of betalain pigment from red dragon fruit (*Hylocereus polyrhizus*). In *American Journal of Food Technology* (Vol. 6, Issue 2, pp. 140–148). <https://doi.org/10.3923/ajft.2011.140.148>
- Xie, L., Ying, Y., Lin, H., Zhou, Y., & Niu, X. (2008). Nondestructive determination of soluble solids content and pH in tomato juice using NIR transmittance spectroscopy. *Sensing and Instrumentation for Food Quality and Safety*, 2(2), 111–115. <https://doi.org/10.1007/s11694-008-9031-7>
- Xie, L., Ying, Y., & Ying, T. (2009). Classification of tomatoes with different genotypes by visible and short-wave near-infrared spectroscopy with least-squares support vector machines and other chemometrics. *Journal of Food Engineering*, 94(1), 34–39. <https://doi.org/10.1016/j.jfoodeng.2009.02.023>
- Xu, S., Lu, B., Baldea, M., Edgar, T. F., & Nixon, M. (2018). An improved variable selection method for support vector regression in NIR spectral modeling. *Journal of Process Control*, 67, 83–93. <https://doi.org/10.1016/j.jprocont.2017.06.001>
- Yang, X. tang, Zhang, Z. qi, Joyce, D., Huang, X. mei, Xu, L. ying, & Pang, X. qun. (2009). Characterization of chlorophyll degradation in banana and plantain during ripening at high temperature. *Food Chemistry*, 114(2), 383–390. <https://doi.org/10.1016/j.foodchem.2008.06.006>
- Yusoff, A., Kumara, N. T. R. N., Lim, A., Ekanayake, P., & Tennakoon, K. U. (2014). Impacts of temperature on the stability of tropical plant pigments as sensitizers for dye sensitized solar cells. *Journal of Biophysics*, 2014. <https://doi.org/10.1155/2014/739514>