

DAFTAR PUSTAKA

- Adams, H. P., Bruno, A., Connors, J. J. B., Demaerschalk, B. M., Khatri, P., McMullan, P. W., *et al.*, (2013). AHA / ASA Guideline Guidelines for the Early Management of Patients With Acute Ischemic Stroke. *Stroke*. doi: 10.1161/STR.0b013e318284056a.
- Akimoto, T., Ito, C., Kato, M., Ogura, M., Muto, S. and Kusano, E. (2011). Reduced hydration status characterized by disproportionate elevation of blood urea nitrogen to serum creatinine among the patients with cerebral infarction. *Med Hyp.*, 77(4), pp. 601–604. doi: 10.1016/j.mehy.2011.06.044.
- Amstrong, L. (2007). Assessing Hydration Status: The Elusive Gold Standard. *J Am Col Nut.*, 26(5), pp. 26 (14): 575-84.
- Appelros, P., Stegmayr, B. and Terent, A. (2009b). Sex differences in stroke epidemiology: A systematic review. *Stroke.*, 40(4), pp. 1082–1090. doi: 10.1161/STROKEAHA.108.540781.
- Asim, M., Alkadi, M. M., Asim, H. and Ghaffar, A. (2019). Dehydration and volume depletion: How to handle the misconceptions. *World J Nephrol.*, 8(1), pp. 23–32. doi: 10.5527/wjn.v8.i1.23.
- Bak, A., Tsiami, A. and Greene, C. (2017). Methods of assessment of hydration status and their usefulness in detecting dehydration in the elderly. *Curr. Res. in Nutr Food Sci Jour.*, 5(2), pp. 43–54. doi: 10.12944/CRNFSJ.5.2.01.
- Baskurt, O. K. and Meiselman, H. J. (2012). Iatrogenic hyperviscosity and thrombosis. *Semin Thrombo Hemost.*, 38(8), pp. 854–864. doi: 10.1055/s-0032-1325616.
- Begum, M. N. and Johnson, C. S. (2010). A review of the literature on dehydration in the institutionalized elderly. *e-SPEN.*, 5(1), pp. e47–e53. doi: 10.1016/j.eclnm.2009.10.007.
- Bevers, M. B., Vaishnav, N. H., Pham, L., Battey, T. W. K. and Kimberly, W. T. (2016). Hyperglycemia is associated with more severe cytotoxic injury after stroke. *J Cer Blood Flow & Met.*, 0(00), pp 1-7. doi: 10.1177/0271678X16671730.
- Bhalla, A., Sankaralingam, S., Dundas, R., Swaminathan, R., Wolfe, C. D. A. and Rudd, A. G. (2000b). Influence of Raised Plasma Osmolality on Clinical Outcome After Acute Stroke. *Stroke.*, pp. 2043–2049.
- Bhatia, K., Mohanty, S., Tripathi, B. K., Gupta, B. and Mittal, M. K. (2015). Predictors of early neurological deterioration in patients with acute ischaemic stroke with special reference to blood urea nitrogen (BUN)/ creatinine ratio & urine specific gravity. *India J Med Res*, (March), pp. 299–307.
- Black-Schaffer, R. M. and Winston, C. (2004). Age and Functional Outcome After Stroke. *Top Stroke Rehabil.*, 11(2), pp. 23–32. doi: 10.1310/DNJU-9VUH-BXU2-DJYU.
- Bravata, D. M., Kim, N., Concato, J. and Brass, L. M. (2003). Hyperglycaemia in patients with acute ischaemic stroke : how often do we screen for undiagnosed diabetes ?. *Q J Med*, 96:pp. 491–497. doi: 10.1093/qjmed/hcg087.
- Broadley, S. A. and Thompson, P. D. (2003). Time to hospital admission for acute stroke : an observational study. *Med J Aust.*, 178(April), pp. 329–331.
- Brott, T., Jr, H. P. A., Olinger, C. P., Marler, J. R., Barsan, W. G., Biller, J., *et al.*, (1989). Measurements of Acute Cerebral Infarction : A Clinical Examination Scale. *Stroke.*, 1, pp. 864–871.
- Carrool, H. and James, L. (2019). Hydration, Arginine Vasopressin, and Glucoregulatory Health in Humans: A Critical Perspective. *Nutr.*, 11. doi: 10.4324/9780203593493-28.

- Chang, S. W., Huang, Y. C., Lin, L. C., Yang, J. T., Weng, H. H., Tsai, Y. H., *et al.*, (2016). Effect of dehydration on the development of collaterals in acute middle cerebral artery occlusion. *Eur Jour Neurol.*, 23(3), pp. 494–500. doi: 10.1111/ene.12841.
- Cuadrado-Godia, E., Ois, A. and Roquer, J. (2010). Heart Failure in Acute Ischemic Stroke. *Curr Cardiol Rev.*, 6(3), pp. 202–213. doi: 10.2174/157340310791658776.
- Dahlan, M. S. (2014). *Statistik untuk Kedokteran dan Kesehatan*. 6th ed. Jakarta: Epidemiologi Indonesia.
- Dahlan, M. S. (2016). *Besar Sampel dalam Penelitian Kedokteran dan Kesehatan*. 4th ed. Jakarta: Epidemiologi Indonesia.
- Danziger, J. and Zeidel, M. L. (2015). Osmotic homeostasis. *Clin J Am Soc Nephrol.*, 10(5), pp. 852–862. doi: 10.2215/CJN.10741013.
- Dasgupta, A. and Wahed, A. (2014). *Clinical Chemistry, Immunology and Laboratory Quality Control*. Elsevier, pp. 197–212. doi: <http://dx.doi.org/10.1016/B978-0-12-407821-5.00011-5>.
- Deb, P., Sharma, S. and Hassan, K. M. (2010). Pathophysiologic mechanisms of acute ischemic stroke: An overview with emphasis on therapeutic significance beyond thrombolysis. *Pathophys.*, 17(3), pp. 197–218. doi: 10.1016/j.pathophys.2009.12.001.
- Elnan, C. and Logallo, N. (2013). Elevated Admission Blood Pressure and Stroke Severity in Acute Ischemic Stroke : The Bergen NORSTROKE Study. *Cerebrovas Dis.*, (5021), pp. 351–354. doi: 10.1159/000355685.
- Etteldorf, J. and Sweeney, M. (2017). *Disturbances of Free Water, Electrolytes, Acid-Base Balance, and Oncotic Pressure*. Elsevier Ltd, p. 113. doi: 10.1016/j.solener.2019.02.027.
- Grabska, K., Gromadzka, G. and Członkowska, A. (2011). Infections and Ischemic Stroke Outcome. *Neurol Res Int.*, 2011, pp. 1–8. doi: 10.1155/2011/691348.
- Harvey, G., Meir, R., Brooks, L. and Holloway, K. (2008). The use of body mass changes as a practical measure of dehydration in team sports. *J Sci Med Sport.*, 11(6), pp. 600–603. doi: 10.1016/j.jsams.2007.05.012.
- Hasibuan, A. E., Fitri, A., Surbakti, K. P. and Rambe, A. S. (2015). Hubungan Kadar Low-Density Lipoprotein Cholesterol dengan Kejadian dan Keparahan Stroke Akut. *Neurona.*, 32(July).
- Hooper, L., Abdelhamid, A., Ali, A., Bunn, D. K., Jennings, A., John, W. G., *et al.* (2015). Diagnostic accuracy of calculated serum osmolality to predict dehydration in older people: adding value to pathology laboratory reports. *BMJ open.*, 5(10), p. e008846. doi: 10.1136/bmjopen-2015-008846.
- Hughes, F., Mythen, M. and Montgomery, H. (2018). The sensitivity of the human thirst response to changes in plasma osmolality: a systematic review. *Periop Med.*, 7(1), pp. 1–11. doi: 10.1186/s13741-017-0081-4.
- Ionita, C. C., Siddiqui, A. H., Levy, E. I., Hopkins, L. N., Snyder, K. V. and Gibbons, K. J. (2011). Acute ischemic stroke and infections. *J Stroke Cerebrovasc Dis.*, 20(1), pp. 1–9. doi: 10.1016/j.jstrokecerebrovasdis.2009.09.011.
- Ishitsuka, K., Kamouchi, M., Hata, J., Fukuda, K., Matsuo, R., Kuroda, J., *et al.*, (2014). High blood pressure after acute ischemic stroke is associated with poor clinical outcomes: Fukuoka stroke registry. *Hypertension.*, 63(1), pp. 54–60. doi: 10.1161/HYPERTENSIONAHA.113.02189.
- Jimenez, C., Melin, B., Koulmann, N., Allevard, A. M., Launay, J. C. and Savourey, G. (1999). Plasma volume changes during and after acute variations of body hydration level in

- humans. *Eur J Appl Physiol.*, 80(1), pp. 1–8. doi: 10.1007/s004210050550.
- Kasner, S. E., Chalela, J. A. and Luciano, J. M. (2007). Reliability and validity of estimating the NIH Stroke Scale score from medical records. *International Classification of Functioning, Disability and Health. Disabil Rehabil.*, 29, pp. 221–230.
- Kavouras, S. A. (2002). Assessing hydration status. *Curr Opin Clin Nutr Metab Care.*, 5 pp. 519–524.
- Kim, H., Lee, K., Choi, H. A., Samuel, S., Park, J. H. and Jo, K. W. (2017). Elevated blood urea nitrogen/creatinine ratio is associated with venous thromboembolism in patients with acute ischemic stroke. *J Korean Neurosurg Soc.*, 60(6), pp. 620–626. doi: 10.3340/jkns.2016.1010.009.
- Kim, W. and Kim, E. J. (2018). Heart Failure as a Risk Factor for Stroke. *Stroke*, 20(1), pp. 33–45.
- Kimura, K. (2005). Atrial fibrillation as a predictive factor for severe stroke and early death in 15 831 patients with acute ischaemic stroke. *J Neurol Neurosurg Psychiatry.*, 76(5), pp. 679–683. doi: 10.1136/jnnp.2004.048827.
- Kreimeier, U. (2000). Pathophysiology of Fluid Imbalance. *Crit care.*, 4(2), pp. 3–7. doi: 10.1186/cc968.
- Kumar, A. D., Boehme, A. K., Siegler, J. E., Gillete, M., Albright, K. C. and Martin-Schild, S. (2016). Leukocytosis in Patients with Neurologic Deterioration after Acute Ischemic Stroke is Associated with Poor Outcomes. *J Stroke Cerebrovasc Dis.*, 72(21), pp. 4139–4156. doi: 10.1007/s00018-015-1997-9.Epigenetic.
- Lee, M., Saver, J. L., Chang, K.-H., Liao, H.-W., Chang, S.-C. and Ovbiagele, B. (2010). Low glomerular filtration rate and risk of stroke: meta-analysis. *Bmj.*, 341, p. c4249. doi: 10.1136/bmj.c4249.
- Li, S.-S., Yin, M.-M., Zhou, Z.-H. and Chen, H.-S. (2017). Dehydration is a strong predictor of long-term prognosis of thrombolysed patients with acute ischemic stroke. *Brain and Behav.*, 7(11), p. e00849. doi: 10.1002/brb3.849.
- Lin, L. C., Yang, J. T., Weng, H. H., Hsiao, C. T., Lai, S. L. and Fann, W. C. (2011). Predictors of early clinical deterioration after acute ischemic stroke. *Am J Emerg Med.*, 29(6), pp. 577–581. doi: 10.1016/j.ajem.2009.12.019.
- Lin, W. C., Shih, H. M. and Lin, L. C. (2015). Preliminary Prospective Study to Assess the Effect of Early Blood Urea Nitrogen/Creatinine Ratio-Based Hydration Therapy on Poststroke Infection Rate and Length of Stay in Acute Ischemic Stroke. *J Stroke Cerebrovasc Dis.*, 24(12), pp. 2720–2727. doi: 10.1016/j.jstrokecerebrovasdis.2015.08.002.
- Liu, C.-H., Wei, Y.-C., Lin, J.-R., Chang, C.-H., Chang, T.-Y., Huang, K.-L., *et al.*, (2016). Initial blood pressure is associated with stroke severity and is predictive of admission cost and one-year outcome in different stroke subtypes: a SRICHS registry study. *BMC Neurol.*, 16(1), p. 27. doi: 10.1186/s12883-016-0546-y.
- Liu, C. H., Lin, S. C., Lin, J. R., Yang, J. T., Chang, Y. J., Chang, C. H., *et al.*, (2014). Dehydration is an independent predictor of discharge outcome and admission cost in acute ischaemic stroke. *Eur J Neurol.*, 21(9), pp. 1184–1191. doi: 10.1111/ene.12452.
- Long, M., Tao, S., Vega, D., Jiang, T., Wen, Q. and Sophia, L. (2016). Persistent Leukocytosis—Is this a Persistent Problem for Patients with Acute Ischemic Stroke? *J Stroke Cerebrovasc Dis.*, 8(5), pp. 444–454. doi: 10.1158/1940-6207.CAPR-14-0359.Nrf2-dependent.
- Martha, S., Collier, L., Davis, S. and Seifert, H. (2019). Translational Evaluation of Acid/Base and Electrolyte Alterations in Rodent Model of Focal Ischemia. *J Stroke Cerebrovasc Dis.*,

- 27(10), pp. 2746–2754. doi: 10.1016/j.jstrokecerebrovasdis.2018.05.045. Translational.
- McManus, M. and Liebeskind, D. S. (2016). Blood Pressure in Acute Ischemic Stroke. *J Clin Neurol.*, 12(2), pp. 137–46. doi: 10.3988/jcn.2016.12.2.137.
- Menon, B. K., Puetz, V., Kochar, P. and Demchuk, A. M. (2011). ASPECTS and Other Neuroimaging Scores in the Triage and Prediction of Outcome in Acute Stroke Patients. *Neuroimag Clin N Am.*, 21(2), pp. 407–423. doi: 10.1016/j.nic.2011.01.007.
- Miller, J. B., Lee, A., Suszanski, J. P., Tustian, M., Corcoran, J. L., Moore, S., *et al.*, (2018). Challenge of intravascular volume assessment in acute ischemic stroke. *Am J Emerg Med.* doi: 10.1016/j.ajem.2018.01.071.
- Minaoki, A., Ghofir, A., Nuradyo, D. (2017). *Korelasi antara indeks aterogenik plasma dengan derajat defisit neurologis pasien stroke iskemik akut*. Tesis. *Unpublished*. Fakultas Kedokteran, Keperawatan dan Kesehatan Masyarakat. Universitas Gadjah Mada: Yogyakarta.
- Mogensen, U. B., Olsen, T. S., Andersen, K. K. and Gerds, T. A. (2013). Cause-specific mortality after stroke: Relation to age, sex, stroke severity, and risk factors in a 10-year follow-up study. *J Stroke Cerebrovasc Dis.*, 22(7), pp. e59–e65. doi: 10.1016/j.jstrokecerebrovasdis.2012.04.006.
- Muchada, M., Rubiera, M., Rodriguez-Luna, D., Pagola, J., Flores, A., Kallas, J., *et al.*, (2014). Baseline national institutes of health stroke scale-adjusted time window for intravenous tissue-type plasminogen activator in acute ischemic stroke. *Stroke.*, 45(4), pp. 1059–1063. doi: 10.1161/STROKEAHA.113.004307.
- Mucke, S., Grottemeyer K., Stahlhut, L., Husstedt, I. W., Evers, S., The Influence of Fluid Intake on Stroke Recurrence- A Prospective Study. *J Neurol Sci.*, 315 (2012) 82-85.
- Nag, C., Das, K., Ghosh, M., Khandajar, M.R., 2012. Plasma osmolality in acute spontaneous intracerebral hemorrhage: Does it influence hematoma volume and clinical outcome?. *J Res Med Sci.* 17(6): 548-551.
- Nardi, K., Milia, P., Eusebi, P., Paciaroni, M., Caso, V. and Agnelli, G. (2012). Admission Leukocytosis in Acute Cerebral Ischemia: Influence on Early Outcome. *J Stroke Cerebrovasc Dis.*, 21(8), pp. 819–824. doi: 10.1016/j.jstrokecerebrovasdis.2011.04.015.
- Nezu, T., Hosomi, N., Kondo, K., Aoki, S., Matsumoto, M. and Kobayashi, S. (2016). Greater Severity of Neurological Defects in Women Admitted With Atrial Fibrillation-Related Stroke. *Circ J.*, 80(1), pp. 250–255. doi: 10.1080/10739149709351455.
- Ortiz, G. . and Sacco, R. . (2008). National Institute of Health Stroke Scale (NIHSS). *Wiley Encyclo Clin Trial.*, pp. 1–9.
- Paper, C. (2008). Guidelines for Management of Ischaemic Stroke and Transient Ischaemic Attack 2008. *Cerebrovasc Dis.*, 25(5), pp. 457–507. doi: 10.1159/000131083.
- Persky, R. W., Turtzo, L. C. and McCullough, L. D. (2010). Stroke in women: Disparities and outcomes. *Curr Cardiol Rep.*, 12(1), pp. 6–13. doi: 10.1007/s11886-009-0080-2.
- Popkin, B. M. and Rosenberg, I. H. (2011). Water, Hydration and Health. *NIH Public Access.*, 68(8), pp. 439–458. doi: 10.1111/j.1753-4887.2010.00304.x. Water.
- Ramírez-Moreno, J. M., Casado-Naranjo, I., Portilla, J. C., Calle, M. L., Tena, D., Falcón, A., *et al.*, (2009). Serum cholesterol LDL and 90-day mortality in patients with intracerebral hemorrhage. *Stroke.*, 40(5), pp. 1917–1920. doi: 10.1161/STROKEAHA.108.536698.
- Rasouli, M. (2016). Basic concepts and practical equations on osmolality: Biochemical approach. *J Clin Biochem.*, 49(12), pp. 936–941. doi: 10.1016/j.clinbiochem.2016.06.001.
- Rodriguez, G. J., Cordina, S. M., Vazquez, G., Suri, M. F. K., Kirmani, J. F., Ezzeddine, M. A., *et*

- al., (2009). The hydration influence on the risk of stroke (THIRST) study. *Neurocrit Care.*, 10(2), pp. 187–194. doi: 10.1007/s12028-008-9169-5.
- Rowat, A., Graham, C. and Dennis, M. (2012). Dehydration in hospital-admitted stroke patients: Detection, frequency, and association. *Stroke.*, 43(3), pp. 857–859. doi: 10.1161/STROKEAHA.111.640821.
- Ryu, C.-W., Shin, H. S., Park, S., Suh, S. H., Koh, J. S. and Choi, H.-Y. (2017). Alberta Stroke Program Early CT Score in the Prognostication after Endovascular Treatment for Ischemic Stroke: A Meta-analysis. *Neurointervent.*, 12(1), pp. 20–30. doi: 10.5469/neuroint.2017.12.1.20.
- Scherbakov, N., Haeusler, K. G., Doehner W., Ischaemic Stroke and Heart Failure: Facts and Numbers. *ESC Heart Fail.*, 2 pp. 1–4
- Schrock, J. W., Glasenapp, M. and Drogell, K. (2012). Elevated blood urea nitrogen/creatinine ratio is associated with poor outcome in patients with ischemic stroke. *Clin Neurol Neurosurg.*, 114(7), pp. 881–884. doi: 10.1016/j.clineuro.2012.01.031.
- Setyopranoto, I., Wibowo, S., Sadewa, A. H., Lamsudin, R. The Interaction of Stroke Risk Factors, Clinical Outcomes and Matrix Mettaloproteinase 9 Levels with the Reduction of Vascular Endothelial Growth Factor A Levels at the Incidence of Brain Edema in Acute Ischemic Stroke Patients. *BJMMR.*, 11(12), pp. 1–14
- Shah, R., Vyas, C. and Vora, J. (2014). NIHSS Score : A handy tool to predict vascular occlusion in acute ischemic stroke. *J Med Sci.*, pp. 18–22.
- Shirreffs, S. M. (2003). Markers of hydration status. *Eur J Clin Nutr.*, 57, pp. S6–S9. doi: 10.1038/sj.ejcn.1601895.
- Sico, J. J., Phipps, M. S., Concato, J., Brandt, C., Wells, C. K., Lo, A. C., *et al.*, (2016). Using Radiological Data to Estimate Ischemic Stroke Severity. *J Stroke Cerebrovasc Dis.*, 25(4), pp. 792–798. doi: 10.1016/j.jstrokecerebrovasdis.2015.12.002.
- Simpson, A., Romer, L. and Kippelen, P. (2017). Exercise-induced dehydration alters pulmonary function but does not modify airway responsiveness to dry air in athletes with mild asthma. *J Applied Physiol.*, 122(5), pp. 1329–1335.
- Sohail, A., Khatri, I. A. and Mehboob, N. (2013). Effect of dyslipidemia on severity and outcome of stroke using mRS scores in Northern Pakistani population. *RMJ.*, 38(4), pp. 345–350.
- Sohrabji, F., Bake, S. and Lewis, D. K. (2013). Age-related changes in brain support cells: Implications for stroke severity. *Neurochem Int.*, 63(4), pp. 291–301. doi: 10.1016/j.neuint.2013.06.013.
- Soriano-tárraga, C., Giralt-steinhauer, E., Mola-caminal, M., Ois, A., Cuadrado-godia, E., Fernández-cadenas, I., *et al.*, (2018). Biological Age is a predictor of mortality in Ischemic Stroke. *Sci Rep.*, pp. 1–8. doi: 10.1038/s41598-018-22579-0.
- Swerdel, J. N., Janevic, T. M., Kostis, W. J., Faiz, A., Cosgrove, N. M. and Kostis, J. B. (2016). Association Between Dehydration and Short-Term Risk of Ischemic Stroke in Patients with Atrial Fibrillation. *Transl Stroke Res.* doi: 10.1007/s12975-016-0471-9.
- Tomita, H., Hagii, J., Metoki, N., Saito, S., Shiroto, H., Hitomi, H., *et al.*, (2015). Impact of Sex Difference on Severity and Functional Outcome in Patients with Cardioembolic Stroke. *J Stroke Cerebrovasc Dis.*, 24(11), pp. 2613–2618. doi: 10.1016/j.jstrokecerebrovasdis.2015.07.016.
- Verbalis JG. Brain volume regulation in response to changes in osmolality. *Neuroscience.* 2010;168:862–70
- Wann, L. S., Curtis, A. B., January, C. T., Ellenbogen, K. A., Lowe, J. E., Estes, N. A. M., *et al.*,

- (2011). 2011 ACCF/AHA/HRS focused update on the management of patients with atrial fibrillation (Updating the 2006 Guideline): A report of the American college of cardiology foundation/American heart association task force on practice guidelines. *Circ.*, 123(1), pp. 104–123. doi: 10.1161/CIR.0b013e3181fa3cf4.
- Wartenberg, K. E., Stoll, A., Funk, A., Meyer, A., Schmidt, J. M. and Berrouschot, J. (2011). Infection after Acute Ischemic Stroke: Risk Factors, Biomarkers, and Outcome. *Stroke Res Treat.*, 2011(1), pp. 1–8. doi: 10.4061/2011/830614.
- Wicaksana, H. C., Adrianto, Y., Rehatta, N. M. Correlation Between White Blood Cell Count and Clinical Severity Based on NIHSS in Acute Ischemic Stroke Patients. *Bali Med J.*, 6(1), pp. 130–134
- Williamson, C. A., Sheehan, K. M., Tipirneni, R., Roark, C. D., Pandey, A. S., Thompson, B. G., *et al.* (2015). The Association Between Spontaneous Hyperventilation, Delayed Cerebral Ischemia, and Poor Neurological Outcome in Patients with Subarachnoid Hemorrhage. *Neurocrit Care.*, 23(3), pp. 330–338. doi: 10.1007/s12028-015-0138-5.
- Xing, Y. (2016). Low Density Lipoprotein Cholesterol and the Outcome of Acute Ischemic Stroke : Results of a Large Hospital-Based Study *Eur Neurol.*, 76(5-6) pp. 195–201. doi: 10.1159/000450604.
- Xu, Tian, Zhang, J. T., Yang, M., Zhang, H., Liu, W. Q., Kong, Y., *et al.*, (2014). Dyslipidemia and outcome in patients with acute ischemic stroke. *Biomed Environ Sci.*, 27(2), pp. 106–10. doi: 10.3967/bes2014.023.
- Yu, J. G., Zhou, R. R. and Cai, G. J. (2011). From Hypertension to Stroke: Mechanisms and Potential Prevention Strategies. *CNS Neurosci Ther.*, 17(5), pp. 577–584. doi: 10.1111/j.1755-5949.2011.00264.x.
- Zanzmera, P., Srivastava Prof., P., Garg, A., Bhatia, R., Singh, M., Tripathi, M., *et al.*, (2012). Prediction of stroke outcome in relation to Alberta Stroke Program early CT score (ASPECTS) at admission in acute ischemic stroke: A prospective study from tertiary care hospital in north India. *Neurol Asia.*, 17(2), pp. 101–107.