

TABLE OF CONTENTS

ACKNOWLEDGEMENT	i
TABLE OF CONTENTS	iii
LIST OF FIGURES	vii
LIST OF TABLES	xvii
LIST OF ABBREVIATION	xviii
ABSTRACT	xxi
CHAPTER 1 – INTRODUCTION	1
1.1 Background	1
1.2 Location and Accessibility	4
1.3 Previous Researchers	7
1.4 Problem Statement	10
1.5 Objectives of Research	10
1.6 Scope of Research	11
1.7 Expected Benefits of Research	12
1.8 Authenticity of Research	13
CHAPTER 2 – LITERATURE REVIEW	14
2.1 Introduction	14
2.2 Geology of Myanmar	15
2.2.1 Tectonic Setting and Tectonic Provinces of Myanmar	15
2.2.2 Magmatism	18
2.2.3 Gold Deposits of Myanmar	23
2.3 Central Volcanic Belt	26
2.3.1 Introduction	26
2.3.2 Tectonic Setting	28
2.3.3 Geological Structure	31
2.3.4 Magmatism	33
2.3.5 Mineralization	36
2.4 Regional Geology of Monywa District	38
2.4.1 Introduction	38
2.4.2 Stratigraphy	40

2.4.3 Geological Structure	45
2.4.4 Mineralization	46
CHAPTER 3 – THEORETICAL BACKGROUND	52
3.1 Epithermal Deposits	52
3.1.1 Low-sulfidation Epithermal Deposits	54
3.1.1.1 Characteristics of Vein Texture	55
3.1.1.2 Ore-Gangue Mineralogy	56
3.1.1.3 Alteration Mineralogy and Zoning	57
3.1.2 High-Sulfidation Epithermal Deposits	58
3.1.2.1 Characteristics of Vein Texture	59
3.1.2.2 Ore-Gangue Mineralogy	59
3.1.2.3 Alteration Mineralogy and Zoning	59
3.1.3 Fluid Inclusion in Epithermal System	61
3.1.3.1 Fluid Inclusion in Low Sulfidation	63
3.1.3.2 Fluid Inclusion in High Sulfidation	64
3.1.4 Origin of Epithermal Deposits	64
3.1.5 Gold Transportation and Deposition in Epithermal System	66
3.2 Hypothesis	67
CHAPTER 4 – RESEARCH METHODOLOGY	69
4.1 Research Methods	69
4.2 Desk Study	69
4.3 Field Investigation	70
4.4 Laboratory Analysis	70
4.4.1 Mineralogical Study	71
4.4.1.1 Petrographic Study from Thin Section	71
4.4.1.2 Ore Microscopy	72
4.4.1.3 X-ray Diffraction	72
4.4.1.4 SEM-EDX Analysis	73
4.4.2 Geochemical Study	74
4.4.2.1 X-ray Fluorescence Analysis	74
4.4.2.2 Inductively Couple Plasma Mass Spectrometry	75
4.4.3 Fluid Inclusion Study (Fluid Inclusion Microthermometry)	75

4.5 Data Analyses and Interpretation	77
CHAPTER 5 – FIELD WORKS AND SAMPLE COLLECTION	81
5.1 Data	81
5.1.1 Field Works	81
5.2 Outcrop Data of Research Areas	82
5.2.1 Kyaukmyet Prospect.....	82
5.2.2 Shwebontha Prospect	84
5.3 Sampling.....	86
CHAPTER 6– LOCAL GEOLOGY	88
6.1 General	88
6.2 Physiography	88
6.3 Geology of the Kyaukmyet Prospect.....	90
6.3.1 Lithostratigraphic Units.....	92
6.3.2 Lithochemistry.....	99
6.3.2.1 Major and Trace Elements Geochemistry.....	99
6.4 Geology of the Shwebontha Prospect	104
6.4.1 Lithostratigraphic Units.....	105
6.4.2 Lithochemistry.....	112
6.4.2.1 Major and Trace Elements Geochemistry.....	112
6.5 Geological Structure.....	115
CHAPTER 7 – ALTERATION AND MINERALIZATION	119
7.1 Introduction	119
7.2 Alteration at Kyaukmyet Prospect	120
7.2.1 Silicic alteration.....	121
7.2.2 Argillic Alteration	123
7.2.3 Propylitic Alteration	126
7.3 Mineralization at Kyaukmyet Prospect	127
7.3.1 Ore and Gangue Mineralogy of the Veins.....	130
7.3.2 Ore and Gangue Minerals Paragenesis.....	138
7.3.3 Quartz Veins Texture	140
7.4 Alteration at Shwebontha Prospect	154
7.4.1 Argillic Alteration	154

7.4.2 Propylitic Alteration	157
7.4.3 Silicification	158
7.5 Mineralization at Shwebontha Prospect	159
7.5.1 Ore and Gangue Mineralogy of the Veins	161
7.5.2 Ore and Gangue Minerals Paragenesis.....	168
7.6 Geochemistry of Ore	170
CHAPTER 8 – FLUID INCLUSION STUDY	173
8.1 Introduction	173
8.2 Fluid Inclusion Petrography	174
8.3 Fluid Inclusion Microthermometry	180
8.4 Interpretation and Discussion.....	184
8.4.1 Hydrothermal Fluid Activities and Evolution Process.....	184
8.4.2 Trapping Pressure and Mineralization Depth.....	190
CHAPTER 9 – DISCUSSION.....	194
9.1 Alteration Minerals.....	194
9.2 Thermal Stability and pH Conditions.....	195
9.3 Ore Fluids and Sulfide Minerals Stabilities	197
9.4 Characteristics of Ore-Forming Fluids.....	202
9.5 Deposit Type and Metal Zonation.....	204
9.6 Genetic Model for the Formation of Epithermal Gold Mineralization System in the Kyaukmyet and Shwebontha Prospects.....	209
CHAPTER 10 – CONCLUSIONS AND RECOMMENDATION	213
10.1 Conclusions	213
10.2 Recommendation.....	216
REFERENCES.....	218
APPENDICES	233
Appendix A: Map of studied areas and sampling location.....	234
Appendix B: Summary of selected sample with details of sample locations and the analyses employed	235
Appendix C: Summary of altered minerals based on XRD results	239
Appendix D: Major oxides compound and trace elements analysis results.....	241
Appendix E: Result summary of fluid inclusion analysis.....	248
Appendix F: SEM-EDX results	254

LIST OF FIGURES

Figure		Page
Figure 1.1	Map illustrating the location of N-S trending Central Volcanic Belt and Mogok Metamorphic Belt with some primary deposits of lead-zin, tin-tungsten and gold (right side). Map showing some of primary gold deposits from sub-units of Central Volcanic Belt (left side) (Modify after Gardiner <i>et al.</i> , 2014 and Mitchell <i>et al.</i> , 1999)	2
Figure 1.2	Location map of the study areas	6
Figure 2.1	Regional tectonic setting and domains of Myanmar	16
Figure 2.2	Map showing magmatism of Myanmar and mainland southeast asia region	22
Figure 2.3	Map showing major tectonic belts and location of primary gold prospects area of Myanmar	24
Figure 2.4	Regional tectonic setting of Myanmar	30
Figure 2.5	Structural map of central Myanmar based on landsat and surface geological mapping	32
Figure 2.6	Sketch map of northern and central Myanmar showing location of Monywa copper deposits relative to main structural features	34
Figure 2.7	Regional geological map of Monywa copper-gold district	40
Figure 2.8	General stratigraphic column of the Monywa copper-gold district	44
Figure 2.9	Schematic northwest-southeast section through the Monywa copper-gold mine district	47
Figure 2.10	Schematic stratigraphic column, Monywa copper-gold deposits	49
Figure 2.11	Cartoon schematic cross-section illustrating mineralized structures and leached cap, and relationship to sand-gravel cover at northern end of Sabetaung pit	50
Figure 3.1	Geological setting and characteristics of low and high sulfidation epithermal deposits	53
Figure 3.2	Sketch diagram showing the mineralogical alteration zoning of low-sulfidation	57
Figure 3.3	Sketch diagram showing the mineralogical alteration zoning of high sulfidation	60
Figure 3.4	Schematic diagram showing typical trends in Th-salinity space due to various fluid evolution processes	62
Figure 4.1	Diagram showing the stages of research methods and materials of proposed research	80
Figure 5.1	Field photographs showing the representative sampling outcrops in Kyaukmyet prospect (a) Massive exposure of rhyolite flow at northeastern part of Yama stream, (b) showing banded rhyolite flow nature	83
Figure 5.2	Field photographs showing the representative sampling outcrops in Kyaukmyet prospect (a,b) Massive exposure of lapilli tuff at northeastern part of Kyaukmyet hill, (c) Massive exposure of tuff at northeastern part of Kyaukmyet hill (d) close view of tuff unit, (e) Colloform-crustiform nature of	

	mineralized quartz vein in rhyolite host rock at Kyaukmyet hill, (f) Close view of colloform-crustiform mineralized quartz vein in rhyolite host rock at Kyaukmyet hill	83
Figure 5.3	Field photographs showing the representative sampling outcrops in Shwebontha prospect (a,c) Massive exposure of rhyolite flow at northeastern part of Shwebontha hill (b,d) close view of rhyolite flow, (e) tuff breccias	85
Figure 5.4	Field photographs showing the representative sampling outcrops in Shwebontha prospect (a,b) Massive exposure of mineralized breccias zone/oxidized zone occurred as open space filling and sulfide leaching at northeastern part of Shwebontha hill, (c,d) Brecciated silicified, pyritized chalcedonic quartz veins alternating with strongly argillized zones cut by cherty or sugary quartz veinlets in the rhyolite host rock occurring at near the Shwebontha Moneasary	86
Figure 6.1	Panoramic view of Monywa copper-gold prospects area (viewing west)	89
Figure 6.2	Morphology and location map of the Monywa Cu-Au prospects area. Elevation shows with color scale bar (meter). (SRTM image with atlas shader)	89
Figure 6.3	Geological map and cross-section (A-B) of the Kyaukmyet prospect (modified after Htet, 2008)	91
Figure 6.4	Field photographs of rhyolite unit showing (a) light grey to ash-white colour of banded rhyolite lava at the northern part of Yama stream, (b) an exposure of yellowish to light grey colour and banded rhyolite lava in the northeastern part of Kyaukmyet hill, (c) flow banding outcrop nature of rhyolite at northwestern part of Kyaukmyet hill, (d) comb quartz and multi quartz veinlet (stockwork vein) in rhyolite rock unit	93
Figure 6.5	Photomicrographs of rhyolite lava showing (a,b) phenocrysts of euhedral plagioclase, K-feldspar and anhedral to subhedral quartz in trachytic texture of rhyolite lava	95
Figure 6.6	Photomicrographs of rhyolite showing (a-d) euhedral quartz phenocrysts with hexagonal outline in rhyolite, plagioclase, K-feldspar and biotite in an aphanitic matrix (cross nicol)	95
Figure 6.7	Field photographs of tuff unit showing (a) ash whitish color to yellowish white colour of tuff, (b) hand specimen of tuff	96
Figure 6.8	Photomicrograph of tuff unit showing (a-d) silicified crystal-dominated tuff (cross nicol). Plagioclase, quartz fragments and opaque minerals observed as in the lithific fragments	96
Figure 6.9	Field photographs of lapilli tuff (a) white to reddish colour ash in fresh surface due to alteration effects, (b) hand specimen sample of lapilli tuff	97
Figure 6.10	Photomicrographs of lapilli tuff showing (a,b) plagioclase, quartz and opaque minerals observed in the lithic fragments, (c) plagioclase, quartz and comb quartz occur as boundary of lithic fragments, (d) quartz, opaque minerals (pyrite) and biotite altered to chlorite occur in the lithic fragments (cross nicol)	98

Figure 6.11	Field photographs of (a,b) silicified sandstone, mudstone and siltstone cut by epithermal auriferous quartz veins with comb quartz at Kyaukmyet hill	99
Figure 6.12	Ti vs Nb/Y plot diagram showing volcanic rock classification (after Pearce, 1996)	100
Figure 6.13	The Zr-TiO ₂ tectonic discrimination diagram for volcanoclastic and volcanic rocks of research area (Peace, 1982)	100
Figure 6.14	Alkali-silica diagram for determining subalkaline and alkaline field and AFM diagram for determining submagma type (tholeiitic or calcalkaline) (Irvine and Baragar, 1971)	101
Figure 6.15	Trace element variation diagram for volcanoclastic and volcanic rocks of Kyaukmyet prospect with Zr	102
Figure 6.16	Chondrite-normalized spider diagrams for volcanic and volcanoclastic rocks from the Kyaukmyet prospect. Using the normalization and ordering scheme of Boynton (1984)	103
Figure 6.17	Geological map and cross-section (A-B) of the Shwebontha prospect (modified after Htet, 2008)	104
Figure 6.18	Field photographs of (a) light to reddish brown color of weathered surface on flow banded rhyolite, (b) ash-white colour of rhyolite flow banded nature, (c) An outcrop of planar flow banding in the rhyolite outcrop, (d) an exposure of strongly flow fold in a rhyolite dyke	106
Figure 6.19	Photomicrographs of rhyolite flow showing (a,b) devitrified flow-banded volcanic fragment. Quartz grain in rhyolite with irregular outline with deep embayments filled with ground mass partly resorbed subhedral to anhedral corroded quartz in an aphanitic matix, (c-d) Quartz grain in rhyolite with irregular outline with deep embayments filled with ground mass, (e,f) partly resorbed euhedral quartz and feldspars phenocrysts in an aphanitic matix	107
Figure 6.20	Field photographs showing exposure of the (a) Light to yellowish brown color of weathered surface on tuffaceous sandstone at Shwebontha prospect, (b) hand specimen sample ash white colour of tuffaceous sandstone	108
Figure 6.21	Photomicrographs of tuffaceous sandstone showing (a,b) subrounded and subangular detrital quartz and feldspar grains surrounded by lithic fragments and clay minerals	108
Figure 6.22	Field photographs showing an exposure of tuff breccia (a) light to reddish brown color weathered surface on angular fragment of tuff breccia at east of Shwebontha Monastery, (b) hand specimen sample of tuff breccia	110
Figure 6.23	Photomicrographs of tuff breccias showing (a,b) quartz, plagioclase, opaque mineral (pyrite) coexisting in the volcanic glass matrix/lithic fragments	110
Figure 6.24	Field photographs showing an exposure of (a) hydrothermal breccia occurs as rusty to reddish colour due to the oxidation of iron sulfide, (b) hand specimen sample of sub-angular to sub-rounded clast of hydrothermal breccia	111

Figure 6.25	Photomicrographs of hydrothermal breccias showing (a,b) coarse-grained pyrite coexisting with quartz and sericite/kaolinite in the lithic fragment	111
Figure 6.26	Classification diagram of volcanic rocks using total alkaline versus SiO ₂ (Middlemost, 1994)	113
Figure 6.27	Classification diagram of Nb/Y vs Zr/TiO ₂ for volcanic rocks of the Shwebontha area (after Pearce, 1996)	113
Figure 6.28	Diagram for (a) discriminate between subalkaline and alkaline fields (Irvine and Baragar,197), (b) AFM diagram for determining submagma type (tholeiitic or calcalkaline) (Irvine and Baragar, 1971)	114
Figure 6.29	Variation in Rb, Nb, Y, Ba, Sr, and Zr (all in ppm) for rhyolite host rocks at Shwebontha area	114
Figure 6.30	Chondrite normalized rare earth elements (REE) diagram for rhyolite rocks from the Shwebontha prospect. Using the normalization and ordering scheme of Nakamura (1974)	115
Figure 6.31	Landsat structural lineament map of Monywa copper-gold prospect area based on satellite map	116
Figure 6.32	An exposure of (a) fault running as a mineralization ENE-WSW trending direction at Kyaukmyet hill, (b) silickenside spperance from fault wall at Shwebontha hill, (c) occurrence of fault scarp at beside the Yama river at Kyaukmyet prospect channel	118
Figure 7.1	Hydrothermal alteration map and cross-section (C-D) of the Kyaukmyet area, Monywa Distirct, Central Myanmar	120
Figure 7.2	An outcrop of (a) fracture filling in amorphous silica in silicic altered zone, (b) massive silica alteration and mineralized sulphide vein at Kyaukmyet hill	122
Figure 7.3	Photomicrographs of silicic alteration showing (a-d) quartz–pyrite veinlet and comb textured quartz crystals within silicic alteration	122
Figure 7.4	Photomicrographs of silicic alteration showing (a,b) adularia is found as intergrowth with quartz and lattice bladed calcite replaced quartz in open space filling	123
Figure 7.5	Photographs showing exposure of (a) argillically altered by flow banded rhyolite exposing at the southern part of the Kyaukmyet prospect, (b) Hand specimen of flow banded rhyolite	124
Figure 7.6	Photomicrographs of argillic alteration showing (a) disseminated pyrite, sericite/kaolinite and fine-grained illite clay mineral under X.P.L (KMR-18), (b) relict of feldspar minerals replaced by sericite/kaolinite, quartz, clay minerals and opaque under X.P.L (KMR-18), (c,d) alteration of plagioclase to sericite, illite and small amount of quartz and pyrite	124
Figure 7.7	X-ray diffractograms of argillic alteration includes quartz-kaolinite-illite/smectite-pyrite (RM=random mount, AD=air-dried and EG=ethylene glycolated) (KMR-18)	125

Figure 7.8	X-ray diffractograms of argillic alteration includes quartz-kaolinite-illite/smectite-pyrite (RM=random mount, AD=air-dried and EG=ethylene glycolated) (KMLP-4)	125
Figure 7.9	Photographs showing chlorite-smectite-epidote alteration in the lapilli tuff unit exposed at the northeastern part of Kyaukmyet hill	126
Figure 7.10	Photomicrographs of propylitic alteration showing (a-d) chlorit/epidote were selectively replaced in plagioclase phenocryst and quartz and opaque mineral (pyrite) occur as fine matrix and anhedral grains	127
Figure 7.11	X-ray diffractograms of prophylic alteration includes quartz-chlorite-illite-pyrite (RM=random mount, AD=air-dried and EG=ethylene glycolated) (KMLP-10)	127
Figure 7.12	Photographs showing (a) quartz veins around 1-2 m of thickness, (b) 100-200 m length rhyolite host rock in Kyaukmyet hill	129
Figure 7.13	Photographs showing (a) Massive ore bodies of brecciated sulfide quartz vein, (b) Crustiform texture of gold-bearing brecciated sulfide quartz vein, (c) Outcrop nature of crustiform-colloform texture of quartz vein in rhyolite host rock, (d) Close view of crustiform-colloform quartz vein in mineralization	129
Figure 7.14	Photographs showing (e) Hand specimen of epithermal quartz vein with angular grey and red-purple siliceous fragments rimmed by comb quartz, (f) hand specimen of massive quartz vein with sulfide clusters	130
Figure 7.15	Photomicrographs of polished thin sections showing primary mineral textures: (a,b) colloform texture pyrite, (c-d) fine to medium grained pyrite disseminated in the gangue matix, (e-f) small grain of galena and chalcopyrite occurred in euhedral large grained in pyrite matrix	131
Figure 7.16	Photomicrographs of polished thin sections showing (a) intergrowth of galena and sphalerite is replaced by in the euhedral pyrite crystal, (b) galena (triangular pits) enclose earlier formed pyrite and galena, (c) anhedral grain of galena occurred in the euhedral pyrite matrix, (d) SEM-EDX image of galena and their elemental composition	132
Figure 7.17	Photomicrographs of polished thin sections showing (a,b) galena and sphalerite occurred in euhedral large grained in pyrite matrix, (c) intergrowth of galena and sphalerite occurred in the pyrite matrix, (d) SEM-EDX image of sphalerite and their elemental composition	134
Figure 7.18	Histograms of FeS atomic fraction of sphalerite grains of samples from KMV-008 vein of Kyaukmyet prospect	135
Figure 7.19	Photomicrographs of polished thin sections showing (a,b) anhedral to subhedral grain of chalcopyrite found in fracture of euhedral pyrite matrix	135

Figure 7.20	Photomicrographs of polished thin sections showing (a) Small grained gold blebs occurs as in the euhedral pyrite crystal, (b) Backscattered image of Au, Ag content of gold grain	136
Figure 7.21	Photomicrographs of polished thin-sections showing (a,b) covellite replacement in framboidal texture pyrite within the vug of the silicic alteration	136
Figure 7.22	Photomicrographs of polished thin sections showing (a-f) concentric growth banding showing sequential development of crustiform-colloform nature of hematite and goethite	138
Figure 7.23	Photographs showing representative various epithermal quartz vein texture of (a) Colloform-crustiform quartz vein in mineralization band and comb quartz texture, (b) Colloform-crustiform quartz vein in mineralization band and lattice blade quartz texture, (c-f) banded (colloform) nature in mineralization	141
Figure 7.24	Photographs showing representative various epithermal quartz veins texture of (a) Lattice bladed calcite replaced by quartz, (b,c) lattice bladed calcite, (d,e) clear to light color of quartz band consisting of tightly packed subparallel crystals oriented perpendicular to the band wall giving the appearance of a teeth like comb, (f) cockade texture which emphasizes the concentric banding	142
Figure 7.25	Photomicrographs primary mineral textures showing (a-d) “Teeth-like” comb quartz growing from opposite sides of the vein wall and interlocking in the center	143
Figure 7.26	Photomicrographs of mineral textures (a,b) euhedral zonal quartz crystals with alternating clear and milky zone parallel to growing crystal faces, (c,d) zonal quartz crystals surrounded by crystalline colloform texture and feathery quartz	144
Figure 7.27	Photomicrographs of mineral textures showing (a,b) concentric crustiform bands of cockade texture	145
Figure 7.28	Photomicrographs showing primary mineral textures (a-d) fine-grained quartz grading into more coarse grained and finally colloform quartz with feathery/flamboyant textures, (e,f) symmetrical colloform quartz with plumose and flamboyant textures, (g,h) symmetrical colloform quartz with mosaic texture	146
Figure 7.29	Photomicrographs of primary mineral textures showing (a-d) crustiform texture of quartz and mineralized pyrite bands	147
Figure 7.30	Photomicrographs of mineral textures showing (a) mosaic extension pattern of the crystalline moss spheres, (b-d) an individual crystalline moss sphere showing concentric rims and colloform nature created by impurities	148
Figure 7.31	Photomicrographs of primary mineral textures (a-d) elongate quartz crystals are randomly distributed in a matrix of smaller quartz crystal	149
Figure 7.32	Photomicrographs of mineral textures showing (a-d) mosaic or jigsaw texture with microcrystalline to crystalline quartz grains	150

Figure 7.33	Photomicrographs of mineral texture showing (a) flameboyant quartz texture, (b) feathery quartz/flamboyant extension	151
Figure 7.34	Photomicrographs of mineral textures showing (a) fine-grained quartz grading into more coarse grained and finally colloform quartz with feathery/flamboyant textures, (b) plumose quartz texture	151
Figure 7.35	Photomicrographs of mineral textures showing (a-d) parallel bladed quartz texture, (e-h) lattice bladed calcite replaced by quartz, (i,j) lattice blade quartz	153
Figure 7.36	Summary of the various silica textures observed in the Kyaukmyet prospect (modified from Bodnar et al., 1985; Dong et al., 1995; Sander and Black, 1988). (a) Lattice-bladed calcite replaced by quartz, (b) Jigsaw texture quartz, (c) Colloform-banded plumose texture quartz, (d) Lattice bladed calcite, (e) Feathery texture quartz, (f) Cockade quartz, (g) Comb quartz, (h) Flamboyant texture quartz, (i) Colloform texture quartz, (j) Zonal quartz, (k) Crustiform quartz texture (l) Moss quartz (XPL= view under crossed polars)	154
Figure 7.37	Photographs showing exposure of (a) ash white color of argillically altered rhyolite at the northeastern of the Shwebontha hill, (b) hand specimen of rhyolite	155
Figure 7.38	Photomicrographs of argillic alteration showing (a) Quartz, illite, illite/smectite and opaque mineral occurred as in the altered rhyolite, (b) altered plagioclase replaced by yellowish brown colour of sericite and kaolinite which is associated with quartz, clay mineral and opaque, (c,d) plagioclase phenocrysts replaced fine-grained quartz, illite, illite/smectite, kaolinite and opaque minerals (pyrite)	155
Figure 7.39	X-ray diffractograms of argillic alteration includes quartz-illite/smectite-pyrite (MSBR-21) (RM=random mount, AD=air-dried and EG=ethylene glycolated, HCL=Hydrochloric acid)	156
Figure 7.40	X-ray diffractograms of argillic alteration includes quartz-kaolinite-illite/smectite-plagioclase (MSBR-16) (RM=random mount, AD=air-dried and EG=ethylene glycolated)	156
Figure 7.41	Field photograph showing (a) Chlorite-illite-epidote alteration in the rhyolite exposed at Shwebontha prospect, (b) hand specimen sample of porphyritic rhyolite	157
Figure 7.42	Photomicrographs showing (a,b) chlorite-illite-epidote, pyrite and quartz were selectively replaced by plagioclase in rhyolite	158
Figure 7.43	X-ray diffractograms of quartz-chlorite-illite-pyrite assemblage from propylitic alteration zone. (AD: air-dried, RM: random mount, EG: ethylene glycolated)	158
Figure 7.44	Photomicrographs showing (a,b) quartz occurs as phenocrysts and silicified groundmass and associated with mineralization (pyrite), hematite and fine-medium grained coexisting with stockwork quartz vein-veinlet and comb quartz	159
Figure 7.45	Photographs showing (a,b) Massive ore bodies in the argillic altered rhyolite host rock occurring at near the Shwebontha	

	Monastery, (c) Hand specimen sample of disseminated sulphides in the rhyolite host rock, (d,e) Hand specimen sample of chalcedonic quartz vein with drusy quartz rhyolite host rock, (f,g,h) Hand specimen samples of gold-bearing brecciated quartz veins at Shwebontha prospect	160
Figure 7.46	Photomicrographs of polished thin sections showing (a,b) anhedral pyrite crystal is infilled the vugs and disseminated in the gangue matrix (MSV-5), (c,d) chalcopryrite and sphalerite occurred in the subhedral to euhedral pyrite crystal as inclusion matrix, (e) Euhedral pyrite (cubic form) occurred in the gangue matrix (MSV-7), (f) SEM-EDX image of pyrite and their elemental composition	162
Figure 7.47	Photomicrographs of polished thin sections showing (a-b) Light grey color of galena develops as bulb and elongated grains in euhedral pyrite matrix, (c) Galena occur as bulb with light grey color in euhedral pyrite matrix (MSV-003), (d) SEM-EDX image of galena and their elemental composition	163
Figure 7.48	Photomicrographs of polished thin sections showing (a) Anhedral grain of sphalerite and galena observed as in the subhedral pyrite matrix as inclusion (MSV-003), (b) sphalerite and pyrite enveloped in the gangue matrix (MSV-009), (c) Sphalerite and pyrite are intergrowth in the gangue matrix (MSV-009), (d) SEM-EDX image of sphalerite and their elemental composition	164
Figure 7.49	Histograms of FeS atomic fraction of sphalerite grains of samples from MSV-009 of Shwebontha prospect	165
Figure 7.50	Photomicrograph of polished thin sections showing (a) chalcopryrite occurred as along the margin of subhedral to euhedral pyrite crystal, (b) anhedral grain of chalcopryrite and galena developed in the euhedral pyrite matrix	166
Figure 7.51	Photomicrographs showing (a) Covellite occurred as in the subhedral to euhedral in the pyrite matrix, (b) Intergrowth of covellite, goethite and pyrite occurred as in the gangue matrix, (c) rhythmic goethite replacing more or less completely pyrite in the gange matrix, (d) bladed crystal of hematite and pyrite in gangue matrix	167
Figure 7.52	Photomicrographs of polished thin sections showing (a,b) large euhedral pyrite crystals replaced by chalcopryrite, galena and fine-grained gold, (c,d) Backscattered electron image and Au, Ag content of electrum grain in chalcedonic quartz veins (MSV-009) and brecciated quartz veins (MSV-006)	168
Figure 7.53	Multi-element plots of gold versus Ag, Cu, As, Sb, Sn, Zn and Pb showing their relation at Kyaukmyet and Shwebontha prospects	172
Figure 8.1	Photomicrographs showing of primary fluid inclusions in quartz from the Kyaukmyet prospect (a-e) two phase liquid-rich fluid inclusion, (f-h) coexisting liquid-rich and vapor-rich inclusions (V=vapor, L=liquid)	176

Figure 8.2	Photomicrographs showing of primary fluid inclusions in quartz from the Shwebontha prospect (a-e) two phase liquid-rich fluid inclusion, (g,h) vapor-rich two-phase inclusions (V=vapor, L=liquid)	177
Figure 8.3	Photomicrographs showing (a-h) secondary necking down fluid inclusion in quartz (KMV-007, KMV-008, KMV-004) for the Kyaukmyet prospect and (MSV-006, MSV-009) for the Shwebontha prospect. (V=vapor, L=liquid)	178
Figure 8.4	Fluid inclusion types observed in samples from the Kyaukmyet prospect (a) containing liquid-rich inclusions with consistent liquid-to-vapor ratio, indicative of non-boiling, (b) coexisting liquid-rich and vapor-rich fluid inclusions, indicative of boiling condition, (c) containing only vapor-rich inclusion indicative of boiling or flashing of the system	179
Figure 8.5	Histograms showing (a) first ice-melting temperatures, (b) total homogenization temperatures, (c) salinities for primary fluid inclusions in mineralized quartz veins of the Kyaukmyet prospect (n=number of analyzed fluid inclusion)	182
Figure 8.6	Histograms showing (a) first ice-melting temperatures, (b) total homogenization temperatures, (c) salinities for primary fluid inclusions in mineralized quartz veins of the Shwebontha prospect (n=number of analyzed fluid inclusion)	183
Figure 8.7	T_h (°C) versus salinity diagram illustrating typical range for fluid inclusions from different deposit types and T_h -salinity space due to various fluid evolution process (Wilkinson, 2001). Rhombus shapes are microthermometric results of Kyaukmyet and Shwebontha fluid inclusions	186
Figure 8.8	Fluid inclusion T_h (°C) versus salinity diagrams of each individual mineralized quartz veins (KMV-007, KMV-008, KMV-004) for the Kyaukmyet prospect and (MSV-006, MSV-009) for the Shwebontha prospect	187
Figure 8.9	Homogenization temperature-salinity diagram of Kyaukmyet and Shwebontha prospects fluid inclusions with field of various water and plotted in the meteoric water area; after Kesler (2005)	188
Figure 8.10	Homogeneous temperature versus salinity diagram of Kyaukmyet and Shwebontha prospects for comparison of stabilities of chloride and sulphide complexes with respect to temperature (Pirajno, 2009)	189
Figure 8.11	Temperature-salinity plot diagram of Kyaukmyet versus Shwebontha fluid inclusions and densities (g/cm ³) of vapor-saturated H ₂ O-NaCl solution (Wilkinson, 2001)	190
Figure 8.12	Temperature versus paleo-depth diagram with boiling-point curves for brine of constant composition given in wt.% NaCl equiv. (after Hass, 1971) showing an estimation of average minimum formation depth of quartz veins from the Kyaukmyet and Shwebontha prospects	193

Figure 8.13	Formation temperature measurements, the inferred pressure is estimated at 16-23 bars for Kyaukmyet prospect and 39-54 bars for Shwebontha prospect (after Hass, 1971)	193
Figure 9.1	Common alteration mineralogy in hydrothermal system (Leach and Corbett, 1995), Diagonal shaded color represented to hydrothermal alteration assemblages of Kyaukmyet and Shwebontha prospects and arrows indicate that their alteration trend	195
Figure 9.2	Thermal stability of various hydrothermal minerals that occur in the epithermal environment under acidic and neutral-pH condition, and the typical temperature range for deposition of epithermal deposit (Reys, 1990, Hedenquist <i>et al.</i> , 1996). Minerals from the Kyaukmyet and Shwebontha prospect areas are blue bold in this figure	197
Figure 9.3	Log fO_2 vs. pH diagram (Heald <i>et al.</i> , 1987). Log fO_2 is constrained from Log fO_2 vs. pH diagram. Deep navy-blue shaded area corresponds to sulfide minerals stability fields of Kyaukmyet and Shwebontha prospect areas	198
Figure 9.4	Estimate range of Log fO_2 and Log fS_2 for Kyaukmyet and Shwebontha deposits. The diagram was set up mainly based on (Heald <i>et al.</i> , 1987)	199
Figure 9.5	Log fS_2 -temperature diagram showing the variation of sulfide assemblages in epithermal deposits that reflect sulphidation state from very low-intermediate-high-very high. Shaded area represents compositional field of arc volcanic rocks, volcanic fumaroles, magmatic-hydrothermal and geothermal (Sillitoe and Hedenquist, 2003). Blue colored rectangle refers to Kyaukmyet and Shwebontha prospects area	200
Figure 9.6	T (°C)-Log fS_2 diagram demonstrating the condition of sulfidation state (Einaudi <i>et al.</i> , 2003). Yellow colored rhombous refers to Kyaukmyet prospect and red colored for Shwebontha prospect	201
Figure 9.7	Salinity (wt.% NaCl eq.) versus homogenization temperatures (Th °C) from Wilkinson (2001) of the Kyaukmyet and Shwebontha prospects	204
Figure 9.8	Scale model for zoning of texture, alteration, ore and gangue mineralogy in a typical boiling zone low to intermediate sulfidation epithermal deposit for the Kyaukmyet prospect	207
Figure 9.9	Genetic Model for the Formation of Epithermal Gold Mineralization System in the Kyaukmyet and Shwebontha prospects	212

LIST OF TABLES

Table		Page
Table 1.1	Previous researchers and authors about gold mineralization in the Monywa copper-gold district, central Myanmar	8
Table 3.1	Classification schemes applied to epithermal deposits	53
Table 7.1	Detected galena grains in mineralized quartz veins with PbS atomic ratio by SEM-EDX analysis	133
Table 7.2	Detected sphalerite grains in mineralized quartz veins with (Zn,Fe)S atomic ratio by SEM-EDX analysis	134
Table 7.3	Generalized paragenesis sequence of the Kyaukmyet prospect	139
Table 7.4	Detected galena grains in mineralized quartz veins with PbS atomic ratio by SEM-EDX analysis	164
Table 7.5	Detected sphalerite grains in mineralized quartz veins with (Zn,Fe)S atomic ratio by SEM-EDX analysis	165
Table 7.6	Detected Au-Ag grains in mineralized quartz veins with Au-Ag atomic ratio by SEM-EDX	168
Table 7.7	Generalized paragenesis sequence of the Shwebontha prospect	169
Table 8.1	Results for Th, Tm, and salinity determination in fluid inclusions data from mineralized quartz veins in the Kyaukmyet and Shwebontha prospect	184
Table 9.1	An overview of the general characteristics of epithermal deposits and a comparison to the features of the Kyaukmyet and Shwebontha deposit	208

LIST OF ABBREVIATION

General Abbreviations and Units

SRTM	-Shuttle Radar Topography Mission
TM	-Thematic Mapper
GIS	-Geographic Information System
CVB	-Central Volcanic Belt
UNDP	-United Nations Development Programme
UN	-United Nations
DGSE	-Department of Geological Survey and Mineral Exploration
HS	-High Sulfidation
IS	-Intermediate Sulfidation
LS	-Low Sulfidation
TAS	-Total Alkali Silica
LOI	-Loss on Ignition
AD	-Air Dried
RM	-Random Mount
EG	-Ethylene glycolated
HCL	-Hydrochloric acid
pH	-Potential of hydrogen
REE	-Rare Earth Elements
P.P.L	-Plane Polarized Light
X.P.L	-Cross Polarized Light
et al.,	-And other
L	-Liquid
V	-Vapor
<i>Th</i>	-Homogenization temperature
<i>T_m</i>	-Last ice melting temperature
<i>T_e</i>	-First ice melting temperature
P	-Pressure

T	-Temperature
X	-Chemical composition
p	-Fluid density
d	-depth
μm	-micrometer
mm	-millimeter
m	-meter
km	-kilometer
wt%	-weight percent
at%	-atomic percent
ppm	-part per million
ppb	-part per billion
°	-Degree
°C	-Degree Celsius
Å	- Angstrom
n.d.	- non-detected
fO_2	-oxygen fugacity
fS_2	-sulfur fugacity
K/Ar	-Potassium-Argon
XRD	-X-ray Diffraction
XRF	-X-ray Fluorescence
ICP-MS	-Inductively Couple Plasma-Mass Spectrometry
SEM-EDX	-Scanning Electron Microscopy with Energy Dispersive X-ray Analysis

Mineral Abbreviation

Adl	-Adularia
Bt	-Biotite
Pl	-Plagioclase

Cal	-Calcite
Ep	-Epidote
Qz	-Quartz
Opq	-Opaque
Ilt	-Illite
Sme	-Smectite
Kln	-Kaolinite
Kfs	-K-feldspar
Ser	-Sericite
Chl	-Chlorite
Lf	-Lithic fragment
Py	-Pyrite
Ccp	-Chalcopyrite
Gn	-Galena
Sph	-Sphalerite
Hem	-Hematite
Gth	-Goethite
Au	-Gold
Elt	-Electrum