

DAFTAR PUSTAKA

- Andrew S. Levey, M., Lesley A. Stevens, MD, MS, F. & Christopher H. Schmid, P. 2009. A New Equation to Estimate Glomerular Filtration Rate. *National Institutes of Health*, 9(150): 604–612.
- Bragadottir, G. & Redfors, B. 2013. Assessing glomerular filtration rate (GFR) in critically ill patients with acute kidney injury - true GFR versus urinary creatinine clearance and estimating equations. *Critical Care*, 17(3).
- Bragadottir, G., Redfors, B. & Ricksten, S. 2013. Assessing glomerular filtration rate (GFR) in critically ill patients with acute kidney injury - true GFR versus urinary creatinine clearance and estimating equations. *Critical Care*, 17(3): R108.
- C Florkowski & Chew-Harris, J.S. 2011. Methods of estimating GFR - Different equations including CKD-EPI. *Clinical Biochemist Reviews*, 32(2): 75–79.
- Chawla, L.S. & Abell, L. 2005. Identifying critically ill patients at high risk for developing acute renal failure : A pilot study. *Kidney International*, 68: 2274–2280.
- Chi, X.H., Li, G.P., Wang, Q.S., Qi, Y.S., Huang, K., Zhang, Q. & Xue, Y.M. 2017. CKD-EPI creatinine-cystatin C glomerular filtration rate estimation equation seems more suitable for Chinese patients with chronic kidney disease than other equations. *BMC Nephrology*, 18(1): 1–7.
- Dasta, J.F., Robert, S. & Zarowitz, B.J. 1991. Is there a reliable index of glomerular filtration rate in critically ill patients? *Annals of Pharmacotherapy*, 25(2): 169–178.
- Delanaye, P. & Mariat, C. 2013. The applicability of eGFR equations to different populations. *Nature Reviews Nephrology*, 9(9): 513–522.
- Dharnidharka, V.R., Kwon, C. & Stevens, G. 2002. Serum cystatin C is superior to serum creatinine as a marker of kidney function: A meta-analysis. *American Journal of Kidney Diseases*, 40(2): 221–226.
- Dinsa, H., Nedi, T. & Berha, A.B. 2017. Concordance between modification of diet in renal disease, chronic kidney disease epidemiology collaboration and Cockcroft-Gault equations in patients with chronic kidney disease at St. Paul's hospital millennium medical college, Addis Ababa, Ethiopia. *BMC Nephrology*, 18(1): 1–13.
- Gault, D.W.C. and M.H. 1976. Prediction of Creatinine Clearance from Serum Creatinine. , 41: 31–41.
- Gelsomino, S., Bonacchi, M., Lucà, F., Barili, F., Del Pace, S., Parise, O., Johnson, D.M. & Gulizia, M.M. 2019. Comparison between three different equations for the estimation of glomerular filtration rate in predicting mortality after coronary artery bypass. *BMC Nephrology*, 20(1): 1–13.
- Hapsari, S.N. & Anniwati, L. 2020. MDRD, CKD-Epi and Creatinine Clearance with 24-Hour Urine Collection Results in Patients with Chronic Kidney Disease. *Indonesian Journal of Clinical Pathology and Medical Laboratory*, 27(1): 66–70.
- Koyner, J.L. 2012. Assessment and diagnosis of renal dysfunction in the ICU. *Chest Medicine*, 141(6): 1584–1594.

- Levey, A.S. & William. 2016. GFR as the “Gold Standard”: Estimated, Measured, and True. *Am J Kidney Disease*, 67(1): 9–12.
- Liu, X., Ma, H., Huang, H., Wang, C., Tang, H., Li, M., Wang, Y. & Lou, T. 2013. Is the chronic kidney disease epidemiology collaboration creatinine-cystatin C equation useful for glomerular filtration rate estimation in the elderly? *Clinical Interventions in Aging*, 8: 1387–1391.
- Macedo, E., Bouchard, J., Soroko, S.H., Chertow, G.M., Himmelfarb, J., Ikizler, T.A., Paganini, E.P. & Mehta, R.L. 2010. Fluid accumulation, recognition and staging of acute kidney injury in critically-ill patients. *Critical Care*, 14(3): 1–7.
- Mangan, C., Stott, M.C. & Dhanda, R. 2018. Renal physiology: blood flow, glomerular filtration and plasma clearance. *Anaesthesia and Intensive Care Medicine*, 19(5): 254–257.
- McMahon, R.S. & Bashir, K. 2020. Anatomy, Abdomen and Pelvis, Kidney Collecting Ducts. *National Institutes of Health*: 1–11.
- Michels, W.M., Grootendorst, D.C., Verduijn, M., Elliott, E.G., Dekker, F.W. & Krediet, R.T. 2010. Performance of the Cockcroft-Gault, MDRD, and new CKD-EPI formulas in relation to GFR, age, and body size. *Clinical Journal of the American Society of Nephrology*, 5(6): 1003–1009.
- Motwani, S.S., Choueiri, T.K., Partridge, A.H., Hu, J., Kaymakcalan, M.D., Waikar, S.S. & Curhan, G.C. 2021. Comparison of Equations To Estimate Glomerular Filtration Rate and Their Impact on Frequency of Cisplatin-associated Acute Kidney Injury. *Kidney360*, 2(2): 205–214.
- Munikrishnappa, D. 2009. Chapter 6 : Limitations of Various Formulae and Other Ways of Assessing GFR in the Elderly : Is There a Role for Cystatin C ? *American Society of Nephrology*: 1–7.
- Murty, M.S.N., Sharma, U.K., Pandey, V.B. & Kankare, S.B. 2013. Serum cystatin C as a marker of renal function in detection of early acute kidney injury. *Indian Journal of Nephrology*, 23(3): 180–183.
- Osegbe, I. & Uche, C. 2017. Sick Cell Patients in Southwest Nigeria. *Nigerian Jou*, 20(8): 16–21.
- Scott, R.P. & Quaggin, S.E. 2015. The cell biology of renal filtration. *The Journal of Cell Biology*, 209(2): 199–210.
- Shemesh, O., Golbetz, H., Kriss, J.P. & Myers, B.D. 1985. Limitations of creatinine as a filtration marker in glomerulopathic patients. *Kidney International*, 28(5): 830–838.
- Soriano, R.M., Penfold, D. & Leslie, S.W. 2021. Anatomy, Abdomen and Pelvis, Kidneys. *National Institutes of Health*: 3–8.
- Stevens, L.A. 2008. Estimating GFR Using Serum Cystatin C Alone and in Combination With Serum Creatinine: A Pooled Analysis of 3,418 Individuals With CKD. *American Journal of Kidney Diseases*, 51(3): 395–406.
- Stevens, L.A., Coresh, J., Greene, T. & Levey, A.S. 2006. Assessing Kidney Function — Measured and Estimated Glomerular Filtration Rate. *New England Journal of Medicine*, 354(23): 2473–2483.
- Sunder, S., Jayaraman, R., Mahapatra, H.S., Sathi, S., Ramanan, V. & Kanchi, P.

2014. Estimation of renal function in the intensive care unit : the covert concepts brought to light. *Journal of Intensive Care*, 4(23): 1–7.
- Udy, A.A., Morton, F.J.A., Nguyen-pham, S., Jarrett, P., Lassig-smith, M., Stuart, J., Dunlop, R., Starr, T., Boots, R.J. & Lipman, J. 2013. A comparison of CKD-EPI estimated glomerular filtration rate and measured creatinine clearance in recently admitted critically ill patients with normal plasma creatinine concentrations. *BMC Nephrology*, 14(250): 1471–2369.
- Vaidya, V.S., Ferguson, M.A. & Bonventre, J. V. 2008. Biomarkers of acute kidney injury. *Annual Review of Pharmacology and Toxicology*, 48: 463–493.
- Wang, F., Ye, P., Luo, L., Xiao, W. & Wu, H. 2010. Association of risk factors for cardiovascular disease and glomerular filtration rate: A community-based study of 4925 adults in Beijing. *Nephrology Dialysis Transplantation*, 25(12): 3924–3931.
- Yokota, L.G., Sampaio, B.M., Rocha, E.P., Balbi, A.L., Prado, I.R.S. & Ponce, D. 2018. Acute kidney injury in elderly patients: Narrative review on incidence, risk factors, and mortality. *International Journal of Nephrology and Renovascular Disease*, 11: 217–224.
- Zhu, Y., Ye, X., Zhu, B., Pei, X., Wei, L., Wu, J. & Zhao, W. 2014. Comparisons between the 2012 new CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration) equations and other four approved equations. *PLoS ONE*, 9(1): 1–10.