

REFERENCES

- Abhilash M, Paul MVS, Varghese M V., Nair RH (2011) Effect of long term intake of aspartame on antioxidant defense status in liver. *Food Chem Toxicol* 49:1203–1207. <https://doi.org/10.1016/j.fct.2011.02.019>
- Ahmad SY, Friel J, Mackay D (2020) The Effects of Non-Nutritive Artificial Sweeteners, Aspartame and Sucralose, on the Gut Microbiome in Healthy Adults: Secondary Outcomes of a Randomized Double-Blinded Crossover Clinical Trial. *Nutrients* 12:3408
- Al-Bukhaiti WQ, Noman A, Qasim AS, Al-Farga A (2017) Gas Chromatography: Principles, Advantages and Applications in Food Analysis. *Int J Agric Innov Res* 6:2319–1473
- Al-Rubaye AF, Hameed IH, Kadhim MJ (2017) A Review: Uses of Gas Chromatography-Mass Spectrometry (GC-MS) Technique for Analysis of Bioactive Natural Compounds of Some Plants. *Int J Toxicol Pharmacol Res* 9:. <https://doi.org/10.25258/ijtpr.v9i01.9042>
- Aly AA, Górecki T (2019) *Green Chemistry and Sustainable Technology Green Analytical Chemistry*. Springer Singapore
- Annissa S, Musfiroh I, Indriati L, et al (2020) *Farmaka Farmaka*. 17:189–197
- Arbelález P, Borrull F, Pocurull E, Marcé RM (2015) Determination of high-intensity sweeteners in river water and wastewater by solid-phase extraction and liquid chromatography-tandem mass spectrometry. *J Chromatogr A* 1393:106–114. <https://doi.org/10.1016/j.chroma.2015.03.035>
- Armenta S, Garrigues S, De La Guardia M (2004) Sweeteners determination in table top formulations using FT-Raman spectrometry and chemometric analysis. *Anal Chim Acta* 521:149–155. <https://doi.org/10.1016/j.aca.2004.05.077>
- Azeez, Alkass, Persike (2019) Long-Term Saccharin Consumption and Increased Risk of Obesity, Diabetes, Hepatic Dysfunction, and Renal Impairment in Rats. *Medicina (B Aires)* 55:681. <https://doi.org/10.3390/medicina55100681>
- Barker SA (2007) Matrix solid phase dispersion (MSPD). *J Biochem Biophys Methods* 70:151–162. <https://doi.org/10.1016/j.jbbm.2006.06.005>
- Bathinapatla A, Kanchi S, Sabela MI, Bisetty K (2015) Theoretical principles and applications of high performance capillary electrophoresis. *Capill Electrophor Princ Challenges Appl* 193–229
- Bergamo AB, Fracassi da Silva JA, de Jesus DP (2011) Simultaneous determination of aspartame, cyclamate, saccharin and acesulfame-K in soft drinks and tabletop sweetener formulations by capillary electrophoresis with capacitively coupled contactless conductivity detection. *Food Chem* 124:1714–1717. <https://doi.org/10.1016/j.foodchem.2010.07.107>
- Berset JD, Ochsenbein N (2012) Stability considerations of aspartame in the direct analysis of artificial sweeteners in water samples using high-performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS). *Chemosphere* 88:563–569. <https://doi.org/10.1016/j.chemosphere.2012.03.030>
- Boroumand S, Chamjangali MA, Bagherian G (2017) Double injection/single

- detection asymmetric flow injection manifold for spectrophotometric determination of ascorbic acid and uric acid: Selection the optimal conditions by MCDM approach based on different criteria weighting methods. *Spectrochim Acta - Part A Mol Biomol Spectrosc* 174:203–213. <https://doi.org/10.1016/j.saa.2016.11.031>
- Bos TS, Knol WC, Molenaar SRA, et al (2020) Recent applications of chemometrics in one- and two-dimensional chromatography. *J Sep Sci* 43:1678–1727. <https://doi.org/10.1002/jssc.202000011>
- Bunaciu AA, Aboul-Enein HY, Fleschin S (2010) Application of fourier transform infrared spectrophotometry in pharmaceutical drugs analysis. *Appl Spectrosc Rev* 45:206–219. <https://doi.org/10.1080/00387011003601044>
- Bursać Kovačević D, Barba FJ, Granato D, et al (2018) Pressurized hot water extraction (PHWE) for the green recovery of bioactive compounds and steviol glycosides from *Stevia rebaudiana* Bertoni leaves. *Food Chem* 254:150–157. <https://doi.org/10.1016/j.foodchem.2018.01.192>
- Cabero C, Saurina J, Hernández-Cassou S (1999) Flow-injection spectrophotometric determination of cyclamate in sweetener products with sodium 1,2-naphthoquinone-4-sulfonate. *Anal Chim Acta* 381:307–313. [https://doi.org/10.1016/S0003-2670\(98\)00763-6](https://doi.org/10.1016/S0003-2670(98)00763-6)
- Camel V (1997) The determination of pesticide residues and metabolites using supercritical fluid extraction. *TrAC - Trends Anal Chem* 16:351–369. [https://doi.org/10.1016/S0165-9936\(97\)00040-X](https://doi.org/10.1016/S0165-9936(97)00040-X)
- Can NÖ, Altiokka G, Aboul-Enein HY (2008) Determination of ticlopidine in pharmaceutical tablets by flow injection analysis using UV-detection. *J Liq Chromatogr Relat Technol* 31:3209–3218. <https://doi.org/10.1080/10826070802480230>
- Cantarelli MA, Pellerano RG, Marchevsky EJ, Camiña JM (2009) Simultaneous determination of aspartame and acesulfame-K by molecular absorption spectrophotometry using multivariate calibration and validation by high performance liquid chromatography. *Food Chem* 115:1128–1132. <https://doi.org/10.1016/j.foodchem.2008.12.101>
- Cárdenas-Soracá DM, Singh V, Nazdrajić E, et al (2020) Development of thin-film solid-phase microextraction coating and method for determination of artificial sweeteners in surface waters. *Talanta* 211:1–9. <https://doi.org/10.1016/j.talanta.2020.120714>
- Chauhan A (2014) GC-MS Technique and its Analytical Applications in Science and Technology. *J Anal Bioanal Tech* 5:. <https://doi.org/10.4172/2155-9872.1000222>
- Chen XH, Zhao YG, Shen HY, Jin MC (2012) Application of dispersive solid-phase extraction and ultra-fast liquid chromatography-tandem quadrupole mass spectrometry in food additive residue analysis of red wine. *J Chromatogr A* 1263:34–42. <https://doi.org/10.1016/j.chroma.2012.09.074>
- Cheng C, Wu SC (2011) Simultaneous analysis of aspartame and its hydrolysis products of Coca-Cola Zero by on-line postcolumn derivation fluorescence detection and ultraviolet detection coupled two-dimensional high-performance liquid chromatography. *J Chromatogr A* 1218:2976–2983.

- <https://doi.org/10.1016/j.chroma.2011.03.033>
- Chi L, Bian X, Gao B, et al (2018) Effects of the Artificial Sweetener Neotame on the Gut Microbiome and Fecal Metabolites in Mice. *Molecules* 23:367. <https://doi.org/10.3390/molecules23020367>
- Choi MMF, Hsu MY, Wong SL (2000) Determination of cyclamate in low-calorie foods by high-performance liquid chromatography with indirect visible photometry. *Analyst* 125:217–220. <https://doi.org/10.1039/a906852j>
- Coskun O (2016) Separation Techniques: CHROMATOGRAPHY. *North Clin Istanbul* 3:156–160. <https://doi.org/10.14744/nci.2016.32757>
- Cummins PM, Rochfort KD, O'Connor B (2016) Ion-Exchange Chromatography: Basic Principles and Application. *Protein Chromatogr* 1485:423. <https://doi.org/10.1007/978-1-4939-6412-3>
- Dias CB, Meinhart AD, Pane DQ, et al (2015) Multivariate Optimisation and Validation of a Method for the Separation of Five Artificial Sweeteners by UPLC-DAD in Nine Food Matrices. *Food Anal Methods* 8:1824–1835. <https://doi.org/10.1007/s12161-014-0056-8>
- Dossi N, Toniolo R, Susmel S, et al (2006) Simultaneous RP-LC determination of additives in soft drinks. *Chromatographia* 63:557–562. <https://doi.org/10.1365/s10337-006-0793-y>
- Duarte LM, Paschoal D, Izumi CMS, et al (2017) Simultaneous determination of aspartame, cyclamate, saccharin and acesulfame-K in powder tabletop sweeteners by FT-Raman spectroscopy associated with the multivariate calibration: PLS, iPLS and siPLS models were compared. *Food Res Int* 99:106–114. <https://doi.org/10.1016/j.foodres.2017.05.006>
- Edwards CH, Rossi M, Corpe CP, et al (2016) The role of sugars and sweeteners in food, diet and health: Alternatives for the future. *Trends Food Sci Technol* 56:158–166. <https://doi.org/10.1016/j.tifs.2016.07.008>
- Erkucuk A, Akgun IH, Yesil-Celiktas O (2009) Supercritical CO₂ extraction of glycosides from *Stevia rebaudiana* leaves: Identification and optimization. *J Supercrit Fluids* 51:29–35. <https://doi.org/10.1016/j.supflu.2009.07.002>
- Ferrer I, Thurman EM (2010) Analysis of sucralose and other sweeteners in water and beverage samples by liquid chromatography/time-of-flight mass spectrometry. *J Chromatogr A* 1217:4127–4134. <https://doi.org/10.1016/j.chroma.2010.02.020>
- Fitch C, Keim KS (2012) Position of the Academy of Nutrition and Dietetics: Use of Nutritive and Nonnutritive Sweeteners. *J Acad Nutr Diet* 112:739–758. <https://doi.org/10.1016/j.jand.2012.03.009>
- Frankenfeld CL, Sikaroodi M, Lamb E, et al (2015) High-intensity sweetener consumption and gut microbiome content and predicted gene function in a cross-sectional study of adults in the United States. *Ann Epidemiol* 25:736–742.e4. <https://doi.org/10.1016/j.annepidem.2015.06.083>
- Frazier RA, Inns EL, Dossi N, et al (2000) Development of a capillary electrophoresis method for the simultaneous analysis of artificial sweeteners, preservatives and colours in soft drinks. *J Chromatogr A* 876:213–220. [https://doi.org/10.1016/S0021-9673\(00\)00184-9](https://doi.org/10.1016/S0021-9673(00)00184-9)
- Gallego R, Bueno M, Herrero M (2019) Sub- and supercritical fluid extraction of

- bioactive compounds from plants, food-by-products, seaweeds and microalgae – An update. *TrAC - Trends Anal Chem* 116:198–213. <https://doi.org/10.1016/j.trac.2019.04.030>
- Gama MR, Melchert WR, Paixão TRLC, Rocha FRP (2019) An overview of the brazilian contributions to green analytical chemistry
- Gan Z, Sun H, Wang R, Feng B (2013a) A novel solid-phase extraction for the concentration of sweeteners in water and analysis by ion-pair liquid chromatography-triple quadrupole mass spectrometry. *J Chromatogr A* 1274:87–96. <https://doi.org/10.1016/j.chroma.2012.11.081>
- Gan Z, Sun H, Wang R, Feng B (2013b) A novel solid-phase extraction for the concentration of sweeteners in water and analysis by ion-pair liquid chromatography-triple quadrupole mass spectrometry. *J Chromatogr A* 1274:87–96. <https://doi.org/10.1016/j.chroma.2012.11.081>
- García-Jiménez JF, Valencia MC, Capitán-Vallvey LF (2007) Simultaneous determination of antioxidants, preservatives and sweetener additives in food and cosmetics by flow injection analysis coupled to a monolithic column. *Anal Chim Acta* 594:226–233. <https://doi.org/10.1016/j.aca.2007.05.040>
- García Jiménez JF, Valencia MC, Cápitan-Vallvey LF (2009) Intense sweetener mixture resolution by flow injection method with on-line monolithic element. *J Liq Chromatogr Relat Technol* 32:1152–1168. <https://doi.org/10.1080/10826070902841885>
- Gui JY, Sun W, Zhang CL, et al (2016) An Innovative Approach to Sensitive Artificial Sweeteners Analysis by Ion Chromatography-Triple Quadrupole Mass Spectrometry. *Chinese J Anal Chem* 44:361–366. [https://doi.org/10.1016/S1872-2040\(16\)60914-3](https://doi.org/10.1016/S1872-2040(16)60914-3)
- Guven G, Samkar H (2019) Examination of Dimension Reduction Performances of PLSR and PCR Techniques in Data with Multicollinearity. *Iran J Sci Technol Trans A Sci* 43:969–978. <https://doi.org/10.1007/s40995-018-0565-1>
- Hashemi M, Habibi A, Jahanshahi N (2011) Determination of cyclamate in artificial sweeteners and beverages using headspace single-drop microextraction and gas chromatography flame-ionisation detection. *Food Chem* 124:1258–1263. <https://doi.org/10.1016/j.foodchem.2010.07.057>
- Hawthorne SB, Grabanski CB, Martin E, Miller DJ (2000) Comparisons of Soxhlet extraction, pressurized liquid extraction, supercritical fluid extraction and subcritical water extraction for environmental solids: Recovery, selectivity and effects on sample matrix. *J Chromatogr A* 892:421–433. [https://doi.org/10.1016/S0021-9673\(00\)00091-1](https://doi.org/10.1016/S0021-9673(00)00091-1)
- Herrmannová M, Křivánková L, Bartoš M, Vytřas K (2006) Direct simultaneous determination of eight sweeteners in foods by capillary isotachopheresis. *J Sep Sci* 29:1132–1137. <https://doi.org/10.1002/jssc.200500441>
- Hooks KB, O'Malley MA (2017) Dysbiosis and its discontents. *MBio* 8:1–11. <https://doi.org/10.1128/mBio.01492-17>
- Horie M, Ishikawa F, Oishi M, et al (2007) Rapid determination of cyclamate in foods by solid-phase extraction and capillary electrophoresis. *J Chromatogr A* 1154:423–428. <https://doi.org/10.1016/j.chroma.2007.03.094>
- Horvath IT, Lantos D (2007) Organometallic chemistry in aqueous and biphasic

- media. *Compr Organomet Chem III* 1:823–845. <https://doi.org/10.1016/b0-08-045047-4/00031-5>
- Hutchinson SA, Ho GS (2009) Stability and degradation of the high - intensity sweeteners: Stability And Degradation of The High-Intensity Sweeteners: Aspartame, . 37–41. <https://doi.org/10.1080/87559129909541189>
- Iwakoshi K, Tahara S, Uematsu Y, et al (2019) Development of a highly sensitive liquid chromatography with tandem mass spectrometry method for the qualitative and quantitative analysis of high-intensity sweeteners in processed foods. *J Chromatogr A* 1592:64–70. <https://doi.org/10.1016/j.chroma.2019.01.036>
- Karunathilaka SR, Yakes BJ, Farris S, et al (2017) Quantitation of Saccharin and Cyclamate in Tabletop Formulations by Portable Raman and NIR Spectrometers in Combination with Partial Least Squares Regression
- Kika FS (2009) Low pressure separations using automated flow and sequential injection analysis coupled to monolithic columns. *J Chromatogr Sci* 47:648–655. <https://doi.org/10.1093/chromsci/47.8.648>
- Ko MJ, Nam HH, Chung MS (2020) Subcritical water extraction of bioactive compounds from *Orostachys japonicus* A. Berger (Crassulaceae). *Sci Rep* 10:1–10. <https://doi.org/10.1038/s41598-020-67508-2>
- Kritsunankul O, Jakmunee J (2011) Simultaneous determination of some food additives in soft drinks and other liquid foods by flow injection on-line dialysis coupled to high performance liquid chromatography. *Talanta* 84:1342–1349. <https://doi.org/10.1016/j.talanta.2011.02.045>
- Kubica P, Namieśnik J, Wasik A (2016) Comparison of hydrophilic interaction and reversed phase liquid chromatography coupled with tandem mass spectrometry for the determination of eight artificial sweeteners and common steviol glycosides in popular beverages. *J Pharm Biomed Anal* 127:184–192. <https://doi.org/10.1016/j.jpba.2016.01.006>
- Lakade SS, Zhou Q, Li A, et al (2018) Hypercrosslinked particles for the extraction of sweeteners using dispersive solid-phase extraction from environmental samples. *J Sep Sci* 41:1618–1624. <https://doi.org/10.1002/jssc.201701113>
- Le AVT, Su YL, Cheng SH (2019) A novel electrochemical assay for aspartame determination via nucleophilic reactions with caffeic acid ortho-quinone. *Electrochim Acta* 300:67–76. <https://doi.org/10.1016/j.electacta.2019.01.020>
- Le THH, Nguyen TQH, Tran CS, et al (2017) Screening determination of food additives using capillary electrophoresis coupled with contactless conductivity detection: A case study in Vietnam. *Food Control* 77:281–289. <https://doi.org/10.1016/j.foodcont.2017.02.020>
- Li J, Liu Y, Liu Q, et al (2017) On-cartridge derivatisation using matrix solid phase dispersion for the determination of cyclamate in foods. *Anal Chim Acta* 972:46–53. <https://doi.org/10.1016/j.aca.2017.04.032>
- Liauchonak I, Qorri B, Dawoud F, et al (2019) Non-nutritive sweeteners and their implications on the development of metabolic syndrome. *Nutrients* 11:1–19. <https://doi.org/10.3390/nu11030644>
- Lim HS, Choi EA, Hwang JY, et al (2018) Improved method for the determination of 12 non-nutritive sweeteners and monitoring in various foods using liquid

- chromatography–tandem mass spectrometry. *Food Addit Contam - Part A Chem Anal Control Expo Risk Assess* 35:1674–1688. <https://doi.org/10.1080/19440049.2018.1486043>
- Llamas NE, Di Nezio MS, Palomeque ME, Fernández Band BS (2005) Automated turbidimetric determination of cyclamate in low calorie soft drinks and sweeteners without pre-treatment. *Anal Chim Acta* 539:301–304. <https://doi.org/10.1016/j.aca.2005.03.044>
- Llamas NE, Di Nezio MS, Palomeque ME, Fernández Band BS (2008) Direct Determination of Saccharin and Acesulfame-K in Sweeteners and Fruit Juices Powders. *Food Anal Methods* 1:43–48. <https://doi.org/10.1007/s12161-007-9006-z>
- Lorenzo RA, Pena MT, Fernández P, et al (2015) Artificial sweeteners in beverages by ultra performance liquid chromatography with photodiode array and liquid chromatography tandem mass spectrometry. *Food Control* 47:43–52. <https://doi.org/10.1016/j.foodcont.2014.06.035>
- Lu L, Hu X, Zhu Z (2017) Trends in Analytical Chemistry Biomimetic sensors and biosensors for qualitative and quantitative analyses of five basic tastes. *Trends Anal Chem* 87:58–70. <https://doi.org/10.1016/j.trac.2016.12.007>
- Mai TD, Le MD, Sáiz J, et al (2016) Triple-channel portable capillary electrophoresis instrument with individual background electrolytes for the concurrent separations of anionic and cationic species. *Anal Chim Acta* 911:121–128. <https://doi.org/10.1016/j.aca.2016.01.029>
- Mazurek S, Szostak R (2011) Quantification of aspartame in commercial sweeteners by FT-Raman spectroscopy. *Food Chem* 125:1051–1057. <https://doi.org/10.1016/j.foodchem.2010.09.075>
- Medeiros RA, Carvalho AE de, Rocha-Filho RC, Fatibello-Filho O (2008) Simultaneous square-wave voltammetric determination of aspartame and cyclamate using a boron-doped diamond electrode. *Talanta* 76:685–689. <https://doi.org/10.1016/j.talanta.2008.04.015>
- Medina DA V, Ferreira APG, Cavalheiro DA V (2019) Polymorphism and thermal behavior of sodium cyclamate. *J Therm Anal Calorim* 7:. <https://doi.org/10.1007/s10973-019-08037-7>
- Medrano LC, Flores-Aguilar JF, Islas G, et al (2019) Solid-Phase Extraction and Large-Volume Sample Stacking-Capillary Electrophoresis for Determination of Artificial Sweeteners in Water Samples. *Food Anal Methods* 12:526–533. <https://doi.org/10.1007/s12161-018-1383-y>
- Mehrotra P (2015) Biosensors and their applications – A review. *J Oral Biol Craniofacial Res* 1–7. <https://doi.org/10.1016/j.jobcr.2015.12.002>
- Memon AF, Palabiyik IM, Solangi AR, et al (2019) Large Volume Sample Stacking (LVSS) in Capillary Electrophoresis (CE) with Response Surface Methodology (RSM) for the Determination of Phenolics in Food Samples. *Anal Lett* 52:2853–2867. <https://doi.org/10.1080/00032719.2019.1624371>
- Mendes CB, Laignier EP, Brigagão MRPL, et al (2010) A simple turbidimetric flow injection system for saccharin determination in sweetener products. *Chem Pap* 64:285–293. <https://doi.org/10.2478/s11696-010-0009-7>
- Miró M, Cerdà V, Estela JM (2002) Multisyringe flow injection analysis:

- Characterization and applications. *TrAC - Trends Anal Chem* 21:199–210. [https://doi.org/10.1016/S0165-9936\(02\)00307-2](https://doi.org/10.1016/S0165-9936(02)00307-2)
- Morris R (2015) Spectrophotometry. *Curr Protoc Essent Lab Tech* 11:2.1.1-2.1.30. <https://doi.org/10.1002/9780470089941.et0201s11>
- Murningsih T, Chairul (2000) Introduction to HPLC : Its Roles on Analysis and Isolation Processes of Natural Chemical Product. *Ber Biol* 5:261–272
- Nambiar AP, Sanyal M, Shrivastav PS (2018) Simultaneous densitometric determination of eight food colors and four sweeteners in candies, jellies, beverages and pharmaceuticals by normal-phase high performance thin-layer chromatography using a single elution protocol. *J Chromatogr A* 1572:152–161. <https://doi.org/10.1016/j.chroma.2018.08.059>
- Nollet LML (2001) Food Analysis by HPLC 2nd ed. In: *Carbohydrate Polymers*. p 197
- Núñez M, Borrull F, Pocurull E, Fontanals N (2017) Pressurised liquid extraction and liquid chromatography-high resolution mass spectrometry to determine high-intensity sweeteners in fish samples. *J Chromatogr A* 1479:32–39. <https://doi.org/10.1016/j.chroma.2016.12.006>
- Núñez O (2015) Preconcentration Methods in Capillary Zone Electrophoresis: Principles Complimentary Contributor Copy
- Oliveira DN De, Menezes M De, Catharino RR (2015) Thermal degradation of sucralose: a combination of analytical methods to determine stability and chlorinated byproducts. 1–5. <https://doi.org/10.1038/srep09598>
- Ordoñez EY, Quintana JB, Rodil R, Cela R (2013) Determination of artificial sweeteners in sewage sludge samples using pressurised liquid extraction and liquid chromatography-tandem mass spectrometry. *J Chromatogr A* 1320:10–16. <https://doi.org/10.1016/j.chroma.2013.10.049>
- Ordóñez EY, Quintana JB, Rodil R, Cela R (2012) Determination of artificial sweeteners in water samples by solid-phase extraction and liquid chromatography-tandem mass spectrometry. *J Chromatogr A* 1256:197–205. <https://doi.org/10.1016/j.chroma.2012.07.073>
- Ordoñez EY, Rodil R, Quintana JB, Cela R (2015) Determination of artificial sweeteners in beverages with green mobile phases and high temperature liquid chromatography-tandem mass spectrometry. *Food Chem* 169:162–168. <https://doi.org/10.1016/j.foodchem.2014.07.132>
- Pacáková V, Štulík K (1997) Capillary electrophoresis of inorganic anions and its comparison with ion chromatography. *J Chromatogr A* 789:169–180. [https://doi.org/10.1016/S0021-9673\(97\)00830-3](https://doi.org/10.1016/S0021-9673(97)00830-3)
- Pérez-Ruiz T, Martínez-Lozano C, Tomás V, et al (2000) Quantitative assay for neohesperidin dihydrochalcone in foodstuffs by capillary electrophoresis. *Chromatographia* 51:385–389. <https://doi.org/10.1007/BF02490473>
- Peruzzi A, Mendez-Lango E, Zhang J, et al (2015) Triple Point of Water. 1–15
- Pierini GD, Llamas NE, Fragoso WD, et al (2013) Simultaneous determination of acesulfame-K and aspartame using linear sweep voltammetry and multivariate calibration. *Microchem J* 106:347–350. <https://doi.org/10.1016/j.microc.2012.09.006>
- Podgornik A, Barut M, Jaksa S, et al (2002) Application of very short monolithic

- columns for separation of low and high molecular mass substances. *J Liq Chromatogr Relat Technol* 25:3099–3116. <https://doi.org/10.1081/JLC-120016211>
- Preveena SM, Cheema MS, Guo H-R (2019) Non-nutritive artificial sweeteners as an emerging contaminant in environment: A global review and risks perspectives. *170:699–707*. <https://doi.org/10.1016/j.ecoenv.2018.12.048>
- Radulescu MC, Bucur B, Bucur MP, Lucian Radu G (2014) Biezymatic biosensor for rapid detection of aspartame by flow injection analysis. *Sensors (Switzerland)* 14:1028–1038. <https://doi.org/10.3390/s140101028>
- Ramos L (2012) *Comprehensive Sampling and Sample Preparation*. Elsevier
- Ramsurn DD, Jhuameer S (2015) Research Article Determination of Artificial Sweeteners in Liquid Foods by High Performance Liquid Chromatography. *Int J Pharm Drug Anal* 3:311–320
- Righetti PG, Guttman AG (2012) *Capillary Electrophoresis Capillary electrophoresis*. eLS John Wiley Sons, Ltd Chichester 1997:561–597. <https://doi.org/10.1002/9780470015902.a0002680.pub2>
- Sakai H, Yamashita A, Tamura M, et al (2015) Simultaneous determination of sweeteners in beverages by LC-MS/MS. *Food Addit Contam - Part A Chem Anal Control Expo Risk Assess* 32:808–816. <https://doi.org/10.1080/19440049.2015.1018341>
- Salameh BA, Al-Degs YS, Abu Safieh KA, AL-Zghool AW (2019) Novel application of multivariate standard addition method based on net analyte signal for quantification of artificial sweeteners in complex food matrices. *J Food Meas Charact* 14:78–87. <https://doi.org/10.1007/s11694-019-00269-3>
- Sampaio PS, Soares A, Castanho A, et al (2018) Optimization of rice amylose determination by NIR-spectroscopy using PLS chemometrics algorithms. *Food Chem* 242:196–204. <https://doi.org/10.1016/j.foodchem.2017.09.058>
- Shah R, Farris S, De Jager LS, Begley TH (2015) A novel method for the simultaneous determination of 14 sweeteners of regulatory interest using UHPLC-MS/MS. *Food Addit Contam - Part A Chem Anal Control Expo Risk Assess* 32:141–151. <https://doi.org/10.1080/19440049.2014.994111>
- Sinan KI, Saftić L, Peršurić Ž, et al (2020) A comparative study of the chemical composition, biological and multivariate analysis of *Crotalaria retusa* L. stem barks, fruits, and flowers obtained via different extraction protocols. *South African J Bot* 128:101–108. <https://doi.org/10.1016/j.sajb.2019.10.019>
- Socas-Rodríguez B, Herrera-Herrera A V., Asensio-Ramos M, Hernández-Borges J (2015) Dispersive Solid-Phase Extraction. *Anal Sep Sci* 1525–1570. <https://doi.org/10.1002/9783527678129.assep056>
- Spanjers CS, Beach CA, Jones AJ, Dauenhauer PJ (2017) Increasing flame ionization detector (FID) sensitivity using post-column oxidation–methanation. *Anal Methods* 9:1928–1934. <https://doi.org/10.1039/c6ay03363f>
- Suarsa IW (2015) *Spektroskopie*. Universitas Udayana
- Suez J, Korem T, Zeevi D, et al (2014) Artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature* 514:181–186. <https://doi.org/10.1038/nature13793>

- Sun XD, Wu HL, Liu Z, et al (2019) Rapid and Sensitive Detection of Multi-Class Food Additives in Beverages for Quality Control by Using HPLC-DAD and Chemometrics Methods. *Food Anal Methods* 12:381–393. <https://doi.org/10.1007/s12161-018-1370-3>
- Teo CC, Tan SN, Yong JWH, et al (2010) Pressurized hot water extraction (PHWE). *J Chromatogr A* 1217:2484–2494. <https://doi.org/10.1016/j.chroma.2009.12.050>
- Thammana M (2016) A review on high performance liquid chromatography. *Int J Pharm Res* 5:1–6. <https://doi.org/10.22214/ijraset.2018.2098>
- Thompson CO, Craige Trenerry V, Kemmery B (1995) Micellar electrokinetic capillary chromatographic determination of artificial sweeteners in low-Joule soft drinks and other foods. *J Chromatogr A* 694:507–514. [https://doi.org/10.1016/0021-9673\(94\)01224-3](https://doi.org/10.1016/0021-9673(94)01224-3)
- Tighrine A, Amir Y, Alfaro P, et al (2019) Simultaneous extraction and analysis of preservatives and artificial sweeteners in juices by salting out liquid-liquid extraction method prior to ultra-high performance liquid chromatography. *Food Chem* 277:586–594. <https://doi.org/10.1016/j.foodchem.2018.10.107>
- Todd R, Baroutian S (2017) A techno-economic comparison of subcritical water, supercritical CO₂ and organic solvent extraction of bioactives from grape marc. *J Clean Prod* 158:349–358. <https://doi.org/10.1016/j.jclepro.2017.05.043>
- Tu X, Chen W (2018) A review on the recent progress in matrix solid phase dispersion. *Molecules* 23:1–13. <https://doi.org/10.3390/molecules23112767>
- Uraisin K, Nacapricha D, Lapanantnoppakhun S, et al (2005) Determination of trace amounts of bromide by flow injection/stopped-flow detection technique using kinetic-spectrophotometric method. *Talanta* 68:274–280. <https://doi.org/10.1016/j.talanta.2005.07.055>
- Van Eyk AD (2015) The effect of five artificial sweeteners on Caco-2, HT-29 and HEK-293 cells. *Drug Chem Toxicol* 38:318–327. <https://doi.org/10.3109/01480545.2014.966381>
- Vazquez-Roig P, Picó Y (2015) Pressurized liquid extraction of organic contaminants in environmental and food samples. *TrAC Trends Anal Chem* 118:709–721. <https://doi.org/10.1016/j.trac.2019.06.038>
- Vistuba JP, Dolzan MD, Vitali L, et al (2015) Sub-minute method for simultaneous determination of aspartame, cyclamate, acesulfame-K and saccharin in food and pharmaceutical samples by capillary zone electrophoresis. *J Chromatogr A* 1396:148–152. <https://doi.org/10.1016/j.chroma.2015.03.070>
- Voss S, Newman E, Miller-Schulze JP (2019) Quantification of sucralose in groundwater well drinking water by silylation derivatization and gas chromatography-mass spectrometry. *Anal Methods* 11:2790–2799. <https://doi.org/10.1039/c9ay00442d>
- Wang YT, Li B, Xu XJ, et al (2020) FTIR spectroscopy coupled with machine learning approaches as a rapid tool for identification and quantification of artificial sweeteners. *Food Chem* 303:.. <https://doi.org/10.1016/j.foodchem.2019.125404>
- Wasik A, McCourt J, Buchgraber M (2007) Simultaneous determination of nine

- intense sweeteners in foodstuffs by high performance liquid chromatography and evaporative light scattering detection-Development and single-laboratory validation. *J Chromatogr A* 1157:187–196. <https://doi.org/10.1016/j.chroma.2007.04.068>
- Wicker AP, Carlton DD, Tanaka K, et al (2018) On-line supercritical fluid extraction—supercritical fluid chromatography-mass spectrometry of polycyclic aromatic hydrocarbons in soil. *J Chromatogr B Anal Technol Biomed Life Sci* 1086:82–88. <https://doi.org/10.1016/j.jchromb.2018.04.014>
- Wiśniewska M, Terpilowski K, Huertas SP, et al (2017) Turbidimetric studies of colloidal silica/ aqueous solution system stability. *Surf Innov*
- Wong DWS (2018) Sweeteners. In: *Mechanism and Theory in Food Chemistry, Second Edition*. Springer International Publishing, Cham, pp 309–325
- Xia S, Yin D, Chen Y, et al (2019) Simultaneous determination of three sulfanilamide artificial sweeteners in foodstuffs by capillary electrophoresis coupled with contactless conductivity detection based on porous aromatic frameworks enhanced solid phase extraction. *Can J Chem* 97:344–351. <https://doi.org/10.1139/cjc-2018-0410>
- Yang DJ, Chen B (2009) Simultaneous determination of nonnutritive sweeteners in foods by HPLC/ESI-MS. *J Agric Food Chem* 57:3022–3027. <https://doi.org/10.1021/jf803988u>
- Yang L, Zhou SJ, Xiao Y, et al (2015) Sensitive simultaneous determination of three sulfanilamide artificial sweeteners by capillary electrophoresis with on-line preconcentration and contactless conductivity detection. *Food Chem* 188:446–451. <https://doi.org/10.1016/j.foodchem.2015.04.060>
- Yasuura M, Okazaki H, Tahara Y, et al (2014) Development of sweetness sensor with selectivity to negatively charged high-potency sweeteners. *Sensors Actuators, B Chem* 201:329–335. <https://doi.org/10.1016/j.snb.2014.04.087>
- Yebra-Biurrun MC (2000) Flow injection determinations of artificial sweeteners: A review. *Food Addit Contam* 17:733–738. <https://doi.org/10.1080/026520300415264>
- Yebra MC, Bermejo P (1998) Indirect determination of cyclamate by an on-line continuous precipitation-dissolution flow system. *Talanta* 45:1115–1122. [https://doi.org/10.1016/S0039-9140\(97\)00219-1](https://doi.org/10.1016/S0039-9140(97)00219-1)
- Yildiz-Ozturk E, Tag O, Yesil-Celiktas O (2014) Subcritical water extraction of steviol glycosides from *Stevia rebaudiana* leaves and characterization of the raffinate phase. *J Supercrit Fluids* 95:422–430. <https://doi.org/10.1016/j.supflu.2014.10.017>
- Yoda SK, Marques MOM, Petenate AJ, Meireles MAA (2003) Supercritical fluid extraction from *Stevia rebaudiana* Bertoni using CO₂ and CO₂ + water: Extraction kinetics and identification of extracted components. *J Food Eng* 57:125–134. [https://doi.org/10.1016/S0260-8774\(02\)00281-9](https://doi.org/10.1016/S0260-8774(02)00281-9)
- Yoshikawa K, Saito S, Sakuragawa A (2011) Simultaneous analysis of acidulants and preservatives in food samples by using capillary zone electrophoresis with indirect UV detection. *Food Chem* 127:1385–1390. <https://doi.org/10.1016/j.foodchem.2011.01.126>
- Yu S, Zhu B, Lv F, et al (2012) Rapid analysis of cyclamate in foods and beverages

- by gas chromatography-electron capture detector (GC-ECD). *Food Chem* 134:2424–2429. <https://doi.org/10.1016/j.foodchem.2012.04.028>
- Zeynep F, Sifa T (2014) Determination of the effects of some artificial sweeteners on human peripheral lymphocytes using the comet assay. *J Toxicol Environ Heal Sci* 6:147–153. <https://doi.org/10.5897/JTEHS2014.0313>
- Zhang F, Zhang Q, Zhang D, et al (2014) Biosensor analysis of natural and artificial sweeteners in intact taste epithelium. *Biosens Bioelectron* 54:385–392. <https://doi.org/10.1016/j.bios.2013.11.020>
- Zhao YG, Cai MQ, Chen XH, et al (2013) Analysis of nine food additives in wine by dispersive solid-phase extraction and reversed-phase high performance liquid chromatography. *Food Res Int* 52:350–358. <https://doi.org/10.1016/j.foodres.2013.03.038>
- Zhu Y, Guo Y, Ye M, James FS (2005) Separation and simultaneous determination of four artificial sweeteners in food and beverages by ion chromatography. *J Chromatogr A* 1085:143–146. <https://doi.org/10.1016/j.chroma.2004.12.042>
- Zygler A, Wasik A, Namieśnik J (2009) Analytical methodologies for determination of artificial sweeteners in foodstuffs. *TrAC - Trends Anal Chem* 28:1082–1102. <https://doi.org/10.1016/j.trac.2009.06.008>