



DAFTAR PUSTAKA

- Abdul Gafor, A. H. *et al.* (2018) ‘The role of reticulocyte hemoglobin content in the management of iron deficiency anemia in patients on hemodialysis’, *Nephro-Urology Monthly*, 10(3), pp. 3–7. doi: 10.5812/numonthly.65629.
- Abramowski, S. W. *et al.* (2013) ‘Diagnostic markers of iron deficiency : Which should we choose ?’, *Revue Medicale Suisse*, 9(373), pp. 380–383.
- Achebe, M. M. and Gafter-Gvili, A. (2017) ‘How I treat anemia in pregnancy: Iron, cobalamin, and folate’, *Blood*, 129(8), pp. 940–949. doi: 10.1182/blood-2016-08-672246.
- Anderson, G. J. (2009) ‘Iron absorption and metabolism’, *Current Opinion in Gastroenterology*, 25(2), pp. 129–135. doi: 10.1097/MOG.0b013e32831ef1f7.
- Andriastuti, M., Adiwijaya, M. and Satari, H. I. (2019) ‘Diagnosis of iron deficiency and iron deficiency anemia with reticulocyte hemoglobin content among children aged 6-18 years’, *Iranian Journal of Blood and Cancer*, 11(4), pp. 127–132.
- Andro, M. *et al.* (2013) ‘Anaemia and cognitive performances in the elderly: A systematic review’, *European Journal of Neurology*, 20(9), pp. 1234–1240. doi: 10.1111/ene.12175.
- Auerbach, M. and Adamson, J. W. (2016) ‘How we diagnose and treat iron deficiency anemia’, *American Journal of Hematology*, 91(1), pp. 31–38. doi: 10.1002/ajh.24201.
- Bager, P. (2014) ‘Fatigue and acute/chronic anaemia’, *Danish medical journal*, 61(4), p. B4824.
- Benyamin, B. *et al.* (2014) ‘Novel loci affecting iron homeostasis and their effects in individuals at risk for hemochromatosis’, *Nature Communications*, 5(May). doi: 10.1038/ncomms5926.
- Blanco-Rojo, R. *et al.* (2014) ‘Influence of diet, menstruation and genetic factors on iron status: A cross-sectional study in Spanish women of childbearing age’, *International Journal of Molecular Sciences*, 15(3), pp. 4077–4087. doi: 10.3390/ijms15034077.
- Blanco, E. *et al.* (2009) ‘Not all DMT1 mutations lead to iron overload’, *Blood Cells, Molecules, and Diseases*, 43(2), pp. 199–201. doi: 10.1016/j.bcmd.2009.05.003.
- Bossuyt, P. M. *et al.* (2006) ‘Bossuyt et al. (2006). assessing new tests against existing diagnostic pathways. BMJ-332_1089-1092’, pp. 1089–1092.
- Briissot, P. *et al.* (2012) ‘Non-transferrin bound iron: A key role in iron overload and iron toxicity’, *Biochimica et Biophysica Acta - General Subjects*, 1820(3), pp. 403–410. doi: 10.1016/j.bbagen.2011.07.014.
- Brugnara, C. (2006) ‘Reticulocyte hemoglobin equivalent (Ret He) and assessment of iron-deficient states’, *Clinical and Laboratory Haematology*, 28(5), pp. 303–308. doi: 10.1111/j.1365-2257.2006.00812.x.
- Buehler, A. M. *et al.* (2019) ‘Rational use of diagnostic tests for clinical decision making’, *Revista da Associacao Medica Brasileira*, 65(3), pp. 452–459. doi:



10.1590/1806-9282.65.3.452.

- Buttarello, M. *et al.* (2004) 'The New Reticulocyte Parameter (RET-Y) of the Sysmex XE 2100: Its Use in the Diagnosis and Monitoring of Posttreatment Sideropenic Anemia', *American Journal of Clinical Pathology*, 121(4), pp. 489–495. doi: 10.1309/W65295DTUWK7U1HH.
- Buttarello, M. *et al.* (2016) 'Evaluation of the hypochromic erythrocyte and reticulocyte hemoglobin content provided by the Sysmex XE-5000 analyzer in diagnosis of iron deficiency erythropoiesis', *Clinical Chemistry and Laboratory Medicine*, 54(12), pp. 1939–1945. doi: 10.1515/cclm-2016-0041.
- Buttarello, M. (2016) 'Laboratory diagnosis of anemia: are the old and new red cell parameters useful in classification and treatment, how?', *International Journal of Laboratory Hematology*, 38, pp. 123–132. doi: 10.1111/ijlh.12500.
- Cable, R. G. *et al.* (2016) 'Effect of iron supplementation on iron stores and total body iron after whole blood donation', *Transfusion*, 56(8), pp. 2005–2012. doi: 10.1111/trf.13659.
- Camaschella, C. (2015) 'Iron-deficiency anemia', *New England Journal of Medicine*, 372(19), pp. 1832–1843. doi: 10.1056/NEJMra1401038.
- Camaschella, C. (2017) 'New insights into iron deficiency and iron deficiency anemia', *Blood Reviews*, 31(4), pp. 225–233. doi: 10.1016/j.blre.2017.02.004.
- Camaschella, C. (2019) 'Iron deficiency', *Blood*, 133(1), pp. 30–39. doi: 10.1182/blood-2018-05-815944.
- Camaschella, C. and Nai, A. (2016) 'Ineffective erythropoiesis and regulation of iron status in iron loading anaemias', *British Journal of Haematology*, 172(4), pp. 512–523. doi: 10.1111/bjh.13820.
- Canals, C. *et al.* (2005) 'Clinical utility of the new Sysmex XE 2100 parameter - Reticulocyte hemoglobin equivalent - In the diagnosis of anemia', *Haematologica*, 90(8), pp. 1133–1134.
- Canonne-Hergaux, F. *et al.* (1999) 'Cellular and subcellular localization of the Nramp2 iron transporter in the intestinal brush border and regulation by dietary iron', *Blood*, 93(12), pp. 4406–4417. doi: 10.1182/blood.v93.12.4406.
- Canonne-Hergaux, F. *et al.* (2001) 'Characterization of the iron transporter DMT1 (NRAMP2/DCT1) in red blood cells of normal and anemic mk/mk mice', *Blood*, 98(13), pp. 3823–3830. doi: 10.1182/blood.V98.13.3823.
- Canonne-Hergaux, F. *et al.* (2006) 'Comparative studies of duodenal and macrophage ferroportin proteins', *American Journal of Physiology - Gastrointestinal and Liver Physiology*, 290(1). doi: 10.1152/ajpgi.00227.2005.
- Chaipokam, J. (2016) 'Diagnostic accuracy of reticulocyte hemoglobin content in Thai patients with microcytic red cells as a test for iron deficiency anemia', *Asian Biomedicine*, 10, pp. S31–S37. doi: 10.5372/1905-7415.1000.519.
- Cheng, H. L. *et al.* (2012) 'The relationship between obesity and hypoferaemia in adults: A systematic review', *Obesity Reviews*, 13(2), pp. 150–161. doi: 10.1111/j.1467-789X.2011.00938.x.
- Cherukuri, S. *et al.* (2005) 'Unexpected role of ceruloplasmin in intestinal iron



- absorption’, *Cell Metabolism*, 2(5), pp. 309–319. doi: 10.1016/j.cmet.2005.10.003.
- Chiabrando, D. *et al.* (2012) ‘The mitochondrial heme exporter FLVCR1b mediates erythroid differentiation’, *Journal of Clinical Investigation*, 122(12), pp. 4569–4579. doi: 10.1172/JCI62422.
- Chinudomwong, P. *et al.* (2020) ‘Diagnostic performance of reticulocyte hemoglobin equivalent in assessing the iron status’, *Journal of Clinical Laboratory Analysis*, 34(6), pp. 1–7. doi: 10.1002/jcla.23225.
- Choi, J. *et al.* (2012) ‘Duodenal reductase activity and spleen iron stores are reduced and erythropoiesis is abnormal in Dcytb knockout mice exposed to hypoxic conditions’, *Journal of Nutrition*, 142(11), pp. 1929–1934. doi: 10.3945/jn.112.160358.
- Chung, M. *et al.* (2012) ‘Biomarkers for Assessing and Managing Iron Deficiency Anemia in Late-Stage Chronic Kidney Disease: Future Research Need’, *Future Research Needs Paper*, 33(83).
- Coad, J. and Pedley, K. (2014) ‘Iron deficiency and iron deficiency anemia in women’, *Scandinavian Journal of Clinical and Laboratory Investigation*, 74(SUPPL. 244), pp. 82–89. doi: 10.3109/00365513.2014.936694.
- Cogswell, M. E. *et al.* (2009) ‘Assessment of iron deficiency in US preschool children and nonpregnant females of childbearing age: National Health and Nutrition Examination Survey 2003–2006’, *American Journal of Clinical Nutrition*, 89(5), pp. 1334–1342. doi: 10.3945/ajcn.2008.27151.
- Cohen, L. A. *et al.* (2010) ‘Serum ferritin is derived primarily from macrophages through a nonclassical secretory pathway’, *Blood*, 116(9), pp. 1574–1584. doi: 10.1182/blood-2009-11-253815.
- Constantine, C. C. *et al.* (2009) ‘A novel association between a SNP in CYBRD1 and serum ferritin levels in a cohort study of HFE hereditary haemochromatosis’, *British Journal of Haematology*, 147(1), pp. 140–149. doi: 10.1111/j.1365-2141.2009.07843.x.
- Cook, J. D. (2003) ‘The quantitative assessment of body iron’, *Blood*, 101(9), pp. 3359–3364. doi: 10.1182/blood-2002-10-3071.
- Culleton, B. F. *et al.* (2006) ‘Impact of anemia on hospitalization and mortality in older adults’, *Blood*, 107(10), pp. 3841–3846. doi: 10.1182/blood-2005-10-4308.
- Delaby, C. *et al.* (2012) ‘Subcellular localization of iron and heme metabolism related proteins at early stages of erythrophagocytosis’, *PLoS ONE*, 7(7). doi: 10.1371/journal.pone.0042199.
- Delforge, M. *et al.* (2011) ‘Iron status and treatment modalities in transfusion-dependent patients with myelodysplastic syndromes’, *Annals of Hematology*, 90(6), pp. 655–666. doi: 10.1007/s00277-011-1164-9.
- Donovan, A. *et al.* (2000) ‘Positional cloning of zebrafish ferroportin1 identifies a conserved vertebrate iron exporter’, *Nature*, 403(6771), pp. 776–781. doi: 10.1038/35001596.
- Donovan, A. *et al.* (2005) ‘The iron exporter ferroportin/Slc40a1 is essential for iron



- homeostasis', *Cell Metabolism*, 1(3), pp. 191–200. doi: 10.1016/j.cmet.2005.01.003.
- Drakesmith, H., Nemeth, E. and Ganz, T. (2015) 'Ironing out Ferroportin', *Cell Metabolism*, 22(5), pp. 777–787. doi: 10.1016/j.cmet.2015.09.006.
- Eren, H. (2005) 'Calibration Process', *Handbook of Measuring System Design*, (March 2005). doi: 10.1002/0471497398.mm999.
- de Falco, L. et al. (2014) 'Functional and clinical impact of novel Tmprss6 variants in iron-refractory iron-deficiency Anemia patients and genotype-phenotype studies', *Human Mutation*, 35(11), pp. 1321–1329. doi: 10.1002/humu.22632.
- Falkingham, M. et al. (2010) 'The effects of oral iron supplementation on cognition in older children and adults: A systematic review and meta-analysis', *Nutrition Journal*, 9(1), pp. 1–16. doi: 10.1186/1475-2891-9-4.
- Fernandez-Jimenez, M. C. et al. (2020) 'Iron Deficiency in Menstruating Adult Women: Much More than Anemia', *Women's Health Reports*, 1(1), pp. 26–35. doi: 10.1089/whr.2019.0011.
- Ferrari, M. et al. (2011) 'Evaluation of iron status in European adolescents through biochemical iron indicators: The HELENA Study', *European Journal of Clinical Nutrition*, 65(3), pp. 340–349. doi: 10.1038/ejcn.2010.279.
- Finberg, K. E. et al. (2008) 'Mutations in TMPRSS6 cause iron-refractory iron deficiency anemia (IRIDA)', *Nature Genetics*, 40(5), pp. 569–571. doi: 10.1038/ng.130.
- Fischl, M.A., Richman, D. (1987) 'The New England Journal of Medicine Downloaded from nejm.org at UNIVERSITY OF AUCKLAND on September 15, 2013. For personal use only. No other uses without permission. From the NEJM Archive. Copyright © 2010 Massachusetts Medical Society. All rights reserv', *The New England Journal of Medicine*, pp. 2010–2013.
- Fleming, M. D. et al. (1998) 'Nramp2 is mutated in the anemic Belgrade (b) rat: Evidence of a role for Nramp2 in endosomal iron transport', *Proceedings of the National Academy of Sciences of the United States of America*, 95(3), pp. 1148–1153. doi: 10.1073/pnas.95.3.1148.
- Folgueras, A. R. et al. (2008) 'Membrane-bound serine protease matriptase-2 (Tmprss6) is an essential regulator of iron homeostasis', *Blood*, 112(6), pp. 2539–2545. doi: 10.1182/blood-2008-04-149773.
- Ganz, T. (2013) 'Systemic iron homeostasis', *Physiological Reviews*, 93(4), pp. 1721–1741. doi: 10.1152/physrev.00008.2013.
- Gkouvatsos, K. (2012) 'Regulation of iron transport and the role of transferrin', *Biochimica et Biophysica Acta - General Subjects*, 1820(3), pp. 188–202. doi: 10.1016/j.bbagen.2011.10.013.
- Gottlieb, Y. et al. (2012) 'Physiologically aged red blood cells undergo erythrophagocytosis in vivo but not in vitro', *Haematologica*, 97(7), pp. 994–1002. doi: 10.3324/haematol.2011.057620.
- Harvey, L. J. et al. (2005) 'Impact of menstrual blood loss and diet on iron deficiency among women in the UK', *British Journal of Nutrition*, 94(4), pp. 557–564. doi:



10.1079/bjn20051493.

- Heath, A.-L. M. *et al.* (2001) ‘The role of blood loss and diet in the aetiology of mild iron deficiency in premenopausal adult New Zealand women’, *Public Health Nutrition*, 4(2), pp. 197–206. doi: 10.1079/phn200054.
- Hudson, B. H. *et al.* (2018) ‘Modulation of intestinal sulfur assimilation metabolism regulates iron homeostasis’, *Proceedings of the National Academy of Sciences of the United States of America*, 115(12), pp. 3000–3005. doi: 10.1073/pnas.1715302115.
- Illing, A. C. *et al.* (2012) ‘Substrate profile and metal-ion selectivity of human divalent metal-ion transporter-1’, *Journal of Biological Chemistry*, 287(36), pp. 30485–30496. doi: 10.1074/jbc.M112.364208.
- Iron, F. and Requires, R. (2006) ‘involved . Following a proposal of Roberts and Bomford (31), Konijn and colleagues (46) reported that inhibitors of lysosomal proteolysis (leupeptin , chymostatin , and chloroquine) re- duced the rate of iron accumulation in hemoglobin in primary ’, 291(September).
- Jakobsen, T. H. *et al.* (2013) ‘Complete Genome Sequence of the Cystic Fibrosis Pathogen *Achromobacter xylosoxidans* NH44784-1996 Complies with Important Pathogenic Phenotypes’, *PLoS ONE*, 8(7), pp. 8–11. doi: 10.1371/journal.pone.0068484.
- Kalantar-Zadeh, K. *et al.* (2005) ‘Time-dependent associations between iron and mortality in hemodialysis patients’, *Journal of the American Society of Nephrology*, 16(10), pp. 3070–3080. doi: 10.1681/ASN.2005040423.
- Karlsson, T. (2011) ‘Comparative evaluation of the reticulocyte hemoglobin content assay when screening for iron deficiency in elderly anemic patients’, *Anemia*, 2011. doi: 10.1155/2011/925907.
- Kassebaum, N. J. *et al.* (2015) ‘Plenary Paper RED CELLS , IRON , AND ERYTHROPOIESIS A systematic analysis of global anemia burden from 1990 to 2010’, *Blood Journal*, 123(5), pp. 615–625. doi: 10.1182/blood-2013-06-508325.The.
- Keel, S. B. *et al.* (2008) ‘Corrected 25 july 2008; see last page’, *Science*, 306(July), pp. 825–829.
- Khan, N. *et al.* (2019) ‘Diagnostic Accuracy of Reticulocyte Haemoglobin Equivalent (RET-He) in Detecting Iron Deficiency Anaemia Keeping Serum Feritin as Gold Standart’, 69(5), pp. 1010–1014.
- Kiss, J. E. and Vassallo, R. R. (2018) ‘How do we manage iron deficiency after blood donation?’, *British Journal of Haematology*, 181(5), pp. 590–603. doi: 10.1111/bjh.15136.
- Kiudeliene, R., Griniute, R. and Labanauskas, L. (2008) ‘Prognostic value of reticulocyte hemoglobin content to diagnose iron deficiency in 6-24-month-old children’, *Medicina*, 44(9), pp. 673–677. doi: 10.3390/medicina44090086.
- Knovich, M. A. *et al.* (2009) ‘Feritin for the clinician’, *Blood Reviews*, 23(3), pp. 95–104. doi: 10.1016/j.blre.2008.08.001.
- Kosman, D. J. (2010) ‘Redox cycling in iron uptake, efflux, and trafficking’, *Journal*



- of *Biological Chemistry*, 285(35), pp. 26729–26735. doi: 10.1074/jbc.R110.113217.
- Labbé, R. F. (1999) ‘Zinc protoporphyrin: A metabolite with a mission’, *Clinical Chemistry*, 45(12), pp. 2060–2072. doi: 10.1093/clinchem/45.12.2060.
- Lam, L. F. and Lawlis, T. R. (2017) ‘Feeding the brain – The effects of micronutrient interventions on cognitive performance among school-aged children: A systematic review of randomized controlled trials’, *Clinical Nutrition*, 36(4), pp. 1007–1014. doi: 10.1016/j.clnu.2016.06.013.
- Lee, T. et al. (2016) ‘Oral versus intravenous iron replacement therapy distinctly alters the gut microbiota and metabolome in patients with IBD’, *Gut*, 66(5), pp. 863–871. doi: 10.1136/gutjnl-2015-309940.
- Lian, Y. et al. (2019) ‘Reticulocyte Hemoglobin Equivalent (RET-He) Combined with Red Blood Cell Distribution Width Has a Differentially Diagnostic Value for Thalassemias’, *Hemoglobin*, 43(4–5), pp. 229–235. doi: 10.1080/03630269.2019.1655440.
- Lopez, A. et al. (2016) ‘Iron deficiency anaemia’, *The Lancet*, 387(10021), pp. 907–916. doi: 10.1016/S0140-6736(15)60865-0.
- Ma, Y. et al. (2006) ‘Iron Imports. V. Transport of iron through the intestinal epithelium’, *American Journal of Physiology - Gastrointestinal and Liver Physiology*, 290(3), pp. 417–422. doi: 10.1152/ajpgi.00489.2005.
- Macapagal et al., 2017 (2016) ‘乳鼠心肌提取 HHS Public Access’, *Physiology & behavior*, 176(1), pp. 139–148. doi: 10.1016/j.physbeh.2017.03.040.
- Macdougall, I. C. et al. (2016) ‘Iron management in chronic kidney disease: Conclusions from a “kidney Disease: Improving Global Outcomes” (KDIGO) Controversies Conference’, *Kidney International*, 89(1), pp. 28–39. doi: 10.1016/j.kint.2015.10.002.
- Mast, A. E. et al. (2011) ‘Clinical utility of the reticulocyte hemoglobin content in the diagnosis of iron deficiency Brief report Clinical utility of the reticulocyte hemoglobin content in the diagnosis of iron deficiency’, *Bone*, 99(4), pp. 1489–1491. doi: 10.1182/blood.V99.4.1489.
- Mattiello, V. et al. (2020) ‘Diagnosis and management of iron deficiency in children with or without anemia: consensus recommendations of the SPOG Pediatric Hematology Working Group’, *European Journal of Pediatrics*, MC. doi: 10.1007/s00431-020-03597-5.
- Maxim, L. D., Niebo, R. and Utell, M. J. (2014) ‘Screening tests: A review with examples’, *Inhalation Toxicology*, 26(13), pp. 811–828. doi: 10.3109/08958378.2014.955932.
- McCann, J. C. and Ames, B. N. (2007) ‘An overview of evidence for a causal relation between iron deficiency during development and deficits in cognitive or behavioral function’, *American Journal of Clinical Nutrition*, 85(4), pp. 931–945. doi: 10.1093/ajcn/85.4.931.
- McKie, A. T. (2008) ‘The role of Dcytb in iron metabolism: An update’, *Biochemical*



- Society Transactions*, 36(6), pp. 1239–1241. doi: 10.1042/BST0361239.
- McLean, E. et al. (2009) ‘Worldwide prevalence of anaemia, WHO Vitamin and Mineral Nutrition Information System, 1993-2005’, *Public Health Nutrition*, 12(4), pp. 444–454. doi: 10.1017/S1368980008002401.
- Mehta, S. et al. (2016) ‘Reticulocyte hemoglobin vis-a-vis serum ferritin as a marker of bone marrow iron store in iron deficiency anemia’, *Journal of Association of Physicians of India*, 64(NOVEMBER), pp. 38–42.
- Meyron-Holtz, E. G. (2011) ‘A possible role for secreted ferritin in tissue iron distribution.’, *Journal of neural transmission (Vienna, Austria : 1996)*, 118(3), pp. 337–347. doi: 10.1007/s00702-011-0582-0.
- Mirza, F. G. et al. (2018) ‘Impact and management of iron deficiency and iron deficiency anemia in women’s health’, *Expert Review of Hematology*, 11(9), pp. 727–736. doi: 10.1080/17474086.2018.1502081.
- Mitchell, C. J. et al. (2014) ‘Functional properties of human ferroportin, a cellular iron exporter reactive also with cobalt and zinc’, *American Journal of Physiology - Cell Physiology*, 306(5), pp. 450–459. doi: 10.1152/ajpcell.00348.2013.
- de Montalembert, M. et al. (2012) ‘Exploration d’une anémie microcytaire chez l’enfant’, *Archives de Pédiatrie*, 19(3), pp. 295–304. doi: 10.1016/j.arcped.2011.12.016.
- Musallam, K. M. and Taher, A. T. (2018) ‘Iron deficiency beyond erythropoiesis: should we be concerned?’, *Current Medical Research and Opinion*, 34(1), pp. 81–93. doi: 10.1080/03007995.2017.1394833.
- Nead, K. G. et al. (2004) ‘Overweight children and adolescents: A risk group for iron deficiency’, *Pediatrics*, 114(1), pp. 104–108. doi: 10.1542/peds.114.1.104.
- Origa, R. et al. (2007) ‘Liver iron concentrations and urinary hepcidin in β-thalassemia’, *Haematologica*, 92(5), pp. 583–588. doi: 10.3324/haematol.10842.
- Peyrin-Biroulet, L. (2015) ‘Guidelines on the diagnosis and treatment of iron deficiency across indications: A systematic review’, *American Journal of Clinical Nutrition*, 102(6), pp. 1585–1594. doi: 10.3945/ajcn.114.103366.
- Potter, Y. (2013) ‘Iron deficiency anaemia.’, *Nursing standard (Royal College of Nursing (Great Britain) : 1987)*, 27(23), p. 59. doi: 10.7748/ns2013.02.27.23.59.p10441.
- Qiu, A. et al. (2006) ‘Identification of an Intestinal Folate Transporter and the Molecular Basis for Hereditary Folate Malabsorption’, *Cell*, 127(5), pp. 917–928. doi: 10.1016/j.cell.2006.09.041.
- Rahayu, E. T. (2014) ‘Peran Reticulocyte Hemoglobin Equivalent (RET-He) untuk Skrining Defisiensi Besi pada Anak Usia 6 Bulan- 5 Tahun’, *Sysmex*.
- Rajagopal, A. et al. (2008) ‘Haem homeostasis is regulated by the conserved and concerted functions of HRG-1 proteins’, *Nature*, 453(7198), pp. 1127–1131. doi: 10.1038/nature06934.
- Ramey, G. et al. (2010) ‘Hepcidin targets ferroportin for degradation in hepatocytes’, *Haematologica*, 95(3), pp. 501–504. doi: 10.3324/haematol.2009.014399.
- Rowland, T. W. et al. (1988) ‘The Effect of Iron Therapy on the Exercise Capacity of



- Nonanemic Iron-Deficient Adolescent Runners', *American Journal of Diseases of Children*, 142(2), pp. 165–169. doi: 10.1001/archpedi.1988.02150020067030.
- Russel, R. et al. (2002) *Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc*. National Academy Press Washington DC.
- Rybo, L. H. A. H. L. N. G. (1966) 'Menstrual Blood Loss-a', (1945).
- Schneider, J., et al (2005) 'Children From Low-Income Families 1 – 3', *The American Journal of Clinical Nutrition*, (Id), pp. 1269–1275.
- Shawki, A. et al. (2012) *H + -Coupled Divalent Metal-Ion Transporter-1: Functional Properties, Physiological Roles and Therapeutics, Current Topics in Membranes*. Elsevier. doi: 10.1016/B978-0-12-394316-3.00005-3.
- Shawki, A. et al. (2016) 'Intestinal brush-border Na+/H⁺ exchanger-3 drives H⁺-coupled iron absorption in the mouse', *American Journal of Physiology - Gastrointestinal and Liver Physiology*, 311(3), pp. G423–G430. doi: 10.1152/ajpgi.00167.2016.
- Shayeghi, M. et al. (2005) 'Identification of an intestinal heme transporter', *Cell*, 122(5), pp. 789–801. doi: 10.1016/j.cell.2005.06.025.
- Shi, H. et al. (2008) 'A cytosolic iron chaperone that delivers iron to ferritin', *Science*, 320(5880), pp. 1207–1210. doi: 10.1126/science.1157643.
- Siswosudarmo, R. (2017) 'Tes diagnostik (Diagnostic test)', *Jurnal Metodologi penelitian*. Available at: <http://obgin-ugm.com/wp-content/uploads/2017/09/HRS-Kuliah-Tes-Diagnostik.pdf>.
- Soe-Lin, S. et al. (2009) 'Nramp1 promotes efficient macrophage recycling of iron following erythrophagocytosis in vivo', *Proceedings of the National Academy of Sciences of the United States of America*, 106(14), pp. 5960–5965. doi: 10.1073/pnas.0900808106.
- Sørensen, E. et al. (2016) 'Genetic factors influencing ferritin levels in 14,126 blood donors: Results from the Danish Blood Donor Study', *Transfusion*, 56(3), pp. 622–627. doi: 10.1111/trf.13397.
- Stauder, R. (2018) 'Anemia at older age: etiologies, clinical implications, and management', *Blood*, 131(5), pp. 505–514. doi: 10.1182/blood-2017-07-746446.
- Steenackers, N. et al. (2018) 'Iron deficiency after bariatric surgery: What is the real problem?', *Proceedings of the Nutrition Society*, 77(4), pp. 445–455. doi: 10.1017/S0029665118000149.
- Steinbicker, A. U. and Muckenthaler, M. U. (2013) 'Out of balance-systemic iron homeostasis in iron-related disorders', *Nutrients*, 5(8), pp. 3034–3061. doi: 10.3390/nu5083034.
- Stevens, G. A. et al. (2013) 'Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: A systematic analysis of population-representative data', *The Lancet Global Health*, 1(1), pp. 16–25. doi: 10.1016/S2214-109X(13)70001-9.
- Stoffman, N. (2005) 'An algorithm using reticulocyte hemoglobin content (CHr)



- measurement in screening adolescents for iron deficiency', *Journal of Adolescent Health*, 36(6), pp. 1–6. doi: 10.1016/j.jadohealth.2004.09.011.
- Sudmann, Å. A. (2012) 'Reticulocyte hemoglobin equivalent to detect thalassemia and thalassemic hemoglobin variants', *International Journal of Laboratory Hematology*, 34(6), pp. 605–613. doi: 10.1111/j.1751-553X.2012.01442.x.
- Susanne, F. (2004) 'Potential Utility of Ret-Y in the Diagnosis of Iron- Restricted Erythropoiesis', *Clinical Chemistry*, 50(7), pp. 1237–1240. doi: 10.1373/clinchem.2004.031237.
- Terrier, B. et al. (2012) 'Prevalence, characteristics and prognostic significance of anemia in daily practice', *Qjm*, 105(4), pp. 345–354. doi: 10.1093/qjmed/hcr230.
- Theil, E. C. et al. (2012) 'Absorption of iron from ferritin is independent of heme iron and ferrous salts in women and rat intestinal segments', *Journal of Nutrition*, 142(3), pp. 478–483. doi: 10.3945/jn.111.145854.
- Thuret, I. (2017) 'Diagnostic biologique de la carence martiale chez l'enfant', *Archives de Pediatrie*, 24(5), pp. S56-S13. doi: 10.1016/S0929-693X(17)24003-2.
- Tim Riskesdas (2018) 'Laporan Nasional RISKESDAS 2018.pdf', *Badan Penelitian dan Pengembangan Kesehatan*, p. 198. Available at: http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf.
- Tolosano, E. et al. (2002) 'Enhanced splenomegaly and severe liver inflammation in haptoglobin/hemopexin double-null mice after acute hemolysis', *Blood*, 100(12), pp. 4201–4208. doi: 10.1182/blood-2002-04-1270.
- Torino, A. B. B. et al. (2015) 'Evaluation of erythrocyte and reticulocyte parameters as indicative of iron deficiency in patients with anemia of chronic disease', *Revista Brasileira de Hematologia e Hemoterapia*, 37(2), pp. 77–81. doi: 10.1016/j.bjhh.2015.02.004.
- Trevethan, R. (2017) 'Sensitivity, Specificity, and Predictive Values: Foundations, Pliabilities, and Pitfalls in Research and Practice', *Frontiers in Public Health*, 5(November), pp. 1–7. doi: 10.3389/fpubh.2017.00307.
- Trost, L. B. (2006) 'The diagnosis and treatment of iron deficiency and its potential relationship to hair loss', *Journal of the American Academy of Dermatology*, 54(5), pp. 824–844. doi: 10.1016/j.jaad.2005.11.1104.
- Uijterschout, L. et al. (2014) 'The value of Ret-Hb and sTfR in the diagnosis of iron depletion in healthy, young children', *European Journal of Clinical Nutrition*, 68(8), pp. 882–886. doi: 10.1038/ejcn.2014.70.
- Ullrich, C. et al. (2005) 'Using Reticulocyte Hemoglobin Content', 294(8), pp. 924–930.
- Urrechaga, E. (2013) 'Erythrocyte and reticulocyte indices in the assessment of erythropoiesis activity and iron availability', *International Journal of Laboratory Hematology*, 35(2), pp. 144–149. doi: 10.1111/ijlh.12013.
- Urrechaga, E. et al. (2014) 'Looking Back To Our Roots : 80 Years of Wintrobe ' S Indices', *European Medical Journal*, (July), pp. 7–9.
- Urrechaga Igartua, E. et al. (2017) 'Reticulocyte hemoglobin content (MCHr) in the



- detection of iron deficiency', *Journal of Trace Elements in Medicine and Biology*, 43, pp. 29–32. doi: 10.1016/j.jtemb.2016.11.001.
- Vanoaica, L. et al. (2010) 'Intestinal feritin H is required for an accurate control of iron absorption', *Cell Metabolism*, 12(3), pp. 273–282. doi: 10.1016/j.cmet.2010.08.003.
- Vázquez-López, M. A. et al. (2019) 'The usefulness of reticulocyte haemoglobin content, serum transferrin receptor and the sTfR-feritin index to identify iron deficiency in healthy children aged 1–16 years', *European Journal of Pediatrics*, 178(1), pp. 41–49. doi: 10.1007/s00431-018-3257-0.
- Vos, T. et al. (2017) 'Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: A systematic analysis for the Global Burden of Disease Study 2016', *The Lancet*, 390(10100), pp. 1211–1259. doi: 10.1016/S0140-6736(17)32154-2.
- Wang, H. et al. (2004) 'Detection and monitoring of SARS coronavirus in the plasma and peripheral blood lymphocytes of patients with severe acute respiratory syndrome', *Clinical Chemistry*, 50(7), pp. 1237–1240. doi: 10.1373/clinchem.2004.031237.
- WHO (2011) 'Serum feritin concentrations for the assessment of iron status and iron deficiency in populations. Vitamin and Mineral Nutrition Information System.', WHO, pp. 1–5. doi: (WHO/NMH/NHD/MNM/11.2).
- World Health Organization (2011) 'Prevention of iron deficiency anaemia in adolescents', *Role of Weekly Iron and Folic Acid Supplementation*, p. 50.
- World Health Organization (2016) 'Guideline Daily Iron', *Daily Iron Supplementation in infants and children*, p. 44.
- Worwood, M. (2017) *Iron Deficiency Anaemia and Iron Overload*. Twelfth Ed, *Dacie and Lewis Practical Haematology: Twelfth Edition*. Twelfth Ed. Elsevier Ltd. doi: 10.1016/B978-0-7020-6696-2.00009-6.
- Yanoff, L. B. et al. (2007) 'Inflammation and iron deficiency in the hypoferremia of obesity', *International Journal of Obesity*, 31(9), pp. 1412–1419. doi: 10.1038/sj.ijo.0803625.
- Zhang, D. L. et al. (2009) 'A Ferroportin Transcript that Lacks an Iron-Responsive Element Enables Duodenal and Erythroid Precursor Cells to Evade Translational Repression', *Cell Metabolism*, 9(5), pp. 461–473. doi: 10.1016/j.cmet.2009.03.006.