

DAFTAR PUSTAKA

- Ahn JH, Hong IP, Bok JI, Kim BY, Song J, Weon HY. 2012. Pyrosequencing analysis of the bacterial communities in the guts of honey bees *Apis cerana* and *Apis mellifera* in Korea. *Journal of Microbiology*. 50: 735-45.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *The Journal of Molecular Biology*. 215: 403–410.
- Atwal AS, Dhaliwal GS. 1969. Some behavioural characteristics of *Apis indica* F and *Apis mellifera* L. *Indian Bee*. 31: 83-90
- Buttel-Reepen H. 1906. Apistica Beitrage zur Systematik Biologie sowie zur geschichtlichen und geographischen Verbreitung der Honigbiene (*Apis mellifica* L) ihrer Varietaten und der iibrigen Apis-Arten. *Veroff Zool Mus Berlin*. 118-120
- Caporaso JG, Bittinger K, Bushman FD, Desantis TZ, Andersen GL, Knight R. 2010. PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics*. 26: 266–267.
- Corby-Harris V, Maes P, Anderson KE. 2014. The bacterial communities associated with honey bee (*Apis mellifera*) foragers. *PLoS ONE*. 9:1-13.
- Damus, M.S. and G.W. Otis. 1997. A morphometric analysis of *Apis cerana*F and *Apis nigrocincta*Smith populations from Southeast Asia. *Apidologie*. 28: 309-323.
- De Graaf DC, Alippi AM, Antúnez K, Aronstein KA, Budge G De, Koker D De, Smet L, Dingman DW, Evans JD, Foster LJ, Fünfhaus A, Garcia-Gonzalez E, Gregorc A, Human H, Murray KD, Nguyen BK, Poppinga L, Spivak M, Vanengelsdorp D, Wilkins S, Genersch E. 2013. Standard methods for American foulbrood research. standard methods for *Apis mellifera* pest and pathogen research. *Journal of Apicultural Research*. 1: 52
- Degewione A, Alamerew S, and Tabor G. 2011. Genetic variability and association of bulb yield and related traits in shallot (*Allium cepa* var *aggregatum* DON.) in Ethiopia. *Inter. J. Agri*. 21: 1-20.
- De Miranda JR, Bailey L, Ball BV, Blanchard P, Budge G, Chejanovsky N, Chen Y-P, Gauthier L, Genersch E, De Graaf D, Ribière M, Ryabov E, De Smet L, Van Der Steen JJ M. 2013. Standard methods for virus research in *Apis*

- mellifera*. Standard methods for *Apis mellifera* pest and pathogen research. *Journal of Apicultural Research*. 4: 52
- Dhaliwal GS, Atwal AS. 1970. Interspecific relations between *Apis cerana indica* and *Apis mellifera*. *J Apic Res*. 9: 53-59
- Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*. 26: 2460–2461.
- Engel MS. 2012. The honey bees of Indonesia (Hymenoptera: Apidae). *Treubia*. 39:1-85
- Engel, P. and Moran, N.A. 2013. The gut microbiota of insects - diversity in structure and function. *FEMS Microbiology Reviews*. 37, 699–735
- Engelbrektson A, Kunin V, Wrighton KC, Zvenigorodsky N, Chen F, Ochman H, Hugenholtz P. 2010. Experimental factors affecting PCR-based estimates of microbial species richness and evenness. *The ISME Journal*. 4: 642–647.
- Excoffier L, Smouse PE, Quattro JM. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics*. 131: 479–491.
- Francoy, TM. 2009. Gender Identification of Five Genera of Stingless Bees (Apidae, Meliponini) based on Wing Morphology. *Genet Mol Res*. 8: 207-214.
- Free, JB. 1982. *Bees and Mankind*. London (GB): George Allen & Unwin.
- Fries I, Chauzat MP, Chen YP, Doublet V, Genersch E, Gisder S, Higes M, McMahon DP, Martín Hernández R, Natsopoulou M, Paxton RJ, Tanner G, Webster TC, Williams, GR. 2013. Standard methods for *Nosema* research. Standard methods for *Apis mellifera* pest and pathogen research. *Journal of Apicultural Research*. 1: 52
- Gotelli N J, Colwell R K. 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters*. 4: 379–391.
- Hadisoesilo, S. and G.W. Otis. 1996. Drone flight times confirm the species status of *Apis nigrocincta* Smith, 1861 to be a species distinct from *Apis cerana* F. In Sulawesi, Indonesia. *Apidologie*. 27: 361-369.
- Hadisoesilo, S. and G.W. Otis. 1998. Differences in drone cappings of *Apis cerana* and *Apis nigrocincta*. *J. Apic. Res*. 37: 11-15.

- Hadisoesilo S. 2001. Keanekaragaman spesies lebah madu asli Indonesia pusat penelitian dan pengembangan hutan dan konservasi alam, Bogor. *Biodiversitas*. 2: 123-128.
- Handelsman, J. 2004. Metagenomics: application of genomics to uncultured microorganisms. *Microbiology and Molecular Biology Reviews*. 68: 669-685.
- Hebert, P.D.N., Cywinska, A., and Ball, S.L. 2003. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*. 270(151): 313-321.
- Jensen AB, Aronstein K, Flores JM, Vojvodic S, Palacio MA, Spivak M. 2013. Standard methods for fungal brood disease research. Standard methods for *Apis mellifera* pest and pathogen research. *Journal of Apicultural Research*. 1: 52
- Kumar MS, Singh AJAR, Alagumuthu G. 2012. Traditional beeskeeping of stingless bees (*Trigona* sp.) by kani tribes of Western Ghats, Tamil Nadu, India. *Indian J Tradit Knowledge*. 11: 342-345.
- Lindauer M. 1956. Dber die Verstlindigung bei indischen Bienen. *Z vgl Physiol*. 38: 521-55
- Lourino MC, Fonseca VLI, Roubik DW, Dollin A, Heard T, Aguilar IB, Venturieri GC, Eardley C, Neto PN. 2006. Global meliponiculture: challenges and opportunities. *Apidologie*. 37: 1-18.
- Maa, T.C. 1953. An inquiry inti the systematics of the tribus Apidini or honeybees (Hymenoptera). *Treubia*. 21: 525-640.
- Mahendran, B, Ghosh SK, Kundu SC. 2006. Molecular phylogeny of silk-producing insects based on 16S ribosomal RNA and cytochrome oxidase subunit I genes. *Journal of Genetics*. 85: 31-38.
- Martinson VG, Moy J, Moran NA. 2012. Establishment of characteristic gut bacteria during development of the honey bee worker. *Applied and Environmental Microbiology*. 78: 2830–2840.
- Mattila HR, Rios D, Walker-Sperling VE, Roeselers G, Newton IL. 2012. Characterization of the active microbiotas associated with honey bees reveals healthier and broader communities when colonies are genetically diverse. *PLoS One*. 7: 32-42.

- Mcfrederick QS, Wcislo WT, Taylor DR, Ishak HD, Dowd SE, Mueller UG. 2012. Environment or kin: whence do bees obtain acidophilic bacteria? *Molecular Ecology*. 21: 1754-1768.
- Menke S, Gillingham MAF, Wilhelm K, Sommer S. 2017. Hpme-Made cost Effective preservation buffer is a better alternative to commercial preservation methods for microbiome research. *Fronties in Microbiology*. 8:1-12.
- Michener CD. 2007. *The Beess of The World*. Second Edition. Baltimore (US): The Johns Hopkins Univ. Pr.
- Mishra RC, Dogra GS. 1983. Post-embryonic development of *Apis cerana indica* F worker bee.
- Moran NA, Hansen AK, Powell JE, Sabree ZL. 2012. Distinctive gut microbiota of honey bees assessed using deep sampling from individual worker bees. *PLoS One*. 7: 36-39.
- Muttoo RN. 1956. Facts about beekeeping in India. *Bee World*. 37: 125-133; 154-157.
- Na-nokorn U, Sripairoj K, Sukmanomon S, Poompuang S, Kamonrat W. 2006. Polymorphic microsatellite from DNA of the endangered Mekong giant catfish, *Pangasianodon gigas* (Chevey) and cross-species amplification in three species of *Pangasius*. *Mol. Ecol*. 6: 1174-1176.
- Newton IL, Roeselers G. 2012. The effect of training set on the classification of honey bee gut microbiota using the Naïve Bayesian Classifier. *BMC Microbiology*. 12: 221-225.
- Okada I, Sakai T. 1960. A comparative study on natural comb of Japanese and European honeybee, with special difference in cell number. *Bull Fac Agr Tamagawa Univ*. 1: 1-11
- Olofsson CT, Butler E, Markowicz P, Lindholm C, Larsson L, Vásquez A. 2016. Lactic acid bacterial symbionts in honeybees – an unknown key to honey's antimicrobial and therapeutic activities. *Int Wound J*. 13: 668–679.
- Otis, G. W. 1996. Distributions of recently recognized species of honey bees (*Apis* spp.) in Asia. *J. Kansas Entomol. Soc. Supp*. 69: 311-333.
- Rahman KA. 1945. Progress of beekeeping in the Punjab. *Bee World*. 26: 42-44; 50-52.
- Roubik DW. 1989. Stingless bee nesting biology. *Apidologie*. 37: 124-143.

- Ruttner, F. 1988. *Biogeography and Taxonomy of Honeybees*. Springer-Verlag, Berlin.
- Sakagami, F.S., T. Matsumura, K. Ito. 1980. *Apis laboriosa* Himalaya, the little known world largest honeybee (Hymenoptera, Apidae). *Insect Matsumurana*. 19: 47-77.
- Salmah, S., T. Inoue, dan S.F. Sakagami. 1990. An analysis of apid bee richness (Apidae) in Central Sumatra. Dalam Sakagami, S.F., R. Ogushi, dan D.W. Roubik (eds.), *Natural History of Social Wasps and Bees in Equatorial Sumatra*. 139-174.
- Seeley TD, Seeley RH, Akrotanakul P. 1982. Colony defense strategies of the honeybee in Thailand. *Ecol Monogr*. 52: 43-63
- Smith F. 1858. *Catalogue of the Hymenopterous insects collected at Sarawak*. Borneo: Mount
- Smith FG. 1961. Races of honeybees in East Africa. *Bee World*. 42: 255-260
- Smith DR, Hagen RH. 1996. The biogeography of *Apis cerana* as revealed by mitochondrial DNA sequence data. *Journal of the Kansas Entomological Society*. 69: 294-310.
- Sokal RR Michener C. 1958. A statistical method for evaluating systematic relationships. *The University of Kansas Science Bulletin*. 38: 1409–1438.
- Supriatna J, Wahyono EH. 2000. *Panduan Lapangan Primata Indonesia*. Jakarta: Yayasan Obor Indonesia.
- Thummajitsakul S, Silprasit K, Klinbunga S, Sittipraneed S. 2013. The partial mitochondrial sequence of the Old World stingless bee, *Tetragonula pagdeni*. *Journal of Genetics*. 92: 1-5.
- Trianto, M., & Purwanto, H. 2020. Molecular Phylogeny of Stingless Bees in the Special Region of Yogyakarta Revealed Using Partial 16S rRNA Mitochondrial Gene. *Buletin Peternakan*. 44(4): 186–93.
- Wen Y, Wang L, Jin Y, Zhang J, Su L, Zhang X, Zhou J, Li Y. 2017. The microbial community dynamics during the vitex honey ripening process in the honeycomb. *Front Microbiol*. 8: 1-12.