

## DAFTAR PUSTAKA

- Albert, P.R. (2015). Why is depression more prevalent in women? *J Psychiatry Neurosci*, 40(4):219–221.
- Amaltinga, A.P.M. dan Mbinta, J.F. (2020). Factors associated with depression among young people globally: a narrative review. *International Journal of Community Medicine and Public Health*, 7(9): 3711–3721.
- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*. Arlington, VA: American Psychiatric Association.
- Andresen, E.M., Malmgren, J.A., Carter, W.B., dan Patrick, D.L. (1994). Screening for Depression in Well Older Adults: Evaluation of a Short Form of the CES-D. *Am J Prev Med*, 10(2):77–84.
- Angraini, D.I. (2014). Hubungan Depresi dengan Status Gizi. *Medula*, 2(2):39–46.
- Appelhans, B.M., Whited, M.C., Schneider, K.L., Ma, Y., Oleski, J.L., Merriam, P.A., Waring, M.E., Olendzki, B.C., Mann, D.M., Ockene, I.S., dan Pagoto, S.L. (2012). Depression severity, diet quality, and physical activity in women with obesity and depression. *J Acad Nutr Diet*, 12(5):693–698.
- Arjadi, R., Nauta, M.H., Utoyo, D.B., dan Bockting, C.L.H. (2017). The Inventory of Depressive Symptomatology Self Report (IDS-SR): Psychometric properties of the Indonesian version. *PLoS ONE*, 12(10): e0187009.
- Assefa, T., Seid, O., Tadese, F., dan Gebremedhin, T. (2020). Nutritional Status and Associated Factors among Adult Psychiatric Patients in Dessie Referral Hospital, Northeast Ethiopia. *Psychiatry Journal*.  
<https://doi.org/10.1155/2020/5087573>
- Ball, K. dan Crawford, D. (2005). Socioeconomic status and weight change in adults: a review. *Social science & medicine*, 60:1987–2010.

- Bazire, S. (2016). *Psychotropic Drug Directory*. Lloyd-Reinhold Publications.
- Beccuti, G. dan Pannain, S. (2011). Sleep and obesity. *Current Opinion in Clinical Nutrition and Metabolic Care*, 14(4):402–412.
- Blüher, M. (2019). Obesity: global epidemiology and pathogenesis. *Nature Reviews Endocrinology*. doi:10.1038/s41574-019-0176-8.
- Budiarto, E. (2015). *Biostatistika untuk Kedokteran dan Kesehatan Masyarakat*. Jakarta: EGC.
- Buscemi, J., Pugach, O., Springfield, S., Jang, J., Tussing-Humphreys, L., Schiffer, L., Stolley, M.R., dan Fitzgibbon, M.L. (2018). Associations between fiber intake and Body Mass Index (BMI) among African-American women participating in a randomized weight loss and maintenance trial. *Eating Behaviors*, 29:48–53.
- Casadei, K. dan Kiel, J. (2020). Anthropometric Measurement. Dalam: *StatPearls*. [Internet]. Treasure Island, Florida: StatPearls Publishing. Dapat diakses di: <https://www.ncbi.nlm.nih.gov/books/NBK537315/>
- Center for Disease Control and Prevention (CDC). (2007). *National Health and Nutrition Examination Survey Anthropometric Procedures Manual*. Hyattsville: CDC.
- Christensen, H., Form, A.F., Mackinnon, A.J., Korten, A.E., Jacomb, P.A., Henderson, A.S., dan Rodgers, B. (1999). Age differences in depression and anxiety symptoms: a structural equation modelling analysis of data from a general population sample. *Psychological Medicine*, 29:325–339.
- Dagnino, P., Ugarte, M.J., Morales, F., González, S., Saralegui, D., dan Ehrental, J.C. (2020). Risk Factors for Adult Depression: Adverse Childhood Experiences and Personality Functioning. *Frontiers in Psychology*, 11:594698.

- de Wit, L.M., van Straten, A., van Hertem, m., Penninx, B.W.J.H., dan Cuijpers, P. (2009). Depression and body mass index, a u-shaped association. *BMC Public Health*, 9(14).
- Faramarzi, E., Ostadrahimi, A., Nikniaz, Z., Jafarabadi, M.A., Fakhari, A., dan Somi, M. (2018). Determination of the Best Anthropometric Index of Obesity for Prediction of Prehypertension and Hypertension in a Large Population-Based-Study; the Azar-Cohort, *Iran Red Crescent Med J*, 20(3):359911.
- Fardet, L. dan Fève, B. (2014). Systemic Glucocorticoid Therapy: a Review of its Metabolic and Cardiovascular Adverse Events. *Drugs*, 74:1731–1745.
- Flórez K.R., Dubowitz, T., Ghosh-Dastidar, M.B., Beckman, R., dan Collins, R.L. (2015). Associations between Depressive Symptomatology, Diet, and Body Mass Index among Participants in the Supplemental Nutrition Assistance Program. *Journal of the Academy of Nutrition and Dietetics*, 115(7).
- Ford, E.S., Mokdad, A.H., dan Giles, W.H. (2003). Trends in waist circumference among U.S. adults. *Obesity Research*, 11(10):1223–1231.
- Forman-Hoffman, V.L., Yankey, J.W., Hillis, S.L., Wallace, R.B., dan Wolinsky, F.D. (2007). Weight and Depressive Symptoms in Older Adults: Direction of Influence?. *Journal of Gerontology: Social Sciences*, 62B(1):S43–S51.
- Giannelli, F.R. (2020). Major depressive disorder. *Journal of the American Academy of Physician Assistants*, 33(4):19–20.
- Gluck, M.E., Geliebter, A., dan Satov, T. (2001). Night Eating Syndrome Is Associated with Depression, Low Self-Esteem, Reduced Daytime hunger, and Less Weight Loss in Obese Outpatients. *Obesity Research*, 9(4):264–267.

- Gulliford, M.C., Mahabir, D., dan Rocke, B. (2003). Food insecurity, food choices, and body mass index in adults: nutrition transition in Trinidad and Tobago. *International Journal of Epidemiology*, 32:508–516.
- He, X., Li, Z., Tang, X., Zhang, L., Wang, L., He, Y., Jin, T., dan Yuan, D. (2018). Age- and sex-related differences in body composition in healthy subjects aged 18 to 82 years. *Medicine*, 97(25):e11152.
- Hermann, S., Rohrmann, S., Linseisen, J., May, A.M., Kunst, A., Besson, H., Romaguera, D., Travier, N., Tormo, M-J., Molina, E., Dorronsoro, M., Barricarte, A., Rodríguez, L., Crowe, F.L., Khaw, K-T., Wareham, N.J., van Boeckel, P.G.A., Bueno-de-Mesquita, H.B., Overvad, K., Jakobsen, M.U., Tjønneland, A., Halkjær, J., agnoli, C., Mattiello, A., Tumino, R., Masala, G., Vineis, P., Naska, A., Orfanos, P., Trichopoulou, A., Kaaks, R., Bergmann, M.M., Steffen, A., Guelpen, B.V., Johansson, I., Borgquist, S., Manjer, J., Braaten, T., Fagherazzi, G., Clavel-Chapelon, F., Mouw, T., Norat, T., Riboli, E., Rinaldi, S., Slimani, N., dan Peeters, P.H.M. (2011). The association of education with body mass index and waist circumference in the EPIC-PANACEA study. *BMC Public Health*, 11:169.
- Hruby, A. dan Hu, F.B. (2015). The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics*, 33(7): 673-689.
- Insel, P., Ross, D., McMahon, K., dan Bernstein, M. (2017). *Nutrition Sixth Edition*. Burlington, Massachusetts: Jones & Bartlett Learning.
- Isaura, E.R., Chen, Y.-C., Adi, A.C., Fan, H.-Y., Li, C.-Y., dan Yang, S.-H. (2019). Association between Depressive Symptoms and Food Insecurity among Indonesian Adults: Results from the 2007–2014 Indonesia Family Life Survey. *Nutrients*, 11(3026).

- Ishtiaq, M., Afridi, M.I., Imranullah, Khan, S.A. (2018). Depression among adult population; prevalence & risk factors of depression among adult population of district Peshawar. *The Professional Medical Journal*, 25(8):1229–1234.
- Jahrami, H., Saif, Z., AlHaddad, M., Faris, M.A., Hammad, L., dan Ali, B. (2020). Assessing dietary and lifestyle risk behaviours and their associations with disease comorbidities among patients with depression: A case-control study from Bahrain. *Heliyon*, 6.
- Johnston, E., Johnson, S., McLeod, P., dan Johnston, M. (2004). The Relation of Body Mass Index to Depressive Symptoms. *Canadian Journal of Public Health*, 95(3): 179-183.
- Kaila, B. dan Raman, M. (2008). Obesity: A review of pathogenesis and management strategies, *Can J Gastroenterol*, 22(1): 61-68.
- Kaner, G., Soyulu, M., Yüksel, N., Inanç, N., Ongan, D., dan Başmisirli, E. (2015). Evaluation of Nutritional Status of Patients with Depression. *BioMed Research International*, 2015.
- Kementerian Kesehatan RI. (2018). *Hasil Utama Riset Kesehatan Dasar (Riskesdas) 2018*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Republik Indonesia.
- Kessler, R.C. dan Bromet, E.J. (2013). The epidemiology of depression across cultures. *Annu Rev Public Health*, 34: 119–138.
- Lakkis, N.A. dan Mahmassani, D.M. (2015). Screening instruments for depression in primary care: a concise review for clinicians. *Postgraduate Medicine*, 127(1):99–106.
- Lamers, F., Vogelzangs, N., Merikangas, K.R., de Jonge, P., Beekman, A.T.F., dan Penninx, B.W.J.H. (2013). Evidence for a differential role of HPA-axis

- function, inflammation and metabolic syndrome in melancholic versus atypical depression. *Molecular Psychiatry*, 18:692–699.
- Lazarevich, I., Camacho, M.E.I., Velázquez-Alva, M.d.C., dan Zepeda, M.Z. (2016). Relationship among obesity, depression, and emotional eating in young adults. *Appetite*, 107:639–644.
- Lee, J.-H., Park, S.K., Ryoo, J.-H., Oh, C.-M., Choi, R.S., McIntyre, R.S., Mansur, R.B., Kim, H., Hales, S., dan Jung, J.Y. (2017). U-shaped relationship between depression and body mass index in the Korean adults. *European Psychiatry*, 3531:1–9.
- Letamo, G. dan Navaneetham, K. (2014). Prevalence and Determinants of Adult Under-Nutrition in Botswana. *PLoS One*, 9(7).
- Lim, J.U., Lee, J.H., Kim, J.S., Hwang, Y.I., Kim, T.-H., Lim, S.Y., Yoo, K.H., Jung, K.-S., Kim, Y.K., Rhee, C.K. (2017). Comparison of World Health Organization and Asia-Pacific body mass index classifications in COPD patients, *International Journal of COPD*, 12: 2465–2475.
- Liu, C., Xie, B., Chou, C-P., Koprowski, C., Zhou, D., Palmer, P., Sun, P., Guo, Q., Duan, L., Sun, X., dan Johnson, C.A. (2007). Perceived stress, depression and food consumption frequency in the college students of China seven cities. *Physiology & Behavior*, 92:748–754.
- Liu, Q., He, H., Yang, J., Feng, X., Zhao, F., dan Lyu, J. (2020). Changes in the global burden of depression from 1990 to 2017: Findings from the Global Burden of Disease study. *Journal of Psychiatric Research*, 126: 134–140.
- Lorem, G.F., Schirmer, H., dan Emaus, N. (2017). What is the impact of underweight on self-reported health trajectories and mortality rates: a cohort study. *Health and Quality of Life Outcomes*, 15:191.

- Markowitz, S., Friedman, M.A., dan Arent, S.M. (2008). Understanding the Relation Between Obesity and Depression: Causal Mechanisms and Implications for Treatment. *Clinical Psychology: Science and Practice*, 15(1).
- McNeil, J.K. dan Harsany, M. (1989). An Age Difference View of Depression. *Canadian Psychology/Psychologie Canadienne*, 30(3):608–615.
- Mikolajczyk, R.T., Ansari, W.E., dan Maxwell, A.E. (2009). Food consumption frequency and perceived stress and depressive symptoms among students in three European countries. *Nutrition Journal*, 8(31).
- Milaneschi, Y., Simmons, W.K., van Rossum, E.F.C., dan Penninx, B.W.J.H. (2019). Depression and obesity: evidence of shared biological mechanisms. *Molecular Psychiatry*, 24:18 – 33.
- Milano, W., Ambrosio, P., Carizzone, F., De Biasio, v., Di Munzio, W., Foia, M.G., dan Capasso, A. (2020). Depression and Obesity: Analysis of Common Biomarkers. *Diseases*, 8(23).
- Mulrow, C.D., Williams Jr., J.W., Gerety, M.B., Ramirez, G., Montiel, O.M., dan Kerber, C. (1995). Case-finding instruments for depression in primary care settings. *Ann. Intern. Med*, 122: 913–921.
- Nonterah, E.A., Debpuur, C., Agongo, G., Amenga-Etego, L., Crowther, N.J., Ramsay, M., dan Oduro, A.R. (2018). Socio-demographic and behavioural determinants of body mass index among an adult population in rural Northern Ghana: the AWI-Gen study. *Global Health Action*, 11:68–79.
- Oliver, G. dan Wardle, J. (1999). Perceived Effects of Stress on Food Choice. *Physiology & Behavior*, 66(3):511–515.

- Oussaada, S.M., Galen, K.A.v., Cooman, M.I., Kleinendorst, L., Hazebroek, E.J., Haelst, M.M.v., Horst, K.W.t., dan Serlie, M.J. (2019). The pathogenesis of obesity, *Metabolism Clinical and Experimental*, 92: 26-36.
- Paans, N.P., Bot, M., Gibson-Smith, D., der Does, W.V., Spinhoven, P., Brouwer, I., Visser, m., Penninx, B.W.J.H. (2016). The association between personality traits, cognitive reactivity and body mass index is dependent on depressive and/or anxiety status. *Journal of Psychosomati Research*, 89(2016): 26-31.
- Par'i, H.M., Wiyono, S., dan Harjatmo, T.P. (2017). *Penilaian Status Gizi*. Jakarta: Pusat Pendidikan Sumber Daya Manusia Kesehatan Badan Pengembangan dan Pemberdayaan Sumber Daya Manusia Kesehatan.
- Pariante, C.M. dan Lightman, S.L. (2008). The HPA axis in major depression: classical theories and new developments. *Trends in Neurosciences*, 31(9).
- Peltzer, K. dan Pengpid, S. (2018). High prevalence of depressive symptoms in a national sample of adults in Indonesia: Child adversity, sociodemographic factors and health risk behaviour. *Asian Journal of Psychiatry*, 33: 52–59.
- Pengpid, S. dan Peltzer, K. (2017). The Prevalence of Underweight, Overweight/Obesity and Their Related Lifestyle Factors in Indonesia, 2014–2015. *AIMS Public Health*, 4(6):633–649.
- Piirtola, M., Jelenkovic, A., Latvala, A., Sund, R., Honda, C., Inui, F., Watanabe, M., Tomizawa, R., Iwatani, Y., Ordoñana, J.R., Sánchez-Romera, J.F., Colodro-Conde, L., Tamoki, A.D., Tarnoki, D.L., Martin, N.G., Montgomery, G.W., Medland, S.E., Rasmussen, F., Tynelius, P., Tan, Q., Zhang, D., Pang, Z., Rebato, E., Stazi, M.A., Fagnani, C., Brescianini, S., Busjahn, A., Harris, J.R., Brandt, I., Nilsen, T.S., Cutler, T.L., Hopper, J.L., Corley, R.P., Huibregtse, B.M., Sung, J., Kim, J., Lee, J., Lee, S., Gatz, M., Butler, D.A., Franz, C.E., Kremen,

- W.S., Lyons, M.J., Magnusson, P.K.E., Pedersen, N.L., Aslan, A.K.D., Öncel, S.Y., Aliev, F., Derom, C.A., Vlietinck, R.F., Loos, R.J.F., Silberg, J.L., Maes, H.H., Boomsma, D.I., Sørensen, T.I.A., Korhonen, T., Kaprio, J., dan Silventoinen, K. (2018). Association of current and former smoking with body mass index: A study of smoking discordant twin pairs from 21 twin cohorts. *PLoS One*, 13(7): e0200140. <https://doi.org/10.1371/journal.pone.0200140>
- Pine, D.S., Cohen, P., Brook, J., dan Coplan, J.D. (1997). Psychiatric Symptoms in Adolescence as Predictors of Obesity in Early Adulthood: A Longitudinal Study. *American Journal of Public Health*, 87:1303–1310.
- Qin, T., Liu, W., Yin, M., Shu, C., Yan, M., Zhang, J., dan Yin, P. (2017). Body mass index moderates the relationship between C-reactive protein and depressive symptoms: evidence from the China Health and Retirement Longitudinal Study. *Scientific Reports*, 7: 39940.
- Radloff, L.S. (1977). The CES-D Scale: A Self-Report Depression Scale for Research in the General Population. *Applied Psychological Measurement*, 1(3):385–401.
- Razzak, H.A., Harbi, A., dan Ahli, S. (2019). Depression: Prevalence and Associated Risk Factors in the United Arab Emirates. *Oman Medical Journal*, 34(4):274–283.
- Roberts, R.E., Deleger, S., Strawbridge, W.J., dan Kaplan, G.A. (2003). Prospective association between obesity and depression: evidence from the Alameda county study. *International Journal of Obesity*, 27:514–521.
- Romain, K., Webb, T., dan Kumar, M. (2018). Depression and obesity: can the cycle be broken? *BJPsych Advances*, 24:132–140.

- Seigel, K., Broman, J-E., dan Hetta, J. (2002). Behavioral activation or inhibition during emotional stress—implications for exercise habits and emotional problems among young females. *Nord J Psychiatry*, 56(6):441–446.
- Silva, D.A., Coutinho, E.d.S.F., de Figueiredo, R.C., Molina, M.d.C.B., Mill, J.G., de Faria, C.P., Nunes, M.A., Moreno, A.B., Giatti, L., Barreto, S.M., dan Viana, M.C. (2021). Depression is associated with increased adiposity in a 4-year follow-up: results from the ELSA-Brasil. *Journal of Affective Disorders*, 282:179–189.
- Simon, G.E., Ludman, E.J., Linde, J.A., Operskalski, B.H., Ichikawa, L., Rohde, P., Finch, E.A., dan Jeffery, R.W. (2008). Association between obesity and depression in middle-aged women. *General Hospital Psychiatry*, 30(2008):32–39.
- Stevens, J., Katz, E.G., dan Huxley, R.R. (2010). Associations between gender, age and waist circumference. *Eur J Clin Nutr*, 64(1):6–15.
- Strauss, J., Witoelar, F., dan Sikoki, B. (2016). *The Fifth Wave of the Indonesia Family Life Survey (IFLS5): Overview and Field Report*. WR-1143/1-NIA/NICHD.
- Stults-Kolehmainen, M.A. dan Sinha, R. (2014). The Effects of Stress on Physical Activity and Exercise. *Sports Med*, 44(1):81–121.
- Sudargo, T., Muhammad, H.F.L., Rosiyani, F., Kusmayanti, N.A. (2016). *Pola Makan dan Obesitas*. Yogyakarta: Gadjah Mada University Press.
- Supriasa, I.D.N., Bakri, B., dan Fajar, I. (2017). *Penilaian Status Gizi*. Edisi 2. Jakarta: EGC.
- Supriasa, I.D.N. (2017). Penilaian Status Gizi Secara Antropometri. Dalam: Hardinsyah, Supriasa, I.D.N., ed., *Ilmu Gizi: Teori & Aplikasi*, Edisi 1. Jakarta: EGC.

- Thau, L., Gandhi, J., dan Sharma, S. (2021). Physiology, Cortisol. Dalam: *StatPearls*. [Internet]. Treasure Island, Florida: StatPearls Publishing. Dapat diakses di: [https://www.ncbi.nlm.nih.gov/books/NBK538239/#\\_NBK538239\\_pubdet\\_](https://www.ncbi.nlm.nih.gov/books/NBK538239/#_NBK538239_pubdet_)
- Voinov, B., Richie, W.D., dan Bailey, R.K. (2013). Depression and Chronic Diseases: It Is Time for a Synergistic Mental Health and Primary Care Approach. *The Primary Care Companion for CNS Disorders*, 15(2).
- Wahyudi, B.F., Sriyono, dan Indarwati, R. (2015). Analisis Faktor yang Berkaitan dengan Kasus Gizi Buruk Pada Balita. *Jurnal Pedimaternal*, 3(1):83–91.
- WHO Expert Consultation. (2004). Appropriate body mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*, 363:157–163.
- World Health Organization. (2005). *WHO STEPS Surveillance Manual: The WHO STEPwise Approach to Chronic Disease Risk Factor Surveillance*. Geneva: World Health Organization.
- World Health Organization. (2008). *Waist Circumference and Waist-Hip Ratio: Report of a WHO Expert Consultation*. Geneva: World Health Organization.
- World Health Organization. (2017). *Depression and Other Common Mental Disorders: Global Health Estimates*. Geneva: World Health Organization. CC BY-NC-SA 3.0 IGO.