

## REFERENCES

- Acharya, B.N., Saraswat, D., Tiwari, M., Shrivastava, A.K., Ghorpade, R., Bapna, S., and Kaushik, M.P., 2010, Synthesis and antimalarial evaluation of 1, 3, 5-trisubstituted pyrazolines, *Eur. J. Med. Chem.*, 45 (2), 430–438.
- Albuquerque, H., Santos, C., Cavaleiro, J., and Silva, A., 2014, Chalcones as versatile synthons for the synthesis of 5- and 6-membered nitrogen heterocycles, *Curr. Org. Chem.*, 18 (21), 2750–2775.
- Antonio-Arias, J.E., DíazOoliva, V.C., Romero-Ceronio, N., Gómez-Rivera, A., Aguilar-Mariscal, H., de la Fuente, L.F.R., and Lobato-García, C.E., 2018, Monomodal vs multimodal microwave irradiation applied in the synthesis of fluorochalcones, *Am. J. Org. Chem.*, 8 (1), 8–12.
- Basco, L.K., Mitaku, S., Skaltsounis, A.L., Ravelomanantsoa, N., Tillequin, F., Koch, M., and Le Bras, J., 1994, *In vitro* activities of furoquinoline and acridone alkaloids against *Plasmodium falciparum*, *Antimicrob. Agents Chemother.*, 38 (5), 1169–1171.
- Batista, R., De Jesus Silva Júnior, A., and De Oliveira, A.B., 2009, Plant-derived antimalarial agents: New leads and efficient phytomedicines. part II. non-alkaloidal natural products, *Molecules*, 14 (8), 3037–3072.
- Bickii, J., Basco, L.K., and Ringwald, P., 1998, Assessment of three *in vitro* tests and an *in vivo* test for chloroquine resistance in *Plasmodium falciparum* clinical isolates, *J. Clin. Microbiol.*, 36 (1), 243–247.
- Bitew, H., Mammo, W., Hymete, A., and Yeshak, M.Y., 2017, Antimalarial activity of acetylenic thiophenes from *echinops hoehnelii schweinf*, *Molecules*, 22 (11), 1–10.
- Burrows, J.N., Hooft Van Huijsduijnen, R., Möhrle, J.J., Oeuvray, C., and Wells, T.N., 2013, Designing the next generation of medicines for malaria control and eradication, *Malar. J.*, 12 (1), 1–20.
- Chavan, B.B., Gadekar, A.S., Mehta, P.P., Vawhal, P.K., Kolsure, A.K., and A. R. Chabukswar, 2016, Synthesis and medicinal significance of chalcones - A Review, *Asian J. Biomed. Pharm. Sci.*, 6 (56), 1-7.
- Chinchilla, M., Valerio, I., Sánchez, R., Mora, V., Bagnarello, V., Martínez, L., Gonzalez, A., Vanegas, J.C., and Apestegui, Á., 2012, *In vitro* antimalarial activity of extracts of some plants from a biological reserve in Costa Rica, *Rev. Biol. Trop.*, 60 (2), 881–891.

- Dash, B., and Karim, S., 2021, Pyrazoline heterocyclic: A Review, *Int. J. Pharm. Sci. Res.*, 12 (5), 2570–2588.
- De Oliveira, M.E., Cenzi, G., Nunes, R.R., Andrighetti, C.R., De Sousa Valadao, D.M., Dos Reis, C., Simoes, C.M.O., Nunes, R.J., Júnior, M.C., Taranto, A.G., Sanchez, B.A.M., Viana, G.H.R., and De Pilla Varotti, F., 2013, Antimalarial activity of 4-metoxychalcones: Docking studies as falcipain/plasmeprin inhibitors, ADMET and lipophilic efficiency analysis to identify a putative oral lead candidate, *Molecules*, 18 (12), 15276–15287.
- Gaikwad, P.L., Gandhi, P.S., Jagdale, D.M., and Kadam, V.J., 2012, The use of biososterism in drug design and molecular modification, *Am. J. Pharmtech Res.*, 2 (4), 2249–3387.
- Gil, A., Pabón, A., Galiano, S., Burguete, A., Pérez-Silanes, S., Deharo, E., Monge, A., and Aldana, I., 2014, Synthesis, biological evaluation and structure-activity relationships of new quinoxaline derivatives as anti-Plasmodium falciparum agents, *Molecules*, 19 (2), 2166–2180.
- Hassan, S.Y., 2011, Synthesis and biological activity of some new pyrazoline and pyrimidine derivatives, *J. Braz. Chem. Soc.*, 22 (7), 1286–1298.
- Ipa, M., Widawati, M., Laksono, A.D., Kusri, I., and Dhewantara, P.W., 2020, Variation of preventive practices and its association with malaria infection in eastern Indonesia: Findings from community-based survey, *PLoS One*, 15 (5), 1–18.
- Jasril, J., Teruna, H.Y., Aisyah, A., Nurlaili, N., and Hendra, R., 2019, Microwave assisted synthesis and evaluation of toxicity and antioxidant activity of pyrazoline derivatives, *Indones. J. Chem.*, 19 (3), 583–591.
- Kumar, G., Tanwar, O., Kumar, J., Akhter, M., Sharma, S., Pillai, C.R., Alam, M.M., and Zama, M.S., 2018, Pyrazole-pyrazoline as promising novel antimalarial agents: A mechanistic study, *Eur. J. Med. Chem.*, 149 139–147.
- Kumar, R., Mohanakrishnan, D., Sharma, A., Kaushik, N.K., Kalia, K., Sinha, A.K., and Sahal, D., 2010, Reinvestigation of structure-activity relationship of methoxylated chalcones as antimalarials: Synthesis and evaluation of 2,4,5-trimethoxy substituted patterns as lead candidates derived from abundantly available natural  $\beta$ -asarone, *Eur. J. Med. Chem.*, 45 (11), 5292–5301.
- Lohidashan, K., Rajan, M., Ganesh, A., Paul, M., and Jerin, J., 2018, Pass and Swiss ADME collaborated in silico docking approach to the synthesis of certain

pyrazoline spacer compounds for dihydrofolate reductase inhibition and antimalarial activity, *Bangladesh J. Pharmacol.*, 13 (1), 23–29.

Marella, A., Shaquiquzzaman, M., Akhter, M., Verma, G., and Alam, M.M., 2015, Novel pyrazole-pyrazoline hybrids endowed with thioamide as antimalarial agents: Their synthesis and 3D-QSAR studies, *J. Enzyme Inhib. Med. Chem.*, 30 (4), 597–606.

Mishra, V.K., Mishra, M., Kashaw, V., and Kashaw, S.K., 2017, Synthesis of 1,3,5-trisubstituted pyrazolines as potential antimalarial and antimicrobial agents, *Bioorganic Med. Chem.*, 25 (6), 1949–1962.

Muriithi, M.W., Abraham, W., Addae-kyereme, J., Scowen, I., Croft, S.L., Gitu, P.M., Kendrick, H., Njagi, E.N.M., and Wright, C.W., 2002, Isolation and *in vitro* antiplasmodial activities of alkaloids from *Teclea trichocarpa*: *in vivo* antimalarial activity and x-ray crystal structure of normelicopicine, *J. Nat. Prod.*, 2 (65), 956–959.

Nandiyanto, A.B.D., Oktiani, R., and Ragadhita, R., 2019, How to read and interpret ftr spectroscopy of organic material, *Indones. J. Sci. Technol.*, 4 (1), 97–118.

Nevozhay, D., 2014, Cheburator software for automatically calculating drug inhibitory concentrations from *in vitro* screening assays, *PLoS One*, 9 (9), 1–10.

Nyandwaro, K., Oyweri, J., Kimani, F., and Mbugua, A., 2020, Evaluating antiplasmodial and antimalarial activities of soybean (glycine max) seed extracts on *P. falciparum* parasite cultures and *P. berghei* - infected mice, *J. Pathog.*, 2020, 1–8.

Okolo, C., Eban, L., Amazu, L., Chukwu, L., Ohadoma, S., and Osuala, F., 2020, *In-vitro* anti-malarial activity of Chikadoma plant from the rainforest of Southern Nigeria, *J. Drug Deliv. Ther.*, 10 (5), 251–254.

Omodior, O., Luetke, M.C., and Nelson, E.J., 2018, Mosquito-borne infectious disease, risk-perceptions, and personal protective behavior among U.S. international travelers, *Prev. Med. Reports*, 12 (July), 336–342.

Powers, D.G., Casebier, D.S., Fokas, D., Ryan, W.J., Troth, J.R., and Coffen, D.L., 1998, Automated parallel synthesis of chalcone-based screening libraries, *Tetrahedron*, 54 (16), 4085–4096.

Prudêncio, M., 2020, Letter in fairness to mosquitoes, *Trends Parasitol.*, 36 (11), 876–877.

- Rani, M., and Mohamad, Y., 2014, Synthesis, studies and *in vitro* antibacterial activity of some 5-(thiophene-2-yl)-phenyl pyrazoline derivatives, *J. Saudi Chem. Soc.*, 18 (5), 411–417.
- Rocha e Silva, L.F., de Magalhães, P.M., Costa, M.R.F., Alecrim, M. das G.C., Chaves, F.C.M., Hidalgo, A. de F., Pohlit, A.M., and Vieira, P.P.R., 2012, *In vitro* susceptibility of *Plasmodium falciparum* Welch field isolates to infusions prepared from *Artemisia annua* L. cultivated in the Brazilian Amazon, *Mem. Inst. Oswaldo Cruz*, 107 (7), 859–866.
- Santalova, E.A., and Denisenko, V.A., 2017, Analysis of the configuration of an isolated double bond in some lipids by selective homonuclear decoupling, *Nat. Prod. Commun.*, 12 (12), 1913–1916.
- Sethiya, J.P., Bhavsar, S.P., and Shahare, H. V, 2019, Synthesis of potential anti-inflammatory pyrazoline derivatives under ultrasonic irradiation, *Int. J. Pharm. Sci. Res.*, 10 (7), 3290–3294.
- Shibeshi, M.A., Kifle, Z.D., and Atnafie, S.A., 2020, Antimalarial drug resistance and novel targets for antimalarial drug discovery, *Infect. Drug Resist.*, 13 4047–4060.
- Syahri, J., Yuanita, E., Nurohmah, B.A., Armunanto, R., and Purwono, B., 2017, Chalcone analogue as potent anti-malarial compounds against *Plasmodium falciparum*: Synthesis, biological evaluation, and docking simulation study, *Asian Pac. J. Trop. Biomed.*, 7 (8), 675–679.
- Talapko, J., Škrlec, I., Alebić, T., Jukić, M., and Včev, A., 2019, Malaria: The past and the present, *Microorganisms*, 7 (6), 1-17.
- Tanner, M., Greenwood, B., Whitty, C.J.M., Ansah, E.K., Price, R.N., Dondorp, A.M., von Seidlein, L., Baird, J.K., Beeson, J.G., Fowkes, F.J.I., Hemingway, J., Marsh, K., and Osier, F., 2015, Malaria eradication and elimination: Views on how to translate a vision into reality, *BMC Med.*, 13 (1), 1–22.
- Widyawaruyanti, A., Devi, A.P., Fatria, N., Tumewu, L., Tantular, I.S., and Hafid, A.F., 2014, *In vitro* antimalarial activity screening of several Indonesian plants using HRP2 assay, *Int. J. Pharm. Pharm. Sci.*, 6 (6), 125–128.
- Yadav, N., Dixit, S.K., Bhattacharya, A., Mishra, L.C., Sharma, M., Awasthi, S.K., and Bhasin, V.K., 2012, Antimalarial activity of newly synthesized chalcone derivatives *in vitro*, *Chem. Biol. Drug Des.*, 80 (2), 340–347.