

## DAFTAR PUSTAKA

- A, A. Q., & N, A. Q. (2017). Prevention of Depression: A Review of Literature. *Journal of Depression & Anxiety*, 06(04), 6–11. <https://doi.org/10.4172/2167-1044.1000292>
- Abhishek, G., Uday, M., Singh Shivendra, K., Manish, M. K., Chandra, T. S., & Kumar, S. V. (2015). Screening depression among elderly in a city of southeast Asia. *Journal of Clinical and Diagnostic Research*, 9(9), 3–7. <https://doi.org/10.7860/JCDR/2015/14100.6426>
- Ahmadi, B., Amini Sanii, N., Allahverdi pour, H., Bani, F., & Bakhtari, F. (2018). Predictors of Physical Activity in Older Adults in Northwest of Iran. *Elderly Health Journal*, 4(2), 75–80. <https://doi.org/10.18502/ehj.v4i2.260>
- Andresen, E. M., Malmgren, J. A., Carter, W. B., & Patrick, D. L. (1994). Screening for depression in well older adults: evaluation of a short form of the CES-D (Center for Epidemiologic Studies Depression Scale). *American journal of preventive medicine*, 10(2), 77–84.
- Albert, P. R. (2015). Why is depression more prevalent in women? *Journal of Psychiatry and Neuroscience*, 40(4), 219–221. <https://doi.org/10.1503/jpn.150205>
- Arai, T., Obuchi, S., Kojima, M., Nishizawa, S., Matsumoto, Y., & Inaba, Y. (2009). The relationship between age and change in physical functions after exercise intervention. Trainability of Japanese community-dwelling older elderly. *Journal of the Japanese Physical Therapy Association*, 12(1), 1–8. <https://doi.org/10.1298/jjpta.12.1>
- Aritonang, J. M. P., Soewadi, S., & Wirasto, R. T. (2018). Korelasi Tingkat Kebermaknaan Hidup Dengan Depresi Pada Lansia Di Posyandu Lansia Padukuhan Soro Padan, Sleman, Yogyakarta. *Berkala Ilmiah Kedokteran Duta Wacana*, 3(1), 25. <https://doi.org/10.21460/bikdw.v3i1.77>
- Australian Longitudinal Study on Women's Health. (2020). ALSWH Data Dictionary Supplement. Global Recommendations On Physical Activity For Health. <https://alswh.org.au/wp-content/uploads/2020/08/DDSSection2.7CESD.pdf>
- Badan Pusat Statistik (BPS). (2019). Statistik Penduduk Lanjut Usia. *Katalog: 4104001*. <https://www.bps.go.id/publication/2019/12/20/ab17e75dbe630e05110ae53b/statistik-penduduk-lanjut-usia-2019.html>
- Badan Pusat Statistik (BPS). (2020). Statistik Penduduk Lanjut Usia. *Katalog: 4104001*. <https://www.bps.go.id/publication/2020/12/21/0fc023221965624a644c1111/statistik-penduduk-lanjut-usia-2020.html>
- Baron, E. C., Davies, T., & Lund, C. (2017). Validation of the 10-item Centre for Epidemiological Studies Depression Scale (CES-D-10) in Zulu, Xhosa and Afrikaans populations in South Africa. *BMC Psychiatry*, 17(1), 1–14. <https://doi.org/10.1186/s12888-016-1178-x>
- Bartha, C., Parker, C., Thomson, C., & Kitchen, K. (2013). *Depression: an Information Guide*.

- Bembnowska, M., & Joško-Ochojska, J. (2015). What causes depression in adults? *Polish Journal of Public Health*, 125(2), 116–120. <https://doi.org/10.1515/pjph-2015-0037>
- Berkowitz, R. I., & Fabricatore, A. N. (2005). Obesity, psychiatric status, and psychiatric medications. *Psychiatric Clinics of North America*, 28(1), 39–54. <https://doi.org/10.1016/j.psc.2004.10.005>
- Boschloo, L., Reeuwijk, K. G., Schoevers, R. A., & W J H Penninx, B. (2014). The impact of lifestyle factors on the 2-year course of depressive and/or anxiety disorders. *Journal of affective disorders*, 159, 73–79. <https://doi.org/10.1016/j.jad.2014.01.019>
- Branco, J. C., Jansen, K., Sobrinho, J. T., Carrapatoso, S., Spessato, B., Carvalho, J., Mota, J., & da Silva, R. A. (2015). Benefícios físicos e redução de sintomas depressivos em idosos: Resultados do Programa Nacional de Caminhada Português. *Ciencia e Saude Coletiva*, 20(3), 789–796. <https://doi.org/10.1590/1413-81232015203.09882014>
- Byeon, H. (2019). Relationship between physical activity level and depression of elderly people living alone. *International Journal of Environmental Research and Public Health*, 16(20). <https://doi.org/10.3390/ijerph16204051>
- Campbell, Collen Marie. (2012). Methods of physical activity assessment for older adults. *Graduate Theses and Dissertation*. 12809. <https://lib.dr.iastate.edu/etd/12809>
- Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public health reports (Washington, D.C. : 1974)*, 100(2), 126–131.
- Centre for Clinical Interventions. (2018). Depression Information Sheet - 02 - What Causes Depression. *Centre for Clinical Interventions*.
- Centers for Disease Control and Prevention.(1999). Physical Activity and Health Older Adults. A Report of the Surgeon General. <https://www.cdc.gov/nccdphp/sgr/pdf/olderad.pdf>
- Chang, Y. C., Lu, M. C., Hu, I. H., Wu, W. C. I., & Hu, S. C. (2017). Effects of different amounts of exercise on preventing depressive symptoms in community-dwelling older adults: A prospective cohort study in Taiwan. *BMJ Open*, 7(4), 1–10. <https://doi.org/10.1136/bmjopen-2016-014256>
- Cole, M. G., & Dendukuri, N. (2003). Risk factors for depression among elderly community subjects: A systematic review and meta-analysis. *American Journal of Psychiatry*, 160(6), 1147–1156. <https://doi.org/10.1176/appi.ajp.160.6.1147>
- Craft, L. L., & Perna, F. M. (2004). The Benefits of Exercise for the Clinically Depressed. Primary care companion to the Journal of clinical psychiatry, 6(3), 104–111. <https://doi.org/10.4088/pcc.v06n0301>
- Cruz-Jentoft, A. J., Baeyens, J. P., Bauer, J. M., Boirie, Y., Cederholm, T., Landi, F., ... Zamboni, M. (2010). Sarcopenia: European consensus on definition and diagnosis. *Age and Ageing*, 39(4), 412–423. <https://doi.org/10.1093/ageing/afq034>
- Daud, K. A. M., Khidzir, N. Z., Ahmad Rasdan, I., & Fadhilahanim Aryani, A. (2018). Validity and reliability of instrument to measure social media skills

- among small and medium entrepreneurs at Pengkalan Datu River. *International Journal of Development and Sustainability*, 7(3), 1026–1037. [www.isdsnet.com/ijds](http://www.isdsnet.com/ijds)
- Darton, Katherine. (2012). *Understanding Depression*. London: Mind. [https://www.mind.org.uk/media/42904/understanding\\_depression\\_2012.pdf](https://www.mind.org.uk/media/42904/understanding_depression_2012.pdf)
- Dunlop, D. D., Song, J., Semanik, P. A., Sharma, L., & Chang, R. W. (2011). Physical activity levels and functional performance in the osteoarthritis initiative: A graded relationship. *Arthritis and Rheumatism*, 63(1), 127–136. <https://doi.org/10.1002/art.27760>
- IFLS. (2014). Kuesioner ComFas IFLS5.
- Irawan, H. (2013). Gangguan Depresi pada Lanjut Usia. *Cermin Dunia Kedokteran*-210, 40(11), 815–819. [http://www.kalbemed.com/Portals/6/06\\_210Gangguan\\_Depresi\\_pada\\_Lanjut\\_Usia.pdf](http://www.kalbemed.com/Portals/6/06_210Gangguan_Depresi_pada_Lanjut_Usia.pdf)
- Irwin, M., Artin, K. H., & Oxman, M. N. (1999). Screening for Depression in the Older Adult. *Archives of Internal Medicine*, 159(15), 1701. <https://doi.org/10.1001/archinte.159.15.1701>
- Jongeneelis, K., Pot, A. M., Eisses, A. M. H., Beekman, A. T. F., Kluiter, H., & Ribbe, M. W. (2004). Prevalence and risk indicators of depression in elderly nursing home patients: The AGED study. *Journal of Affective Disorders*, 83(2–3), 135–142. <https://doi.org/10.1016/j.jad.2004.06.001>
- Kepres RI. (1998). *UU NO 13 TAHUN 1998 tentang Kesejahteraan Lansia*. I(disitasi 2019 Maret 4), 3. <http://bphn.go.id>
- Kemenkes, RI. (2016). Infodatin Situasi Lanjut Usia (Lansia) di Indonesia. Jakarta: Pusat data dan informasi Kemenkes RI. Tersedia di: <https://pusdatin.kemkes.go.id/resources/download/pusdatin/infodatin/Infodatin-lansia-2016.pdf>
- Kementrian Kesehatan RI.(2016). Kurikulum Pelatihan Bagi Pelatih Pelayanan Kesehatan Lanjut Usia dan Geriatri untuk Petugas Puskesmas. Jakarta:Bakti Husada.Tersedia di: [http://siakpel.bppsdmk.kemkes.go.id:8102/akreditasi\\_kurikulum/kurikulum\\_1911131042097162625d1586b1fdccc7fe400aae45af.pdf](http://siakpel.bppsdmk.kemkes.go.id:8102/akreditasi_kurikulum/kurikulum_1911131042097162625d1586b1fdccc7fe400aae45af.pdf)
- Kemenkes, RI.(2013). RISKESDAS. Jakarta: Indonesia. Tersedia di: [https://labmandat.litbang.kemkes.go.id/images/download/laporan/RKD/2013/Laporan\\_riskesdas\\_2013\\_final.pdf](https://labmandat.litbang.kemkes.go.id/images/download/laporan/RKD/2013/Laporan_riskesdas_2013_final.pdf)
- Kurdi, F. N., & Flora, R. (2019). Physical exercise increased brain-derived neurotrophic factor in elderly population with depression. *Open Access Macedonian Journal of Medical Sciences*, 7(13), 2057–2061. <https://doi.org/10.3889/oamjms.2019.574>
- Kurlowicz, L. H. (1993). Social factors and depression in late life. *Archives of Psychiatric Nursing*, 7(1), 30–36. [https://doi.org/10.1016/0883-9417\(93\)90020-W](https://doi.org/10.1016/0883-9417(93)90020-W)
- Kurniawan, B. (2013). Hubungan Antara Aktivitas Fisik Dengan Tingkat Depresi Pada Lansia Di Panti Werdha Budi Luhur, Bantul.*Skripsi*.
- Madyaningrum, E., Chuang, Y. C., & Chuang, K. Y. (2019). Prevalence and related factors of depression among the elderly in Indonesia. *International*

- Journal of Gerontology*, 13(3), 202–206.  
[https://doi.org/10.6890/IJGE.201909\\_13\(3\).0004](https://doi.org/10.6890/IJGE.201909_13(3).0004)
- Mao, H. Y., Hsu, H. C., & Lee, S. Da. (2020). Gender differences in related influential factors of regular exercise behavior among people in Taiwan in 2007: A cross-sectional study. *PLoS ONE*, 15(1), 1–13.  
<https://doi.org/10.1371/journal.pone.0228191>
- McAuley, E., Mihalko, S. L., & Bane, S. M. (1997). Exercise and self-esteem in middle-aged adults: Multidimensional relationships and physical fitness and self-efficacy influences. *Journal of Behavioral Medicine*, 20(1), 67–83.  
<https://doi.org/10.1023/A:1025591214100>
- Musich, S., Wang, S. S., Hawkins, K., & Greame, C. (2017). The Frequency and Health Benefits of Physical Activity for Older Adults. *Population Health Management*, 20(3), 199–207. <https://doi.org/10.1089/pop.2016.0071>
- National Collaborating Centre for Mental Health (UK). Depression: The Treatment and Management of Depression in Adults (Updated Edition). Leicester (UK): British Psychological Society; 2010. (NICE Clinical Guidelines, No. 90.) Available from:  
<https://www.ncbi.nlm.nih.gov/books/NBK63748/>
- National Institute of Mental Health. (2016). Depression Basics. *National Institutes of Health*, 6. <https://doi.org/10.1039/c3cc46800c>
- Nazzari, H., Isserow, S. H., Heilbron, B., Chb, M. B., McKinney, J., Lithwick, D. J., Morrison, B. N., Nazzari, H., Isserow, S. H., Heilbron, B., & Krahm, A. D. (2016). The health benefits of physical activity and cardiorespiratory fitness. *British Columbia Medical Journal*, 58(3), 131–137.
- New England Research Institute. (1991). *Physical Activity Scale for the Elderly (PASE) Administration and Scoring Instruction Manual*. [https://www.physio-pedia.com/Physical Activity Scale for the Elderly \(PASE\)](https://www.physio-pedia.com/Physical_Activity_Scale_for_the_Elderly_(PASE)).
- Nurullah, F. E., Nuripah, G., & Dewi, M. K. (2015). Hubungan Olahraga Rutin dengan Tingkat Depresi pada Lansia di Kecamatan Coblong Kota Bandung. *Prosiding Pendidikan Dokter*.  
<http://karyailmiah.unisba.ac.id/index.php/dokter/article/download/1386/pdf>
- Park, C.-H., Elavsky, S., & Koo, K.-M. (2014). Factors influencing physical activity in older adults. *Journal of Exercise Rehabilitation*, 10(1), 45–52.  
<https://doi.org/10.12965/jer.140089>
- Pettee, K. K., Brach, J. S., Kriska, A. M., Boudreau, R., Richardson, C. R., Colbert, L. H., Satterfield, S., Visser, M., Harris, T. B., Ayonayon, H. N., & Newman, A. B. (2006). Influence of marital status on physical activity levels among older adults. *Medicine and Science in Sports and Exercise*, 38(3), 541–546. <https://doi.org/10.1249/01.mss.0000191346.95244.f7>
- Qamar, M., Zahirah, N., & Ahmad, S. (2020). Assessment of Depression Among Older Adults Staying in Old Home Care in Klang, Malaysia. *Pharmaceutical Sciences and Research*, 7(1), 66–74. <https://doi.org/10.7454/psr.v7i1.1062>
- Primasari, Ameta. (2018). Proses Penuaan dari Aspek Kedokteran Gigi / Ameta Primasari – Edisi Ke-2 -- Medan: USU Press 2018.  
<http://repository.usu.ac.id/handle/123456789/70769>

- Roadolff, L. S. (1977). The CES-D scale: a self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1, 385–401.
- Rosyanti, L. Hadi, I. F. (2018). *Memahami Gangguan Depresi Mayor (Major Depression Disorder)*. 10–15.
- Sharma, A. M., & Padwal, R. (2010). Obesity is a sign - Over-eating is a symptom: An aetiological framework for the assessment and management of obesity. *Obesity Reviews*, 11(5), 362–370. <https://doi.org/10.1111/j.1467-789X.2009.00689.x>
- Shaw, B. A., & Spokane, L. S. (2008). Examining the association between education level and physical activity changes during early old age. *Journal of Aging and Health*, 20(7), 767–787. <https://doi.org/10.1177/0898264308321081>
- Sherwood, N. E., & Jeffery, R. W. (2000). The behavioral determinants of exercise: implications for physical activity interventions. *Annual review of nutrition*, 20, 21–44. <https://doi.org/10.1146/annurev.nutr.20.1.21>
- Singh, D. K. A., Rahman, N. N. A. A., Rajaratnam, B. S., Yi, T. C., & Shahar, S. (2018). Validity and reliability of physical activity scale for elderly in Malay language (PASE-M). *Malaysian Journal of Public Health Medicine*, 2018(Specialissue1), 116–123.
- Stahl, S. M. (2002). The psychopharmacology of painful physical symptoms in depression. *Journal of Clinical Psychiatry*, 63(5), 382–383. <https://doi.org/10.4088/JCP.v63n0501>
- Sumantri, A. (2011). *Metodologi Penelitian Kesehatan*. Jakarta:Kencana PrenadaMedia Group.
- Thivel, D., Tremblay, A., Genin, P. M., Panahi, S., Rivière, D., & Duclos, M. (2018). Physical Activity, Inactivity, and Sedentary Behaviors: Definitions and Implications in Occupational Health. *Frontiers in Public Health*, 6(October), 1–5. <https://doi.org/10.3389/fpubh.2018.00288>
- The National Institute on Aging. (2020). Strategic Directions for Research 2020-2025. <https://www.nia.nih.gov/sites/default/files/2020-05/nia-strategic-directions-2020-2025.pdf>
- UN DESA's Population Division. (2019). World Population Ageing 2019. In *United Nations*. <https://www.un.org/development/desa/en/news/population/our-world-is-growing-older.html>
- U.S. Department of Health and Human Services. (2018). Physical Activity Guidelines for Americans, 2<sup>nd</sup> edition. *Department of Health and Human Services. Washington, DC: U.S*
- Van Damme, A., Declercq, T., Lemey, L., Tandt, H., & Petrovic, M. (2018). Late-life depression: Issues for the general practitioner. *International Journal of General Medicine*, 11, 113–120. <https://doi.org/10.2147/IJGM.S154876>
- Varma, G. S. (2012). Depression in the Elderly: Clinical Features and Risk Factors. *Aging and Disease*, 3(6), 465–471.
- Wang, S., Ma, W., Wang, S. M., & Yi, X. (2018). A cross sectional examination of the relation between depression and frequency of leisure time physical



- exercise among the elderly in Jinan, China. *International Journal of Environmental Research and Public Health*, 15(9). <https://doi.org/10.3390/ijerph15092041>
- Wang, R., Bishwajit, G., Zhou, Y., Wu, X., Feng, D., Tang, S., Chen, Z., Shaw, I., Wu, T., Song, H., Fu, Q., & Feng, Z. (2019). Intensity, frequency, duration, and volume of physical activity and its association with risk of depression in middle- And older-aged Chinese: Evidence from the China Health and Retirement longitudinal study, 2015. *PLoS ONE*, 14(8), 1–17. <https://doi.org/10.1371/journal.pone.0221430>
- Washburn, R. A., McAuley, E., Katula, J., Mihalko, S. L., & Boileau, R. A. (1999). The Physical Activity Scale for the Elderly (PASE): Evidence for validity. *Journal of Clinical Epidemiology*, 52(7), 643–651. [https://doi.org/10.1016/S0895-4356\(99\)00049-9](https://doi.org/10.1016/S0895-4356(99)00049-9)
- Watson, K. B., Carlson, S. A., Gunn, J. P., Galuska, D. A., O'Connor, A., Greenlund, K. J., & Fulton, J. E. (2016). Physical Inactivity Among Adults Aged 50 Years and Older - United States, 2014. *MMWR. Morbidity and mortality weekly report*, 65(36), 954–958. <https://doi.org/10.15585/mmwr.mm6536a3>
- Weeks, L. E., Profit, S., Campbell, B., Graham, H., Chircop, A., & Sheppard-LeMoine, D. (2008). Participation in physical activity: Influences reported by seniors in the community and in long-term care facilities. *Journal of Gerontological Nursing*, 34(7), 36–43. <https://doi.org/10.3928/00989134-20080701-11>
- WHO. (2020). Depression. Diakses 4 Januari 2021 dari <https://www.who.int/news-room/fact-sheets/detail/depression>
- WHO. (2017). Depression and Other Common Mental Disorders: Global Health Estimates. Geneva: World Health Organization. <https://apps.who.int/iris/bitstream/handle/10665/254610/WHO-MSD-MER-2017.2-eng.pdf?sequence=1>
- WHO. (2011). Global Recommendations On Physical Activity For Health. <https://www.who.int/dietphysicalactivity/leaflet-physical-activity-recommendations.pdf>
- WHO. (2016). Preventing depression in the WHO European Region. *World Health Organization*. [https://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0003/325947/New-Preventing-depression.pdf](https://www.euro.who.int/__data/assets/pdf_file/0003/325947/New-Preventing-depression.pdf)
- WHO. (2002). Proposed working definition of an older person in Africa for the MDS Project. <https://www.who.int/healthinfo/survey/ageingdefnolder/en/>
- Yusuf, Ah., PK, Rizky Fitryasari., & Nihayati, Hanik Endang. (2014). *Buku Ajar Keperawatan Kesehatan Jiwa*. Salemba Medika.
- Zhang, W., O'Brien, N., Forrest, J. I., Salters, K. A., Patterson, T. L., Montaner, J. S. G., Hogg, R. S., & Lima, V. D. (2012). Validating a shortened depression scale (10 item CES-D) among HIV-Positive people in British Columbia, Canada. *PLoS ONE*, 7(7). <https://doi.org/10.1371/journal.pone.0040793>